Published in final edited form as:

Tetrahedron Lett. 2008 March 31; 49(14): 2316–2319.

Synthesis of a novel benzyl-octahydropyrazino[1,2-a]pyrimidin-6-one derivative as a convenient internal bicyclic peptidomimetic

Byoung J. Min, Xuyuan Gu, Takashi Yamamoto, Ravil R. Petrov, Hongchang Qu, Yeon Sun Lee, and Victor J. Hruby*

Department of Chemistry, University of Arizona, Tucson, AZ 85721, USA.

Abstract

A substituted hydropyrazino[1,2-a]pyrimidin-6-one derivative was synthesized stereoselectively *via* the intramolecular N-acyliminium ion cyclization between an amide nitrogen and an N^α-acetal derived from Cbz-protected aminopropyl-phenylalaninamide in very good yields. The formation of a single diastereomer is due to the low energy chairlike conformation of its bicyclic structure. This methodology provides a convenient tool to build internal bicyclic peptidomimetics.

Keywords

internal bicyclic peptidomimetics; N-acyliminium ion cyclization; acetals

It has been well recognized that β -turns are important structural features in biologically active peptides and proteins in terms of their function.¹ A large amount of research has been pursued to investigate peptide-protein or protein-protein interactions that also could be induced by small molecules bearing similar local structural features.² Recently, our group has developed the external β -turn mimetics **1** (Figure 1) containing the thiazolo[3,2-a]pyridine-5-one moiety *via* both solution and solid phase synthesis.³ It has been known that for interactions with a target protein, side chains of internal β -turn mimetics are more accessible than those of external mimetics, which are more sterically hindered by the support.⁴ Therefore, we have designed and synthesized a novel internal bicyclic scaffold for peptidomimetic **2** (Figure 1). For structure characterization and configuration assignment, a density functional theory (DFT) calculation, solution NMR spectroscopy, and X-ray crystallography were performed.

Compound **8** was designed as a model compound to investigate the synthesis of 1,3,6,8-substituted tetrahydro-2*H*-pyrazino[1,2-a]pyrimidine-4,7-diones (**2**) derivatives (Scheme 1). The straight forward approach to this bicyclic scaffold was to introduce an acetal moiety at the α -amino group of the phenylalanine, which can be attacked first by the amide to yield the acyliminium ion **7** and then by the carbamate nitrogen atoms in a one-pot fashion to construct the bicyclic structure. The reaction was expected to give a corresponding diastereomeric mixture, as the nucleophilic attack may take place from both the *si*-face and *re*-face of the planar iminium ion double bond.^{4,5} Surprisingly, a single diastereomer was generated and isolated from the reaction, and its purity was confirmed by HPLC. Here we provide details for the synthesis, the rationalization of diastereoselectivity, and its absolute stereochemistry.

*Corresponding Author. Phone: 520-621-6332; Fax: 520-621-8407; email: hruby@u.arizona.edu

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

(3-Aminopropyl)-carbamic acid benzyl ester **3** was conjugated with N^{α} -Boc-*L*-Phe-OH using BOP, HOBt, and NMM to afford the amide **4** in 92 % isolated yield (Scheme 1).⁶ After deprotection of the N^{α} -Boc group by TFA, reductive amination was performed by sequential additions of 2,2-dimethoxyacetaldehyde and sodium triacetoxyborohydride in the presence of a catalytic amount of acetic acid in THF.⁷ This one-pot 3-step reaction generated the secondary amine **5** with an acetal moiety in 90% yield after purification. The Cbz group was chosen to protect the secondary amine, though other protection groups could be used for the orthogonal protection. After protection, the corresponding acetal **6** was subjected to acid-catalyzed cyclization in the presence of formic acid to give bicyclic product **8** in 89% yield. Interestingly, this bicyclic product **8** only showed a single diastereomer both in crude NMR and HPLC analysis. In order to confirm the stereochemistry and the optical purity of the diastereomer, various NMR analyses were performed, but the result was not conclusive due to the existence of rotamers. Thus, the two Cbz groups of **8** were removed by hydrogenation. The NMR spectrum as well as HPLC analysis of the product **9** clearly indicated that the compound **9** was a single diastereomer.⁸ The product was crystallized in a mixed solvent CH_2Cl_2 , EtOAc, and MeOH. The X-ray crystal structure of **9** confirmed that the bridged head hydrogen has a *cis*-configuration to the hydrogen at position 7, and that the pyrimidine and piperazine rings have a chairlike conformation with the benzyl group in an axial position (Figure 2).⁹ In the structural conformation of **9**, the anomeric effect could be one possible factor for its chairlike structure, since the lone pair of the nitrogen stabilizes the axial benzyl group in the chairlike conformation more than the equatorial one in the boatlike conformation.¹⁰

To confirm their configurations, density functional theory (DFT) calculations were performed using the Gaussian 03 program.¹¹ Structure optimization followed by a frequency calculation was performed on **9** at the B3LYP/6-31G(d) level (no imaginary frequencies were found). Zero-point energies (ZPEs) of the DFT results were scaled by 0.9806.¹² The compound **9** was studied to compare the obtained calculated conformation with its X-ray crystal structure. It turns out that the calculated structure of **9** was shown to be well-overlapped with the X-ray crystal structure. For simplicity with low calculational costs, the model compounds of two diastereomers, in which a Cbz group at the position 1 in **9** was replaced by an acetyl group, were calculated by the same method.

As anticipated, the diastereomer[y2] **10**, whose pyrimidine ring has a chairlike conformation, is more stable than the diastereomer **11** which has a twist boatlike conformation implying that the bis-acylated benzyl-octahydropyrazino[1,2-*a*]pyrimidin-6-one derivative also prefers a chair-like conformation rather than a boatlike one (Figure 3). The energy difference between the two diastereomers is 1.5 kcal/mol, which leads to a 9/1 product ratio. On the basis of the results from the DFT calculation and X-ray crystal structure, the thermodynamically preferred diastereomer was confirmed.

An interesting result was found when the synthesis of the 6,5-membered ring **19** (Scheme 2) was attempted. The 6,5-bicyclic compound **19** could not be generated by the same strategy (Scheme 2). The acetal **15** did not go on to cyclization even with reaction times of up to 5 days and heating for 4 h, but instead underwent β -elimination to generate enamide product **17** in 10 % yield which was the only isolated product along with the starting material.¹³ We [y3] speculate that the elimination reaction is the major side reaction for this type of tandem bicyclization. Thermodynamically driven 6,6-membered bicyclic formation could be a stable pathway to yield the desired product while the kinetically driven 6,5-membered bicyclic needs higher energy compared to the eliminated enamide product **17**. Therefore, the compound **17** was the only isolated product under the harsh reaction condition. Other ring-size bicyclic constructions currently are under investigation.

In summary, we have developed a simple and efficient methodology to synthesize a substituted octahydropyrazino[1,2-*a*]pyrimidin-6-one as a scaffold for the formation of internal bicyclic peptidomimetics. Installing an acetal moiety on α -amino group following by tandem acid-catalyzed N-acycliminium ion bicyclization was the key for this transformation. The formation of the bicyclic scaffold is considered to follow a thermodynamically driven pathway to yield only a single diastereomer. We expect that this methodology will bring a simple and efficient tool to develop new types of peptide-based ligands.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work was supported by grants from US Public Health Service (DK 17420) and the National Institute on Drug Abuse (DA 06284 and DA 13449). We thank Prof. Richard Glass and Prof. Hamish Christie (The University of Arizona) for helpful discussions regarding the reaction mechanism.

References

1. Rose GD, Gierasch LM, Smith JA. Adv. Protein. Chem 1985;37:1–109. [PubMed: 2865874]
2. Burgess K. Acc. Chem. Res 2001;34:826–835. [PubMed: 11601967]
3. (a) Ndungu JM, Cain J, Davis P, Ma S-W, Vanderah TW, Lai J, Porreca F, Hruby VJ. Tetrahedron, Lett 2006;47:2233–2236. [PubMed: 17203138] (b) Gu X, Ying J, Min B, Cain JP, Davis P, Willey P, Navratilova E, Yamamura HI, Porreca F, Hruby VJ. Biopolymers 2005;80:151–163. [PubMed: 15660379] (c) Ndungu JM, Gu X, Gross DE, Cain JP, Carducci MD, Hruby VJ. Tetrahedron Lett 2004;45:4139–4142.
4. (a) Eguchi M, Lee MS, Nakanishi H, Stasiak M, Lovell S, Kahn M. J. Am. Chem. Soc 1999;121:12204–12205. (b) Kim H.-O, Nakanishi H, Lee MS, Kahn M. Org. Lett 2000;2:301–302. [PubMed: 10814307] (c) Eguchi M, Lee MS, Stasiak M, Kahn M. Tetrahedron Lett 2001;42:1237–1239. (d) Eguchi M, Shen RY, Shea JP, Lee MS, Kahn M. J. Med. Chem 2002;45:1395–1398. [PubMed: 11906279]
5. (a) Speckamp WN, Hiemstra H. Tetrahedron 1985;41:4367–4416. For reviews, see: Hiemstra, H.; Speckamp, WN. Comprehensive Organic Chemistry. Trost, BM., editor. 2. Pergamon Press; Oxford: 1991. p. 1047–1082. (c) Baldwin JE, Humle C, Edwards AJ, Schofield CJ. Tetrahedron Lett 1993;34:1665–1668. (d) Brodney MA, Padwa A. Tetrahedron Lett 1997;38:6153–6156. (e) Lee YS, Min BJ, Park YK, Lee JY, Lee SJ, Park H. Tetrahedron 1999;55:12991–12996. (f) Maryanoff CA, Zhang H-C, Cohen JH, Turchi IJ, Maryanoff CA. Chem. Rev 2004;104:1431–1628. [PubMed: 15008627](b)
6. Castro B, Dormoy JR, Evin G, Selve C. Tetrahedron Lett 1975;16:1219–1222.
7. Ho PT, Chang D, Zhong JW, Musso GF. Peptide Res 1993;6:10. [PubMed: 8439733]
8. For details, see supporting materials.
9. CCDC 639677 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
10. Ghosh SK, Hsung RP, Liu J. J. Am. Chem. Soc 2005;127:8260–8261. [PubMed: 15941240]
11. Frisch, MJ.; Trucks, GW.; Schlegel, HB.; Gill, PMW.; Johnson, BG.; Robb, MA.; Cheeseman, JR.; Keith, T.; Petersson, GA.; Montgomery, JA.; Raghavachari, K.; Al-Laham, MA.; Zakrzewski, VG.; Ortiz, JV.; Foresman, JB.; Cioslowski, J.; Stefanov, BB.; Nanayakkara, A.; Challacombe, M.; Peng, CY.; Ayala, PY.; Chen, W.; Wong, MW.; Andres, JL.; Replogle, ES.; Gomperts, R.; Martin, RL.; Fox, DJ.; Binkley, JS.; Defrees, DJ.; Baker, J.; Stewart, JJP.; Head-Gordon, M.; Gonzalez, C.; Pople, JA. Gaussian 03, revision B.05. Gaussian, Inc.; Pittsburgh, PA: 2003.
12. (a) Scott AP, Radom L. J. Physical Chem 1996;100:16502–16513. http://comp.chem.umn.edu/database/freq_scale.htm

13. Cheng J-F, Chen M, Arrhenius T, Nadzan A. *Tetrahedron Lett* 2002;43:6293–6295. Similar type of the enamide formation has been reported in synthesizing piperazine derivatives where the authors also suggested that the acyliminium ion was the intermediate:

Figure 1.
Bicyclic external (1) and internal (2) β -turn mimetics

Scheme 1. Synthesis of 7-benzyl-octahydropyrazino[1,2-a]pyrimidin-6-one (9)

Reagents and conditions: (a) Boc-*L*-Phe-OH, BOP, HOBT, NMM, DMF, 92%; (b) TFA, CH_2Cl_2 ; (c) 2,2-dimethoxyacetaldehyde, THF; (d) $\text{NaBH}(\text{OAc})_3$, AcOH, THF, 90% (3 steps overall); (e) Cbz-Cl, DIEA, CH_2Cl_2 , 96%; (f) formic acid, 89%; (g) H_2 , Pd/C, MeOH, 95%.

Figure 2.
X-ray crystal structure of (7S, 9aR)-7-benzyl-octahydropyrazino[1,2-a]pyrimidin-6-one, 9

Figure 3.
The lowest energy conformational structures of two diastereomers [y1] (10 and 11, stereoviews) from DFT calculation using Gaussian 03

Scheme 2. Synthesis of 1-(2-aminoethyl)-3-benzylpiperazin-2-one, 18

Reagents and conditions: (a) Boc-L-Phe-OH, BOP, HOBt, NMM, DMF, 95 %; (b) TFA, DCM; (c) 2,2-dimethoxyacetaldehyde, THF; (d) NaBH(OAc)₃, AcOH, THF, 90% (3 steps overall); (e) Cbz-Cl, DIEA, DCM, 81 %; (f) formic acid, 10 %; (g) H₂, Pd/C, MeOH, 60 %.