Review

Progress in the pathogenesis and genetics of Parkinson's disease

Yoshikuni Mizuno*, Nobutaka Hattori, Shin-ichiro Kubo, Shigeto Sato, Kenya Nishioka, Taku Hatano, Hiroyuki Tomiyama, Manabu Funayama, Yutaka Machida and Hideki Mochizuki

Department of Neurology, Juntendo University School of Medicine, 2-1-1 Hongo Bunkyo, Tokyo 113, Japan

Recent progresses in the pathogenesis of sporadic Parkinson's disease (PD) and genetics of familial PD are reviewed. There are common molecular events between sporadic and familial PD, particularly between sporadic PD and *PARK1*-linked PD due to α -synuclein (*SNCA*) mutations. In sporadic form, interaction of genetic predisposition and environmental factors is probably a primary event inducing mitochondrial dysfunction and oxidative damage resulting in oligomer and aggregate formations of α -synuclein. In *PARK1*-linked PD, mutant α -synuclein proteins initiate the disease process as they have increased tendency for self-aggregation. As highly phosphorylated aggregated proteins are deposited in nigral neurons in PD, dysfunctions of proteolytic systems, i.e. the ubiquitin–proteasome system and autophagy–lysosomal pathway, seem to be contributing to the final neurodegenerative process. Studies on the molecular mechanisms of nigral neuronal death in familial forms of PD will contribute further on the understanding of the pathogenesis of sporadic PD.

Keywords: Parkinson's disease; familial Parkinson's disease; pathogenesis; genetics

1. INTRODUCTION

Clinical features of Parkinson's disease (PD) were first described by [Parkinson \(1817\)](#). He reported six patients in his monograph published in 1817 and described most of the typical clinical features such as bradykinesia, rest tremor, postural instability, stooped posture and micrographia. He did not describe rigidity, which was described by [Charcot \(1888\)](#); he proposed to call this disease as PD. In this review, we will focus on the recent progress in the pathogenesis of neuronal death in sporadic as well as familial PD.

2. CLINICAL FEATURES OF PD

Clinical features of PD include bradykinesia, rest tremor, rigidity and postural instability. In addition to these four cardinal symptoms, loss of automatic movements such as loss of arm swing, loss of blinking, reptile stare, masked face and difficulty in two simultaneous motor acts comprise motor features of PD. Furthermore, many non-motor symptoms frequently appear in PD, such as cognitive impairment, hallucination, delusion, behavioural abnormalities, depression, disturbances of sleep and wakefulness, loss of smell, pain, and autonomic dysfunctions such as constipation, hypotension, urinary frequency, impotence and sweating. Definition of PD has been proposed in various ways. For the research purpose, British Brain

Bank criteria ([Hughes et al. 1992](#)) are frequently used, which define PD as those patients who have bradykinesia and at least one of the remaining four cardinal symptoms. Also, other causes of Parkinsonism have to be excluded by appropriate tests and/or clinical observation.

3. NEUROPATHOLOGY OF PD

The most characteristic features of neuropathology are loss of pigmented neurons in the substantia nigra (SN; [Tretiakoff 1919](#)) and the presence of eosinophilic cytoplasmic inclusion bodies (Lewy bodies); [Lewy \(1912\)](#) discovered these inclusions in the substantia innominata. Lewy bodies are usually absent in autosomal recessive young onset Parkinsonism due to *parkin* mutations ([Takahashi et al. 1994](#); [Mori et al. 1998](#)). Also, in *PARK8*-linked PD, Lewy bodies may or may not be present ([Wszolek et al. 2004](#)).

Neuronal loss and Lewy body formation are seen not only in SN but also in locus coeruleus, pedunculopontine nucleus, raphe nucleus, dorsal motor nucleus of the vagal nerve, olfactory bulb, parasympathetic as well as sympathetic post-ganglionic neurons, Meynert nucleus, amygdaloid nucleus and cerebral cortices. These lesions are responsible for non-motor symptoms of PD. [Braak et al. \(2003\)](#) proposed a hypothesis that in PD and PD with dementia (PDD), Lewy bodies were first formed in the dorsal motor nucleus and the olfactory bulb and slowly involved higher structures along the brain stem, diencephalon and the cerebral cortex.

* Author for correspondence (y_mizuno@med.juntendo.ac.jp).

One contribution of 17 to a Theme Issue 'Japan: its tradition and hot topics in biological sciences'.

Figure 1. Aetiological and pathogenetic factors of PD. Aetiology refers to the cause of the disease and pathogenesis represents molecular events that lead nigral neurons to death.

4. NEUROTOXIN-BASED MODELS OF PD

The following substances have been used in producing animal models of PD, i.e. 6-hydroxydopamine (6-OHDA), 1-methyl-4-phenyl-tetrahydropyridine (MPTP), mitochondrial complex I-inhibitors such as rotenone, and α -synuclein overexpression. Systemic administration of MPTP produces selective degeneration of the dopaminergic neurons in SN in both humans and animals (Davis *et al.* 1979; Burns *et al.* 1983; Langston *et al.* 1983). We were interested in how MPTP killed nigral neurons. MPTP is oxidized by glial monoamine oxidase B to 1-methyl-4-phenylpyridinium ion (MPP⁺; Chiba *et al.* 1984), which kills nigral neurons (Chiba *et al.* 1985; Javitch *et al.* 1985). We noted structural similarity between MPP⁺ and NAD⁺, which is an important cofactor of respiratory enzymes. We thought that MPP⁺ might inhibit the activities of mitochondrial NAD⁺-linked respiratory enzymes. Two groups reported inhibition of complex I by MPP⁺ (Nicklas *et al.* 1985; Ramsay *et al.* 1986). We independently made the same observation (Mizuno *et al.* 1987a). Furthermore, we found inhibition of the α -ketoglutarate dehydrogenase complex of the mitochondrial tricarboxylic acid cycle by MPP⁺ (Mizuno *et al.* 1987b). This enzyme synthesizes succinate from α -ketoglutarate; succinate is an electron donor for complex II of the electron transport chain. Thus dual inhibition of the activities of complexes I and II would impair deleteriously the electron transport and ATP synthesis. Inhibition of the electron transport induces oxidative damage by increasing the formation of reactive oxygen species.

5. AETIOLOGY AND PATHOGENESIS OF PD

We thought that mitochondrial function might be impaired in PD. Schapira *et al.* (1989) reported decreased activity of complex I in SN of PD. We found a decrease in complex I proteins by immunoblotting (Mizuno *et al.* 1989) and by immunohistochemistry (Hattori *et al.* 1991). We also found a decrease in the amount of α -ketoglutarate dehydrogenase complex in

SN of PD by immunohistochemistry (Mizuno *et al.* 1994). Thus biochemical changes in PD were essentially similar to those of the MPTP-induced Parkinsonism.

Oxidative damage is also an important factor for nigral neuronal death; increase in iron (Youdim *et al.* 1989), increase in lipid peroxides (Dexter *et al.* 1989), decrease in glutathione (Sofic *et al.* 1992), increase in hydroxynonenal-modified proteins (Yoritaka *et al.* 1996) and increase in 8-hydroxy-deoxy guanine (Shimura-Miura *et al.* 1999) were reported in SN of PD. Reactive oxygen species impair mitochondrial proteins, further aggravating mitochondrial function. Ultimate outcomes are the dissipation of mitochondrial membrane potential and the release of cytochrome *c* into cytoplasm activating the apoptotic cascade. Apoptotic nigral neuronal death has been postulated in PD (Mochizuki *et al.* 1996; Anglade *et al.* 1997). The interaction of genetic predisposition and environmental factors is believed to trigger mitochondrial dysfunction and oxidative damage in sporadic PD. Additional aetiological as well as pathogenetic factors are summarized in figure 1.

6. DYSFUNCTION OF PROTEIN DEGRADATION IN PD

In recent years, dysfunction of protein degradation has emerged as an important contributor to nigral neuronal death in PD. Impaired protein degradation is likely to follow mitochondrial dysfunction and oxidative damage. In the presence of oxidative stress or mutated proteins, the folding process of proteins may be impaired resulting in an increase in misfolded proteins. Misfolded proteins are generally cytotoxic and have to be removed by protein degrading systems. In eukaryotic cells, the ubiquitin–proteasome system and the autophagy–lysosomal pathway are two major protein degradation systems. The ubiquitin system consists of three enzymes, i.e. the ubiquitin-activating enzyme (E1), the ubiquitin-conjugating enzyme (E2) and the ubiquitin-protein ligase; these enzymes work

Figure 2. Schematic presentation of nigral neuronal death in PD. In sporadic PD, interaction of genetic predisposition and environmental risk factors is believed to initiate the pathological cascade. In familial PD due to *SNCA* mutations, mutant α -synuclein probably initiates the pathological process. DA, dopamine.

together to transfer ubiquitin molecules to target proteins that have to be metabolized by the 26S proteasome, which is an ATP-dependent proteolytic enzyme (Tanaka *et al.* 2004). Proteins with four or more than four ubiquitin molecules attached to lysine 48 residue of ubiquitin are recognized by the 26S proteasome. The 26S proteasome predominantly degrades short-lived nuclear and cytosolic proteins and misfolded proteins in the endoplasmic reticulum (Rubinsztein 2006).

The autophagy–lysosome pathway is able to degrade oligomers and aggregates of proteins as well as intracellular organelles (Yorimitsu & Klionsky 2005). In this pathway, double membrane-bound autophagosomes are generated by the elongation of small membranous structures, which encircle proteins to be digested. Then the autophagosome fuses with the lysosome to form the autophagosomes–lysosome. Then acidic hydrolases within the lysosome digest proteins that were incorporated into autophagosomes (Rubinsztein 2006).

What evidence do we have to suggest the dysfunction of protein degradation in PD? First of all, presence of Lewy bodies is strong evidence of impaired protein degradation. Lewy bodies consist of aggregated proteins and α -synuclein is the major component (Spillantini *et al.* 1997). Further interestingly, missense mutations of the *α -synuclein* gene (*SNCA*) cause autosomal dominant familial PD (Polymeropoulos *et al.* 1987; Krueger *et al.* 1998; Zarranz *et al.* 2004). α -Synuclein is a neuron-specific protein expressed predominantly in presynaptic membranes and the nucleus (Maroteaux *et al.* 1988). It is natively unfolded without a significant amount of secondary structure consisting of 140 amino acids (Weinreb *et al.* 1996).

Thus the aggregation of α -synuclein emerged as one of the most important processes in nigral degeneration in PD. The mutant α -synuclein has increased tendency for self-aggregation (El Agnaf *et al.* 1998). α -Synuclein is mainly located in the lipid raft in membranes (Fortin *et al.* 2004; Kubo *et al.* 2005) and this localization appears to be important in the trafficking of α -synuclein and its final localization in presynaptic and synaptic vesicular membranes.

α -Synuclein is degraded by both autophagy and the proteasome; however, mutant forms of α -synuclein and oligomers are dependent on the autophagy–lysosome pathway for their clearance (Webb *et al.* 2003). Wild-type α -synuclein is translocated into lysosomes for degradation by the chaperone-mediated autophagy pathway; however, mutant A53T and A30P proteins can bind to the chaperone-mediated autophagy-pathway receptor on the lysosomal membrane, but act as uptake blockers inhibiting their own degradation and that of other proteins (Cuervo *et al.* 2004). In sporadic PD, dysfunctions of both ubiquitin–proteasome and autophagy–lysosome systems appear to be present. As 26S proteasome is an ATP-dependent enzyme, dysfunction of mitochondria will compromise its function. Furthermore, oxidative stress enhances oligomer formation of α -synuclein. Thus formed oligomers impair membrane structures (Volles & Lansbury 2002) such as synaptic vesicles and mitochondria, and further increase oxidative stress and mitochondrial dysfunction. In this way, vicious cycles will be formed within nigral neurons leading to neuronal death (figure 2). Decrease in the 26S proteasomal activity was reported in PD (McNaught *et al.* 2003).

We have only indirect evidence of lysosomal dysfunction in PD. Activation of lysosomal functions

were reported by treating cell lines with Parkinsonism-inducing neurotoxins, such as overexpression of mutant α -synuclein (Stefanis *et al.* 2001), or proteasomal inhibitors (Ding *et al.* 2003; Rideout *et al.* 2004). Other indirect evidence came from studies on the association of Gaucher disease and PD. Gaucher disease is an autosomal recessive lysosomal lipid storage disease caused by mutations of a lysosomal enzyme, glucocerebrosidase β -glucosidase. Gaucher disease and its carrier state appear to be risk factors for PD (Tayebi *et al.* 2003; Ahron-Peretz *et al.* 2004; Goker-Alpan *et al.* 2004, 2006; Lwin *et al.* 2004; Sato *et al.* 2005; Kono *et al.* 2007). Furthermore, the recently identified *ATP13A2*, the disease gene for *PARK9*-linked PD (Kufor-Rakeb syndrome), encodes a lysosomal membrane protein (Ramirez *et al.* 2006). These two observations indicate the importance of lysosomal function for the maintenance of nigral neurons.

7. PROGRESS IN FAMILIAL PD

Thirteen chromosome loci have been identified to be linked to familial forms of PD (table 1). As *PARK1* and *PARK4* represent the same locus, the number of the familial forms is 12.

(a) *PARK1*- and *PARK4*-linked PD

PARK1- and *PARK4*-linked PD is an autosomal dominant one caused by mutations of the α -synuclein gene (*SNCA*); *PARK1* is caused by missense mutations and *PARK4* by multiplications of *SNCA*.

Three missense mutations, i.e. A53T (Polymeropoulos *et al.* 1997), A30P (Krüger *et al.* 1998) and E46K (Zarranz *et al.* 2004), duplications (Chartier-Harlin *et al.* 2004; Ibanez *et al.* 2004; Nishioka *et al.* 2006; Fuchs *et al.* 2007) and triplications (Singleton *et al.* 2003; Farrer *et al.* 2004) of *SNCA* are known. Missense mutations are very rare; A53T is limited to families with Greece origin; only one German family with A30P mutation and one Spanish family with E46K mutation are known. Multiplications of *SNCA* appear to be more common. Singleton *et al.* (2003) reported triplication of *SNCA* in a large kindred (Iowanian family). The triplication involved the 1.5 Mb region; exons of the adjacent genes on both side of *SNCA* were also triplicated. The amount of protein expressed would be doubled. Duplications of *SNCA* (Chartier-Harlin *et al.* 2004; Ibanez *et al.* 2004; Nishioka *et al.* 2006) were also reported. In the recently reported Swedish-American family (Fuchs *et al.* 2007), patients in the Swedish branch had duplication and those in the American branch (Farrer *et al.* 2004) had triplication. They suggested unequal recombination and unequal crossing over as the potential mechanisms for duplication and triplication, respectively (Fuchs *et al.* 2007).

There is a clinico-genetic correlation. E46K mutation and triplications are associated with Parkinsonism and dementia, and the age of onset is younger than the other mutations; neuropathological changes are those of diffuse Lewy body disease. A30P mutation is usually not associated with dementia. Duplication usually does not cause dementia but it can happen (Nishioka *et al.* 2006; Fuchs *et al.* 2007). A53T

Table 1. Familial forms of PD. (AD, autosomal dominant; AR, autosomal recessive; LB, Lewy bodies; SP, sporadic.)

type	loci	genes	Inh	LB
<i>PARK1</i>	4q21–23	α -synuclein (<i>SNCA</i>)	AD	+
<i>PARK2</i>	6q25.20–27	<i>parkin</i> (<i>PRKN</i>)	AR	±
<i>PARK3</i>	2p13	<i>unknown</i>	AD	+
<i>PARK4</i>	4q21–23	α -synuclein (<i>SNCA</i>)	AD	+
<i>PARK5</i>	4p14	<i>UCH-L1</i>	AD	
<i>PARK6</i>	1p35–36	<i>PINK1</i>	AR	+
<i>PARK7</i>	1p36	<i>Dγ-1</i>	AR	
<i>PARK8</i>	12p11.2–q13.1	<i>LRRK2</i>	AD	±
<i>PARK9</i>	1p36	<i>ATP13A2</i>	AR	
<i>PARK10</i>	1p32	<i>unknown</i>	SP	
<i>PARK11</i>	2q36–37	<i>unknown</i>	AD	
<i>PARK12</i>	Xq21–25	<i>unknown</i>	SP	
<i>PARK13</i>	2p13	Omi/HtrA2	AD?	

mutation may cause dementia and cortical Lewy bodies are reported (Golbe *et al.* 1990).

Regarding the pathogenesis of *PARK1*-linked PD, increased tendency for oligomer and aggregate formations of mutant α -synuclein is likely to be the cause (El Agnaf *et al.* 1998; Fredenburg *et al.* 2007). In duplication and triplication, increased amount of normal α -synuclein is probably predisposing nigral neurons for oligomer and aggregate formations. Recently, two groups independently reported that *SNCA* polymorphic mutations are significant risk factors for sporadic PD (Mueller *et al.* 2005; Mizuta *et al.* 2006); some of those polymorphic mutations were associated with increased α -synuclein expression. Thus the molecular mechanism of nigral degeneration is similar between *SNCA*-mutated and sporadic PDs.

Regarding the toxicity of oligomers, Volles & Lansbury (2002) reported that protofibrillar Ala30Pro and Ala53Thr had greater permeabilizing activities per mole than the wild-type protein. The leakage of vesicular contents induced by protofibrillar α -synuclein exhibited a strong preference for low-molecular mass molecules like dopamine, suggesting a pore-like mechanism for permeabilization.

(b) *PARK2*-linked PD

PARK2-linked PD is an autosomal recessive young onset PD. Clinical features were first described by Yamamura *et al.* in 1973. The usual age of onset is between 20 and 40, but it can be before 10 years and above 60 years. When the age of onset is young, dystonic features and sleep benefits are characteristic symptoms; sleep benefit represents temporal improvement in Parkinsonism after a sleep or nap. They respond well to L-dopa; however, they will soon develop motor fluctuations. Pathologically, SN undergoes severe neuronal loss and gliosis; the locus coeruleus is much less severely involved. Usually no Lewy bodies are seen (Takahashi *et al.* 1994; Mori *et al.* 1998), although rare Lewy body positive cases were reported (Farrer *et al.* 2001).

We identified the disease gene as follows. While we were doing an association study between the genetic polymorphism of the manganese superoxide dismutase

Figure 3. A schematic of the *parkin* gene. The coding region consists of 12 exons. In the amino terminal region, ubiquitin-like domain is indicated as 'Ubl', and in the carboxyl terminal side, two RING finger motives are indicated as 'RING1' and 'RING2'. RING, rare interesting gene; IBR, in between RINGs.

gene (*sod2*) and sporadic PD (Shimoda-Matsubayashi *et al.* 1996), we found a family that appeared to be linked to the *sod2* locus, which had been mapped to the telomeric region of the long arm of chromosome 6. We did linkage analysis on 13 similar families and mapped the disease locus to the long arm of chromosome 6 near the *sod2* locus (6q25.2–27; Matsumine *et al.* 1997).

While we were doing linkage analysis on additional families, we found a patient who showed deletion of one of the microsatellite markers (D6S305) that we were using in the linkage analysis (Matsumine *et al.* 1998). We thought that this microsatellite marker might be located within the disease gene. By screening the Keio BAC library (Asakawa *et al.* 1997) using D6S305, we cloned a cDNA consisting of 2960 base pairs, of which 1395 base pairs constituted the open reading frame (Kitada *et al.* 1998). As this was a novel gene, we named it *parkin*. The total size of *parkin* was 1.4 Mb, the second largest gene after *dystrophin*. The number of exons was 12. The gene product consisted of 462 amino acids. There were unique structures in parkin protein (figure 3). There was 30% homology to ubiquitin in the amino terminal domain and there were two RING-finger-like motifs in the carboxyl half of the protein. RING stands for rare interesting gene and RING-like structures have been found in proteins with ubiquitin-ligase activity (Lorick *et al.* 1999). By northern blot, parkin messengers were ubiquitously expressed including the systemic organs (Kitada *et al.* 1998).

Reported mutations in *parkin* now exceed 100 (Hattori *et al.* 1998; Abbas *et al.* 1999; Klein *et al.* 2000; Kann *et al.* 2002; Khan *et al.* 2003; Hedrich *et al.* 2004). Not only exonic deletions but also missense and nonsense mutations and multiplications of exons were reported.

We thought that parkin might be related to the ubiquitin-proteasome system (UPS); in fact the parkin protein had ubiquitin-protein ligase activity (Shimura *et al.* 2000). Since then many parkin-interacting proteins have been reported; the following are substrate candidates: CDCrel-1 (Zhang *et al.* 2000), which is believed to be negatively regulating transmitter release; glycosylated α -synuclein (Shimura *et al.* 2001); synphilin-1

(Chung *et al.* 2001), which is an α -synuclein-interacting protein; PAEL-receptor (Imai *et al.* 2001), which is an endoplasmic reticulum protein; p38 (Corti *et al.* 2003); polyglutamine (Tsai *et al.* 2003); α - and β -tubulins (Ren *et al.* 2003); cyclin-E (Staropoli *et al.* 2003); SEPT5_v2 (Choi *et al.* 2003), which is also known as cell division-control protein-2; DJ-1 (Moore *et al.* 2005); RanBP2 (Um *et al.* 2006), which is a protein localized in the cytosolic filament of the nuclear core complex; and protein-1 (Ko *et al.* 2006). Also, parkin-regulatory proteins have been reported, i.e. CHIP (Imai *et al.* 2002), which is a chaperone; HSP-70 (Imai *et al.* 2002); Rpn10 subunit (Sakata *et al.* 2003); BAG5 (Kalia *et al.* 2004); Nrdp1/FLRF (Zhong *et al.* 2005); LRRK2 (Smith *et al.* 2005); and 14-3-3 η (Sato *et al.* 2006a,b). In addition, non-lysine 48-related ubiquitylation substrates were reported. Lysine-63 polyubiquitylation is believed to be related to endocytosis, DNA repair, translation, $\text{I}\kappa\text{B}$ activation, DNA silencing, virus budding, protein sorting and protein trafficking (Tanaka *et al.* 2004). Recently, Lim *et al.* (2005) reported that synphilin-1 was polyubiquitylated at lysine 63 residue of ubiquitin. Other lysine sites are mono-ubiquitylated. Despite vast number of parkin-interacting proteins, there is no immunohistochemical proof of accumulation of above parkin-interacting proteins in autopsied patients.

Parkin-knockout (KO) mice do not show nigral neuronal loss or striatal dopamine deficiency (Goldberger *et al.* 2003; von Coelln *et al.* 2004; Perez & Palmiter 2005; Ko *et al.* 2005; Sato *et al.* 2006). What were reported are only subtle changes in dopaminergic functions. von Coelln *et al.* (2004) found some loss of neurons in the locus coeruleus and reduced startle response. Ko *et al.* (2005) found an increase in the amount of the aminoacyl-tRNA synthase cofactor p38 in the midbrain/hindbrain region of both young and old parkin-null mice. They postulated that p38 is a substrate of parkin as E3 ligase. They further showed that overexpression of p38 in the SN in mice lead to loss of dopaminergic neurons. They analysed the level of p38 in the cortical regions of the patients with parkin mutations by Western blotting; they found an increase in p38. We studied striatal dopamine receptors by *ex*

vivo autoradiography. In *parkin*-KO mice, both striatal D1 and D2 receptor bindings were significantly increased when compared with wild mice. Midbrain dopamine content was increased in KO mice. Increase in D1 and D2 receptor bindings in the striatum would indicate reduction in dopamine release; increase in dopamine in nigral neurons would cause oxidative stress.

Ved *et al.* (2005) found increased sensitivity of mitochondria to complex I inhibitors such as rotenone, fenperoximate, pyridaben or stigmatellin in their *parkin*-KO *Caenorhabditis elegans*; they observed similar effects by overexpressing α -synuclein, or knocking down *D β -1*. Further interestingly, *parkin*-KO *Drosophila* produced by Greene *et al.* (2003) exhibited reduced lifespan, locomotor defects and male sterility. Locomotor defects were due to muscle degeneration with mitochondrial damage consisting of disruption and disintegration of the cristae. There was no neuronal loss in the brain including dopaminergic neurons in these flies. Pesah *et al.* (2004) also reported similar findings. Then Clark *et al.* (2006a,b) and Park *et al.* (2006) made *PINK1/parkin* double KO *Drosophila*. Both groups reported that overexpression of parkin rescued *PINK1*-KO-induced muscle damage; but *PINK1* overexpression could not rescue *parkin*-KO-induced damage. Furthermore, they showed interaction of parkin and *PINK1*. They concluded that parkin was functioning in the downstream of *PINK1* in a common pathway to keep mitochondrial integrity. Their flies also showed dopaminergic neuronal degeneration.

At a cellular level, Machida *et al.* (2005) constructed a parkin-knockdown cell line using SH-SY5Y cells, which showed apoptotic cell death. Furthermore, they found increase in the auto-oxidized forms of L-dopa and dopamine (Dopa chrome+dopamine chrome), suggesting the presence of anti-oxidative property in parkin. We reported a profound accumulation of iron in the SN of a *parkin*-mutated patient (Takanashi *et al.* 2001). Taken together, oxidative stress appears to be a pathogenetic pathway common to *PARK2*-linked and sporadic PDs.

(c) *PARK6*-linked PD

PARK6-linked PD is another form of young onset autosomal recessive PD. The age of onset is slightly older than that of *PARK2*, i.e. from 32 to 48 years (Valente *et al.* 2001). Therefore, dystonic features and sleep benefit are uncommon. Affected patients show L-dopa-responsive Parkinsonism. The disease gene was identified as *PINK1* (*PTEN-induced kinase 1*; Valente *et al.* 2004a,b). *PTEN* stands for *phosphatase with tensin homology*. *PINK1* has eight exons and cDNA spans 1.8 kb. It encodes a protein with 581 amino acids. The protein is ubiquitously expressed including brain and systemic organs. Interestingly, it is a mitochondrial protein located in the matrix and the intermembrane space. It has a serine/threonine protein kinase domain. However, its function is not known (Valente *et al.* 2004).

Many missense and nonsense mutations have been reported (Hatano *et al.* 2004; Healy *et al.* 2004; Rohe *et al.* 2004; Valente *et al.* 2004; Li *et al.* 2005). In

contrast to *parkin*, most of the *PINK1* mutations reported are either missense or nonsense mutations, although one family with a large deletion mutation is known (Li *et al.* 2005); this deletion involved exons 6–8 homozygously.

Regarding the molecular mechanisms of nigral neuronal death, as *PINK1* has a protein kinase domain as *LRRK2* has, dysfunction in some phosphorylation reactions may be important in the pathogenesis. *PINK1* is inducible by *PTEN* and *PTEN* is an oncogene mutated in many neoplastic cells (Li *et al.* 1997). As oncogenesis and degeneration are the opposite sides of eukaryotic cell fate, elucidation of the function of *PINK1* in relation to *PTEN* is an interesting topic.

(d) Do heterozygotes for parkin or PINK1 mutations develop PD?

In autosomal recessive diseases, usually both alleles should have a mutation to show the disease. But in cases of *PINK1* (Valente *et al.* 2004; Bonifati *et al.* 2005; Fung *et al.* 2006; Criscuolo *et al.* 2006; Toft *et al.* 2007) as well as *parkin* mutations (Hedrich *et al.* 2002; Khan *et al.* 2002; West *et al.* 2002; Oliveira *et al.* 2003; Clark *et al.* 2006a,b; Schlitter *et al.* 2006), at times only one mutation can be found (heterozygote). The question is, how do they get the disease? There are several possibilities. First of all, the second mutation may be localized in a place that is difficult to find by the currently available methods. The second possibility is the haploinsufficiency (West *et al.* 2002); here the amount of normal gene product is not sufficient to keep nigral neurons alive. But usually in autosomal recessive diseases, only one normal gene is sufficient to prevent the disease; parents (usually carriers) of a patient are normal in most of the cases. The third possibility is the interaction at the protein level; the mutated protein might interfere with the functions of normal protein (dominant-negative effect); but this possibility has not been proved. Finally, single heterozygous state might be acting as a risk factor for sporadic PD (Schlitter *et al.* 2006). Another interesting observation in *parkin* heterozygotes was made by Oliveira *et al.* (2003) who reported that mutations in the first RING finger domain tended to be heterozygotic and associated with later age of onset.

(e) *PARK7*-linked PD

PARK7-linked PD is another young onset PD. Clinical features are very similar to those of *PARK2*-linked PD (Duijin *et al.* 2001); the age of onset is usually 20–40 years. Some atypical features such as psychiatric symptoms (anxiety attacks; Dekker *et al.* 2003), and short stature and brachydactyly (Dekker *et al.* 2004) have been reported. The disease gene was identified as *D β -1* (Bonifati *et al.* 2003), which had been cloned by Nagakubo *et al.* (1997). The size of *D β -1* is 24 kb with eight exons encoding a protein consisting of 189 amino acids. *PARK7*-linked PD is very rare (Bonifati *et al.* 2003; Hague *et al.* 2003; Hering *et al.* 2004).

The function of DJ-1 protein is not well known. The active form of DJ-1 is a dimer of monomeric DJ-1. DJ-1 is a cytoplasmic protein; however, it can translocate into the mitochondria. It has a strong anti-oxidative property (Nagakubo *et al.* 1997; Abou-Sleiman *et al.*

2003; Canet-Aviles *et al.* 2004; Moore *et al.* 2005) that depends on its cysteine residue at 106, which undergoes oxidation to form a disulphide bond (Canet-Aviles *et al.* 2004). Downregulation of endogenous DJ-1 protein of the neuronal cell line by siRNA was reported to enhance the cell death induced by oxidative stress, ER stress and proteasome inhibition, but not by proapoptotic stimulus (Yokota *et al.* 2003). The Leu166-Pro mutant DJ-1 protein has a reduced anti-oxidative activity (Takahashi-Niki *et al.* 2004). DJ-1 protein expression is increased upon oxidative stress induced by paraquat (Mitsumoto *et al.* 2001). As nigral neurons are exposed to high oxidative stress owing to the presence of dopamine, DJ-1 may be acting as a strong anti-oxidative protein. As mutant DJ-1 was reported to interact with parkin (Moore *et al.* 2005), parkin might be acting as E3 ligase to remove mutated DJ-1.

(f) PARK8-linked PD

PARK8-linked PD is an autosomal dominant PD linked to the centromeric region of chromosome 12 (Funayama *et al.* 2002). Clinical features were described back in 1978 on a large Japanese family (Nukada *et al.* 1978); clinical features are essentially similar to those of sporadic PD, except for slightly earlier onset of age. Dementia is not a common feature but it is known to occur (Wszolek *et al.* 1997).

The disease gene was identified as *lrrk2* (Paisan-Ruiz *et al.* 2004; Zimprich *et al.* 2004); *lrrk2* is a huge gene encompassing 144 kb, consisting of 7449 bp and encoding a protein consisting of 2517 amino acids, and has 51 exons. The carboxyl half of the LRRK2 contains several functional domains such as ANK (ankyrin-repeat domain), LRR (leucine-repeat-rich), ROC (Ras of complex proteins), COR (carboxy terminal of ROC), MAPKKK (mitogen activated protein kinase kinase) and WD domain that is rich in tryptophan and aspartate repeats. Pathogenetic mutations are concentrated in these functional domains.

PARK8-linked PD is now believed to be the most common form of autosomal dominant familial PD and 20 missense or nonsense mutations have been reported (Paisan-Ruiz *et al.* 2004; Zimprich *et al.* 2004; Aasly *et al.* 2005; Di Fonzo *et al.* 2005; Funayama *et al.* 2005; Hernandez *et al.* 2005; Kachergus *et al.* 2005; Nichols *et al.* 2005; Paisan-Ruiz *et al.* 2005; Mata *et al.* 2006). *lrrk2* Mutations were also found in some of the apparently sporadic PD patients (Gilks *et al.* 2005). One of the polymorphic mutations, G2385R, is a genetic risk factor for sporadic PD in Asian populations (Di Fonzo *et al.* 2006; Funayama *et al.* 2007; Tan *et al.* 2007).

Four different neuropathologies were reported within the same family (Wszolek *et al.* 2004); one of their patients showed brain stem-type Lewy body disease, the second showed diffuse-type Lewy body disease, the third accumulation of tau in the remaining nigral neurons and the last simple nigral atrophy. This observation tells us the difficulty of defining a disease by neuronal inclusions.

Function of LRRK2 is not well known. The ROC domain is able to bind GTP but it does not have GTPase activity, but GTP binding is essential for the MAPKKK domain to exert kinase activity (Ito *et al.*

2007); some of the mutant LRRK2 have increased kinase activity (Gloeckner *et al.* 2006). Other functional domains are believed to be important in protein-protein interactions (Zimprich *et al.* 2004). LRRK2 also interacts with other familial PD proteins; Smith *et al.* (2005) reported interaction of LRRK2 with parkin through the ROC domain; however, the interaction with parkin did not enhance polyubiquitylation of LRRK2.

Recently, Hatano *et al.* (2007) made a detailed observation on the intracellular distribution of LRRK2. It was found to be present in Golgi apparatus, plasma membrane, synaptic vesicles and particularly in the lipid rafts; presence in the lipid rafts suggests that LRRK2 is probably involved in signal transduction, membrane trafficking and cytoskeletal organization (Brawn & London 1998). Biskup *et al.* (2006) also reported the presence of LRRK2 in membrane structures, such as lysosomes, endosomes, transport vesicles and mitochondria. In this regard it is interesting to note that α -synuclein is also expressed in the presynaptic membranes and lipid rafts (Fortin *et al.* 2004; Kubo *et al.* 2005).

(g) PARK9-linked PD

PARK9-linked PD is an autosomal recessive disorder characterized by L-dopa-responsive Parkinsonism, supranuclear gaze palsy, pyramidal sign and dementia; it is also called as Kufor-Rakeb syndrome; the name of the initial Jordanian family with this disorder (Najim Al-Din *et al.* 1994). Age of onset was very early, between 11 and 16 years. MRI showed significant atrophy of the globus pallidus and the pyramids, as well as generalized brain atrophy in later stages. Some of them developed facial-facial-finger mini-myoclonus, visual hallucinations and oculogyric dystonic spasm (Williams *et al.* 2005).

Hamphire *et al.* (2001) performed linkage analysis on this Kufor-Rakeb family and mapped the disease locus to the short arm of chromosome 1 at 1p36 with a maximum LOD score of 3.6, the hot spot for autosomal recessive familial PD. The disease gene was identified as *ATP13A2* (Ramirez *et al.* 2006), which is a lysosomal membrane protein with an ATPase domain; exact function is still unknown. It is interesting to note that mutations of a lysosomal membrane protein can induce nigral degeneration, suggesting the importance of lysosomes for the maintenance of the integrity of nigral neurons.

(h) Other forms of familial PD

PARK3-linked PD is an autosomal dominant PD. Clinical features are essentially similar to those of sporadic PD with the age of onset between 36 and 89 (Gasser *et al.* 1998). Patients from two out of six families reported in that literature developed dementia. Autopsy findings in two families showed nigral degeneration and neurofibrillary tangles in cortical neurons.

PARK5-linked PD is an autosomal dominant PD. Only one family is reported (Leroy *et al.* 1998). Clinical features are similar to those of sporadic PD with the age of onset from 49 to 50. The disease gene was reported as ubiquitin carboxyl-terminal hydrolase-L1 (*UCH-L1*; Leroy *et al.* 1998). Ile93Met missense mutation

was found in the affected members. Deletion of exons 7 and 8 in mouse *UCH-L1* causes gracile axonal dystrophy (*gad* mouse); this is an autosomal recessive condition characterized by axonal degeneration and formation of spheroid bodies in motor and sensory nerve terminals (Saigo *et al.* 1999).

UCH-L1 is an enzyme that cleaves carboxy-terminal peptide bond of polyubiquitine chains. Thus UCH-L1 is an ubiquitin-recycling enzyme. UCH-L1 is a neuron-specific enzyme. Catalytic activity of Ile93Met-mutated UCH-L1 was reported to be half of the wild enzyme (Leroy *et al.* 1998). Thus the supply of ubiquitin for 26S proteasome may be reduced with this mutation.

PARK10 was found by genome-wide scanning. Hicks *et al.* (2002) studied 51 Icelandic families with more than one PD patient; they analysed 117 patients and 168 of their unaffected relatives using 781 micro-satellite markers. Allele-sharing, model-independent analysis of their results showed linkage to a region on chromosome 1p32 with a LOD score of 4.9. They designate this region *PARK10*. The disease gene has not been identified yet. Clinical features are essentially similar to those of sporadic PD and the mean age of onset was 65.8 years.

PARK11 was also found by genome-wide scanning; Pankratz *et al.* (2003a,b) screened 85 families with a very strong family history of PD and found an evidence of linkage to the long arm of chromosome 2 (LOD=4.9). Clinical features are essentially similar to those of sporadic PD with the mean age of onset at 58 years.

PARK12 was also found by genome-wide scanning on sporadic PD; Pankratz *et al.* (2003a,b) screened 277 families without a strong family history of PD and detected linkage to the long arm of chromosome X (LOD=3.2).

PARK13-linked PD was reported to have a mutation (G399S) in *Omi/HtrA2* (Strauss *et al.* 2005); they found G399S mutation in four German sporadic PD patients. They also identified a novel A141S polymorphism that was associated with PD ($p<0.05$). Both mutations resulted in defective activation of the protease activity of Omi/HtrA2. Further studies are necessary before concluding that this is a new familial PD-inducing protein. A cDNA of HtrA2 was first isolated by Faccio *et al.* (2000); it encoded a protein (Omi) consisting of 458 amino acids and had homology to bacterial HtrA endoprotease and had a PDZ domain. Omi mRNA was expressed ubiquitously, and the gene was localized on human chromosome 2p12 near the *PARK3* locus; however, *PARK3*-linked PD patients did not have this mutation. Omi/HtrA2 is a nuclearly encoded mitochondrial protein localized in the intermembrane space. Further interestingly, it has a serine protease domain and it binds to apoptosis-inhibiting protein upon release into the cytoplasm (Suzuki *et al.* 2001). G399S mutant Omi/HtrA2 induces mitochondrial dysfunction associated with altered mitochondrial morphology and cells over-expressing G399S mutant Omi/HtrA2 are more susceptible to stress-induced cell death than wild-type (Strauss *et al.* 2005).

Finally, there are many families in which linkage analysis failed to show linkage to any one of the

known loci that are associated with familial PD. By elucidating the functions of familial PD proteins, pathogenesis of sporadic PD will be better understood.

This study was in part supported by the grant for research on diseases of the brain from Ministry of Education, Science and Sports in Japan.

REFERENCES

- Aasly, J. O., Toft, M., Fernandez-Mata, I., Kachergus, J., Hulihan, M., White, L. R. & Farrer, M. 2005 Clinical features of LRRK2-associated Parkinson's disease in central Norway. *Ann. Neurol.* **57**, 762–765. (doi:10.1002/ana.20456)
- Abbas, N. *et al.* 1999 A wide variety of mutations in the *parkin* gene are responsible for autosomal recessive Parkinsonism in europe. *Hum. Mol. Gen.* **8**, 567–574. (doi:10.1093/hmg/8.4.567)
- Abou-Sleiman, P. M., Healy, D. G., Quinn, N., Lees, A. J. & Wood, N. W. 2003 The role of pathogenic DJ-1 mutations in Parkinson's disease. *Ann. Neurol.* **54**, 283–286. (doi:10.1002/ana.10675)
- Ahron-Peretz, J., Rosenbaum, H. & Gershoni-Baruch, H. 2004 Mutations in the glucocerebrosidase N370S allele and Parkinson's disease in Ashkenazi Jews. *N. Engl. J. Med.* **351**, 1972–1977. (doi:10.1056/NEJMoa033277)
- Anglade, P. *et al.* 1997 Apoptosis and autophagy in nigral neurons of patients with Parkinson's disease. *Histol. Histopathol.* **12**, 25–31.
- Asakawa, S. *et al.* 1997 Human BAC library, construction and rapid screening. *Gene* **191**, 69–79. (doi:10.1016/S0378-1119(97)00044-9)
- Biskup, S. *et al.* 2006 Localization of LRRK2 to membranous and vesicular structures in mammalian brain. *Ann. Neurol.* **60**, 557–569. (doi:10.1002/ana.21019)
- Bonifati, V. *et al.* 2003 Mutations in the *Dj-1* gene associated with autosomal recessive early-onset Parkinsonism. *Science* **299**, 256–259. (doi:10.1126/science.1077209)
- Bonifati, V. *et al.* 2005 Early-onset Parkinsonism associated with PINK1 mutations. Frequency, genotypes, and phenotypes. *Neurology* **65**, 87–95. (doi:10.1212/01.wnl.0000167546.39375.82)
- Braak, H., Del Tredici, K., Rub, U., de Vos, R. A., Jansen Steur, E. N. & Braak, E. 2003 Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiol. Aging* **24**, 197–211. (doi:10.1016/S0197-4580(02)00065-9)
- Brawn, D. A. & London, E. 1998 Functions of lipid rafts in biological membrane. *Annu. Rev. Cell. Dev. Biol.* **14**, 111–136. (doi:10.1146/annurev.cellbio.14.1.111)
- Burns, R. S., Chiueh, C. C., Markey, S., Ebert, M. H., Jakobowicz, D. & Kopin, I. J. 1983 A primate model of Parkinson's disease, selective destruction of substantia nigra, pars compacta dopaminergic neurons by N-methyl-4-phenyl-1,2,3,6-tetrahydropyridine. *Proc. Natl Acad. Sci. USA* **80**, 4546–4550. (doi:10.1073/pnas.80.14.4546)
- Canet-Aviles, R. M. *et al.* 2004 The Parkinson's disease protein DJ-1 is neuroprotective due to cysteine-sulfinic acid-driven mitochondrial localization. *Proc. Natl Acad. Sci. USA* **101**, 9103–9108. (doi:10.1073/pnas.0402959101)
- Charcot, J.-M. 1888 *Leçons du Mardi à La Salpêtrière*, (1887–1888). Paris, France: Bureaux du Progrès.
- Chartier-Harlin, M. C. *et al.* 2004 Alpha-synuclein locus duplication as a cause of familial Parkinson's disease. *Lancet* **364**, 1167–1169. (doi:10.1016/S0140-6736(04)17103-1)
- Chiba, K., Trevor, A. J. & Castagnoli Jr, N. 1984 Metabolism of the neurotoxic tertiary amine, MPTP, by brain monoamine oxidase. *Biochem. Biophys. Res. Commun.* **120**, 574–578. (doi:10.1016/0006-291X(84)91293-2)

- Chiba, K., Trevor, A. & Castagnoli Jr, N. 1985 Active uptake of MPP⁺, metabolite of MPTP, by brain synaptosomes. *Biochem. Biophys. Res. Commun.* **128**, 1228–1232.
- Choi, P. et al. 2003 SEPT5_v2 is a parkin binding protein. *Mol. Brain Res.* **117**, 179–189. (doi:10.1016/S0169-328X(03)00318-8)
- Chung, K. K. K., Zhang, Y., Lim, K. L., Tanaka, Y., Huang, H., Gao, J., Ross, C. A., Dawson, V. L. & Dawson, T. M. 2001 Parkin ubiquitinates the α -synuclein-interacting protein synphilin-1, implications for Lewy-body formation in Parkinson disease. *Nat. Med.* **7**, 1144–1150. (doi:10.1038/nm1001-1144)
- Clark, I. E., Dodson, M. W., Jiang, G., Cao, J. H., Huh, J. R., Seol, J. H., Yoo, S. J., Hay, B. A. & Guo, M. 2006a *Drosophila* pink1 is required for mitochondrial function and interacts genetically with *Parkin*. *Nature* **441**, 1162–1166. (doi:10.1038/nature04779)
- Clark, L. N. et al. 2006b Case-control study of the *Parkin* gene in early-onset Parkinson disease. *Arch. Neurol.* **63**, 548–552. (doi:10.1001/archneur.63.4.548)
- Corti, O. et al. 2003 The p38 subunit of the aminoacyl-tRNA synthetase complex is a parkin substrate, linking protein biosynthesis and neurodegeneration. *Hum. Mol. Genet.* **12**, 1427–1437. (doi:10.1093/hmg/ddg159)
- Creene, J. C., Whitworth, A. J., Kuo, I., Andrews, L. A., Feany, M. B. & Pallanck, L. J. 2003 Mitochondrial pathology and apoptotic muscle degeneration in *Drosophila* parkin mutants. *Proc. Natl Acad. Sci. USA* **100**, 4078–4083. (doi:10.1073/pnas.0737556100)
- Crisuolo, C. et al. 2006 PINK1 homozygous W437X mutation in a patient with apparent dominant transmission of Parkinsonism. *Mov. Disord.* **21**, 1265–1267. (doi:10.1002/mds.20933)
- Cuervo, A. M., Stefanis, L., Fredenburg, R., Lansbury, P. T. & Sulzer, D. 2004 Impaired degradation of mutant α -synuclein by chaperone-mediated autophagy. *Science* **305**, 1292–1295. (doi:10.1126/science.1101738)
- Davis, G. C., Williams, A. C., Markey, S. P., Ebert, M. H., Caine, E. D., Reichert, C. M. & Kopin, I. J. 1979 Chronic Parkinsonism secondary to intravenous injection of meperidine analogues. *Psych. Res.* **1**, 249–254. (doi:10.1016/0165-1781(79)90006-4)
- Dekker, M. et al. 2003 Clinical features and neuroimaging of PARK7-linked Parkinsonism. *Mov. Disord.* **18**, 751–757. (doi:10.1002/mds.10422)
- Dekker, M. C., Galjaard, R. J., Snijders, P. J., Heutink, P., Oostra, B. A. & van Duijn, C. M. 2004 Brachydactyly and short stature in a kindred with early-onset Parkinsonism. *Am. J. Med. Genet. A* **130**, 102–104. (doi:10.1002/ajmg.a.30021)
- Dexter, D. T., Carter, C. J., Wells, F. R., Javoy-Agid, F., Agid, Y., Lees, A., Jenner, P. & Marsden, C. D. 1989 Basal lipid peroxidation in substantia nigra is increased in Parkinson's disease. *J. Neurochem.* **52**, 381–389. (doi:10.1111/j.1471-4159.1989.tb09133.x)
- Di Fonzo, A. et al. 2005 A frequent *LRRK2* gene mutation associated with autosomal dominant Parkinson's disease. *Lancet* **365**, 412–415.
- Di Fonzo, A. et al. 2006 A common missense variant in the *LRRK2* gene, Gly 2385Arg, associated with Parkinson's disease risk in Taiwan. *Neurogenetics* **7**, 133–138. (doi:10.1007/s10048-006-0041-5)
- Ding, Q., Dimayuga, E., Martin, S., Bruce-Keller, A. J., Nukala, V., Cuervo, A. M. & Keller, J. N. 2003 Characterization of chronic low-level proteasome inhibition on neural homeostasis. *J. Neurochem.* **86**, 489–497.
- Duijn, C. M. V. et al. 2001 PARK7, a novel locus for autosomal recessive early-onset Parkinsonism, on chromosome 1p36. *Am. J. Hum. Genet.* **69**, 629–634. (doi:10.1086/322996)
- El Agnaf, O. M., Jakes, R., Curran, M. D. & Wallace, A. 1998 Effects of the mutations Ala30 to Pro and Ala 53 to Thr on the physical and morphological properties of α -synuclein protein implicated in Parkinson's disease. *FEBS Lett.* **440**, 67–70. (doi:10.1016/S0014-5793(98)01419-7)
- Faccio, L., Fusco, C., Chen, A., Martinotti, S., Bonventre, J. V. & Zervos, A. S. 2000 Characterization of a novel human serine protease that has extensive homology to bacterial heat shock endoprotease HtrA and is regulated by kidney ischemia. *J. Biol. Chem.* **275**, 2581–2588. (doi:10.1074/jbc.275.4.2581)
- Farrer, M. et al. 2001 Lewy bodies and Parkinsonism in families with parkin mutations. *Ann. Neurol.* **50**, 293–300. (doi:10.1002/ana.1132)
- Farrer, M. et al. 2004 Comparison of kindreds with Parkinsonism and α -synuclein genomic multiplications. *Ann. Neurol.* **55**, 174–179. (doi:10.1002/ana.10846)
- Fortin, D. L., Troyer, M. D., Nakamura, K., Kubo, S., Anthony, M. D. & Edwards, R. 2004 Lipid rafts mediate the synaptic localization of α -synuclein. *J. Neurosci.* **24**, 6715–6723. (doi:10.1523/JNEUROSCI.1594-04.2004)
- Fredenburg, R. A., Rospighosi, C., Meray, R. K., Kessler, J., Lashuel, H. A., Elizer, D. & Lansbury Jr, P. T. 2007 The impact of the E46K mutation on the properties of α -synuclein, its monomeric and oligomeric states. *Biochemistry* **46**, 7107–7118. (doi:10.1021/bi7000246)
- Fuchs, J. et al. 2007 Phenotypic variation in a large Swedish pedigree due to SNCA duplication and triplication. *Neurology* **68**, 1–7. (doi:10.1212/01.wnl.0000254458.17630.c5)
- Funayama, M., Hasegawa, K., Kowa, H., Saito, M., Tsuji, S. & Obata, F. 2002 A new locus for Parkinson's disease (PARK 8) maps to chromosome 12p11.2–q13.1. *Ann. Neurol.* **51**, 296–301. (doi:10.1002/ana.10113)
- Funayama, M., Hasegawa, K., Ohta, E., Kawashima, N., Komiyama, M., Kowa, H., Tsuji, S. & Obata, F. 2005 An LRRK2 mutation as a cause for the Parkinsonism in the original PARK8 family. *Ann. Neurol.* **57**, 918–921. (doi:10.1002/ana.20484)
- Funayama, M. et al. 2007 Leucine-rich repeat kinase 2 G2385R variant is a risk factor for Parkinson disease in Asian population. *Neuroreport* **18**, 273–275. (doi:10.1097/WNR.0b013e32801254b6)
- Fung, H. C., Chen, C. M., Hardy, J., Singleton, A. B., Chen, G. J. L. & Wu, Y. R. 2006 Analysis of the *PINK1* gene in a cohort of patients with sporadic early-onset Parkinsonism in Taiwan. *Neurosci. Lett.* **394**, 33–36. (doi:10.1016/j.neulet.2005.10.005)
- Gasser, T. et al. 1998 A susceptibility locus for Parkinson's disease maps to chromosome 2p13. *Nat. Genet.* **18**, 262–265. (doi:10.1038/ng0398-262)
- Gilks, W. P. et al. 2005 A common LRRK2 mutation in idiopathic Parkinson's disease. *Lancet* **365**, 415–416.
- Gloeckner, C. J., Kinkl, N., Schumacher, A., Braum, R., O'Neill, E., Meitinger, T., Kolch, W., Prokisch, H. & Ueffing, M. 2006 The Parkinson disease causing LRRK2 mutation I2020T is associated with increased kinase activity. *Hum. Mol. Genet.* **15**, 223–232. (doi:10.1093/hmg/ddi439)
- Goker-Alpan, O., Schiffmann, R., LaMarca, M. E., Nussbaum, R. L., McInerney-Leo, A. & Sidransky, E. 2004 Parkinsonism among Gaucher disease carriers. *J. Med. Genet.* **41**, 937–940. (doi:10.1136/jmg.2004.024455)
- Goker-Alpan, O., Giasson, B. I., Eblan, M. J., Nguyen, E. J., Hurtig, H. I., Lee, V. M.-Y., Trojanowski, J. Q. & Sidransky, E. 2006 Glucocerebrosidase mutations are important risk factor for Lewy body disorders. *Neurology* **67**, 908–910. (doi:10.1212/01.wnl.0000230215.41296.18)

- Golbe, L. I., Di Iorio, G., Bonavita, V., Miller, D. C. & Duvoisin, R. C. 1990 A large kindred with autosomal dominant Parkinson's disease. *Ann. Neurol.* **27**, 276–282. (doi:10.1002/ana.410270309)
- Goldberg, M. S. *et al.* 2003 Parkin-deficient mice exhibit nigrostriatal deficits but not loss of dopaminergic neurons. *J. Biol. Chem.* **278**, 43 628–43 635. (doi:10.1074/jbc.M308947200)
- Hague, S. *et al.* 2003 Early-onset Parkinson's disease caused by a compound heterozygous DJ-1 mutation. *Ann. Neurol.* **54**, 271–274. (doi:10.1002/ana.10663)
- Hampshire, D. J., Roberts, E., Crow, Y., Hoad, J., Mubaidin, A., Wriekat, A. L., Al-Din, A. & Woods, C. G. 2001 Kufor-Rakeb syndrome, pallido-pyramidal degeneration with supranuclear upgaze paresis and dementia, maps to 1p 36. *J. Med. Genet.* **38**, 680–682. (doi:10.1136/jmg.38.10.680)
- Hatano, Y. *et al.* 2004 Novel PINK1 mutations in early-onset Parkinsonism. *Ann. Neurol.* **56**, 424–427. (doi:10.1002/ana.20251)
- Hatano, T., Kubo, S., Imai, S., Maeda, M., Ishikawa, K., Mizuno, Y. & Hattori, N. 2007 Leucine-rich repeat kinase 2 associates with lipid rafts. *Hum. Mol. Genet.* **16**, 678–690. (doi:10.1093/hmg/ddm013)
- Hattori, N., Tanaka, M., Ozawa, T. & Mizuno, Y. 1991 Immunohistochemical studies on complex I, II, III, and IV of mitochondria in Parkinson's disease. *Ann. Neurol.* **30**, 563–571. (doi:10.1002/ana.410300409)
- Hattori, N. *et al.* 1998 Molecular analysis of a novel ubiquitin-like protein (*PARKIN*) gene in Japanese families with AR-JP, evidence of homozygous deletions in the *parkin* gene in affected individuals. *Ann. Neurol.* **44**, 935–941. (doi:10.1002/ana.410440612)
- Healy, D. G. *et al.* 2004 PINK1 (*PARK6*) associated Parkinson disease in Ireland. *Neurology* **63**, 1486–1488.
- Hedrich, K. *et al.* 2002 Evaluation of 50 probands with early-onset Parkinson's disease for parkin mutations. *Neurology* **58**, 1239–1246.
- Hedrich, K. *et al.* 2004 Distribution, type, and origin of parkin mutations, review and case studies. *Mov. Disord.* **19**, 1146–1157. (doi:10.1002/mds.20234)
- Hering, R. *et al.* 2004 Novel homozygous p.E64D mutation in DJ1 in early onset Parkinson disease (*PARK7*). *Hum. Mutat.* **24**, 321–329. (doi:10.1002/humu.20089)
- Hernandez, D. G. *et al.* 2005 Clinical and positron emission tomography of Parkinson's disease caused by LRRK2. *Ann. Neurol.* **57**, 453–456. (doi:10.1002/ana.20401)
- Hicks, A. A. *et al.* 2002 A susceptibility gene for late-onset idiopathic Parkinson's disease. *Ann. Neurol.* **52**, 549–555. (doi:10.1002/ana.10324)
- Hughes, A. J., Daniel, S. E., Kilford, L. & Lees, A. J. 1992 Accuracy of clinical diagnosis of idiopathic Parkinson's disease, a clinico-pathological study of 100 cases. *J. Neurol. Neurosurg. Psych.* **55**, 181–184.
- Ibanez, P., Bonnet, A. M., Debarges, B., Lohmann, E., Tison, F., Pollak, P., Agid, Y., Durr, A. & Brice, A. 2004 Causal relation between α -synuclein gene duplication and familial Parkinson's disease. *Lancet* **364**, 1169–1171. (doi:10.1016/S0140-6736(04)17104-3)
- Imai, Y., Soda, M., Inoue, H., Hattori, N., Mizuno, Y. & Takahashi, R. 2001 An unfolded putative transmembrane polypeptide, which can lead to endoplasmic reticulum stress, is a substrate of Parkin. *Cell* **105**, 891–902. (doi:10.1016/S0092-8674(01)00407-X)
- Imai, Y., Soda, M., Hatakeyama, S., Akagi, T., Hashikawa, T., Nakayama, K. I. & Takahashi, R. 2002 CHIP is associated with *Parkin*, a gene responsible for familial Parkinson's disease, and enhances its ubiquitin ligase activity. *Mol. Cell* **10**, 55–67. (doi:10.1016/S1097-2765(02)00583-X)
- Ito, G., Okai, T., Fujino, G., Takeda, K., Ichijo, H., Kanada, T. & Iwatsubo, T. 2007 GTP binding is essential to the protein kinase activity of LRRK2, a causative gene product for familial Parkinson's disease. *Biochemistry* **46**, 1380–1382. (doi:10.1021/bi061960m)
- Javitch, J. A., D'Amato, R. J., Strittmatter, S. M. & Snyder, S. H. 1985 Parkinsonism-inducing neurotoxin, *N*-methyl-4-phenyl-1,2,3,6-tetrahydropyridine, uptake of the metabolite *N*-methyl-4-phenylpyridine by dopamine neurons explains selective toxicity. *Proc. Natl Acad. Sci. USA* **82**, 2173–2177. (doi:10.1073/pnas.82.7.2173)
- Kachergus, J. *et al.* 2005 Identification of a novel LRRK2 mutation linked to autosomal dominant Parkinsonism, evidence of a common founder across European populations. *Am. J. Hum. Genet.* **76**, 672–680. (doi:10.1086/429256)
- Kalia, S. K. *et al.* 2004 BAG5 Inhibits parkin and enhances dopaminergic neuron degeneration. *Neuron* **44**, 931–945. (doi:10.1016/j.neuron.2004.11.026)
- Kann, M. *et al.* 2002 Role of parkin mutations in 111 community-based patients with early onset Parkinsonism. *Ann. Neurol.* **51**, 621–625. (doi:10.1002/ana.10179)
- Khan, N. L., Brooks, D. J., Pavese, N., Sweeney, M. G., Wood, N. W., Lees, A. J. & Piccini, P. 2002 Progression of nigrostriatal dysfunction in a parkin kindred, an [18F]dopa PET and clinical study. *Brain* **125**, 2248–2256. (doi:10.1093/brain/awf237)
- Khan, N. L., Graham, E., Critchley, P., Schrag, A. E., Wood, N. W., Lees, A. J., Bhatia, K. P. & Quinn, N. 2003 Parkin disease, a phenotypic study of a large case series. *Brain* **126**, 1279–1292. (doi:10.1093/brain/awg142)
- Kitada, T., Asakawa, S., Hattori, N., Matsumine, H., Yamamura, Y., Minoshima, S., Yokochi, M., Mizuno, Y. & Shimizu, N. 1998 Deletion mutation in a novel protein “Parkin” gene causes autosomal recessive juvenile Parkinsonism (AR-JP). *Nature* **392**, 605–608. (doi:10.1038/33416)
- Klein, C. *et al.* 2000 Parkin deletions in a family with adult-onset, tremor-dominant Parkinsonism, expanding the phenotype. *Ann. Neurol.* **48**, 65–71. (doi:10.1002/1531-8249(200007)48:1<65::AID-ANA10>3.0.CO;2-L)
- Kono, S. *et al.* 2007 Dopaminergic neuronal dysfunction associated with Parkinsonism in both a Gaucher disease patient and a carrier. *J. Neurol. Sci.* **252**, 181–184. (doi:10.1016/j.jns.2006.10.019)
- Ko, H. S. *et al.* 2005 Accumulation of the authentic parkin substrate aminoacyl-tRNA synthase cofactor, p38/JTV-1, leads to catecholaminergic cell death. *Neurobiol. Dis.* **25**, 7968–7978.
- Ko, H. S., Kim, S. W., Sriram, S. R., Dawson, V. L. & Dawson, T. M. 2006 Identification of far upstream element-binding protein-1 as an authentic parkin substrate. *J. Biol. Chem.* **281**, 16193–16196. (doi:10.1074/jbc.C600041200)
- Krüger, R. *et al.* 1998 Ala30Pro mutation in the gene encoding α -synuclein in Parkinson's disease. *Nat. Genet.* **18**, 106–108. (doi:10.1038/ng0298-106)
- Kubo, S., Nemani, V. M., Chalkley, R. J., Anthony, M. D., Hattori, N., Mizuno, Y., Edwards, R. H. & Frotin, D. L. 2005 A combinatorial code for the interaction of α -synuclein with membranes. *J. Biol. Chem.* **280**, 31664–31672. (doi:10.1074/jbc.M504894200)
- Langston, J. W., Ballard, P., Tetrad, J. W. & Irwin, I. 1983 Chronic Parkinsonism in humans due to a product of meperidine-analog synthesis. *Science* **219**, 979–980. (doi:10.1126/science.6823561)
- Leroy, E. *et al.* 1998 The ubiquitin pathway in Parkinson's disease. *Nature* **395**, 451–452. (doi:10.1038/26652)

- Lewy, F. H. 1912 Paralysis agitans, I. Pathologische Anatomie, in Handbuch der Neurologie, herausgegeben von Lewndowsky, 3ter Band, Spezielle Pathologie IIs, pp. 920–933. Berlin, Germany: Springer.
- Li, J. *et al.* 1997 PTEN, a putative protein tyrosine phosphatase gene mutated in human brain, breast, and prostate cancer. *Science* **275**, 1876–1878. (doi:10.1126/science.275.5308.1943)
- Li, Y. *et al.* 2005 Clinicogenetic study of PINK1 mutations in autosomal recessive early-onset Parkinsonism. *Neurology* **64**, 1955–1957. (doi:10.1212/01.WNL.0000164009.36740.4E)
- Lim, K. L. *et al.* 2005 Parkin mediates nonclassical, proteasomal-independent ubiquitination of synphilin-1, implications for Lewy body formation. *Neurobiol. Dis.* **25**, 2002–2009.
- Lorick, K. L., Jensen, J. P., Fang, S., Ong, A. M., Hatakeyama, S. & Weissman, A. 1999 RING fingers mediate ubiquitin-conjugating enzyme (E2)-dependent ubiquitination. *Proc. Natl Acad. Sci. USA* **96**, 11 364–11 369. (doi:10.1073/pnas.96.20.11364)
- Lwin, A., Orvisky, E., Goker-Alpan, O., LaMarca, M. E. & Sidransky, E. 2004 Glucocerebrosidase mutations in subjects with Parkinsonism. *Mol. Genet. Metab.* **81**, 70–73. (doi:10.1016/j.ymgme.2003.11.004)
- Machida, Y. *et al.* 2005 Common anti-apoptotic roles of parkin and α -synuclein in human dopaminergic cell. *Biochem. Biophys. Res. Commun.* **332**, 233–240. (doi:10.1016/j.bbrc.2005.04.124)
- Maroteaux, L., Campanelli, J. T. & Scheller, R. H. 1988 Synuclein, a neuron-specific protein localized to the nucleus and presynaptic nerve terminal. *J. Neurosci.* **8**, 2804–2815.
- Mata, I. F., Wedemeyer, W. J., Farrer, M. J., Taylor, J. P. & Gallo, K. A. 2006 LRRK2 in parkinson's disease, protein domains and functional insights. *Trends Neurosci.* **29**, 286–293. (doi:10.1016/j.tins.2006.03.006)
- Matsumine, H. *et al.* 1997 Localization of a gene for autosomal recessive form of juvenile Parkinsonism (AR-JP) to chromosome 6q25.2–27. *Am. J. Hum. Genet.* **60**, 588–596.
- Matsumine, H., Yamamura, Y., Hattori, N., Kobayashi, T. & Mizuno, Y. 1998 A microdeletion spanning D6S305 co-segregates with autosomal recessive juvenile Parkinsonism (ARJP). *Genomics* **49**, 143–146. (doi:10.1006/geno.1997.5196)
- McNaught, K. S., Belizaire, R., Isacson, O., Jenner, P. & Olanow, C. W. 2003 Altered proteasomal function in sporadic Parkinson's disease. *Exp. Neurol.* **179**, 38–46. (doi:10.1006/exnr.2002.8050)
- Mitsumoto, A., Nakagawa, Y., Takeuchi, A., Okawa, K., Iwamatsu, A. & Takanezawa, Y. 2001 Oxidized forms of peroxiredoxins and DJ-1 on two-dimensional gels increased in response to sublethal levels of paraquat. *Free Radic. Res.* **35**, 301–310. (doi:10.1080/10715760100300831)
- Mizuno, Y., Saitoh, T. & Sone, N. 1987a Inhibition of mitochondrial NADH-ubiquinone oxidoreductase activity by 1-methyl-4-phenylpyridinium ion. *Biochem. Biophys. Res. Commun.* **143**, 294–299. (doi:10.1016/0006-291X(87)90346-9)
- Mizuno, Y., Saitoh, T. & Sone, N. 1987b Inhibition of mitochondrial α -ketoglutarate dehydrogenase by 1-methyl-4-phenylpyridinium ion. *Biochem. Biophys. Res. Commun.* **143**, 971–976. (doi:10.1016/0006-291X(87)90346-9)
- Mizuno, Y., Ohta, S., Tanaka, M., Takamiya, S., Suzuki, K., Sato, T., Oya, H., Ozawa, T. & Kagawa, Y. 1989 Deficiencies in complex I subunits of the respiratory chain in Parkinson's disease. *Biochem. Biophys. Res. Commun.* **163**, 1450–1455. (doi:10.1016/0006-291X(89)91141-8)
- Mizuno, Y., Matuda, S., Yoshino, H., Mori, H., Hattori, N. & Ikebe, S. 1994 An immunohistochemical study on α -ketoglutarate dehydrogenase complex in Parkinson's disease. *Ann. Neurol.* **35**, 204–210. (doi:10.1002/ana.410350212)
- Mizuta, I. *et al.* 2006 Multiple candidate gene analysis identifies α -synuclein as a susceptibility gene for sporadic Parkinson's disease. *Hum. Mol. Genet.* **15**, 1–8. (doi:10.1093/hmg/ddl030)
- Mochizuki, H., Goto, G., Mori, H. & Mizuno, Y. 1996 Histochemical detection of apoptosis in Parkinson's disease. *J. Neurol. Sci.* **137**, 120–123. (doi:10.1016/0022-510X(95)00336-Z)
- Moore, D. J., Zhang, L., Troncoso, J., Lee, M. K., Hattori, N., Mizuno, Y., Dawson, T. M. & Dawson, V. L. 2005 Association of DJ-1 and parkin mediated by pathogenic DJ-1 mutations and oxidative stress. *Hum. Mol. Genet.* **14**, 71–84. (doi:10.1093/hmg/ddi007)
- Mori, H., Kondo, T., Yokochi, M., Matusmine, H., Nakagawa-Hattori, Y., Miyake, T., Suda, K. & Mizuno, Y. 1998 Pathologic and biochemical studies of juvenile Parkinsonism linked to chromosome 6q. *Neurology* **51**, 890–892.
- Mueller, J. C. *et al.* 2005 Multiple regions of alpha-synuclein are associated with Parkinson's disease. *Ann. Neurol.* **57**, 535–541. (doi:10.1002/ana.20438)
- Nagakubo, D., Taira, T., Kitaura, H., Ikeda, M., Tamai, K., Iguchi-Ariga, S. M. & Ariga, H. 1997 DJ-1, a novel oncogene which transforms mouse NIH3T3 cells in cooperation with ras. *Biochem. Biophys. Res. Commun.* **231**, 509–513. (doi:10.1006/bbrc.1997.6132)
- Najim Al-Din, A. S., Wriekat, A., Mubaidin, A., Dasouki, M. & Hiari, M. 1994 Pallido-pyramidal degeneration, supranuclear upgaze paresis and dementia, Kufor-Rakeb syndrome. *Acta Neurol. Scand.* **89**, 347–352.
- Nichols, W. C. *et al.* 2005 Genetic screening for a single common LRRK2 mutation in familial Parkinson's disease. *Lancet* **365**, 410–412.
- Nicklas, W. J., Vyas, I. & Heikkila, R. E. 1985 Inhibition of NADH-linked oxidation in brain mitochondria by 1-methyl-4-phenyl-pyridine, a metabolite of the neurotoxin, 1-methyl-4-phenyl-1,2,5,6-tetrahydropyridine. *Life Sci.* **36**, 2503–2508. (doi:10.1016/0024-3205(85)90146-8)
- Nishioka, K. *et al.* 2006 Clinical heterogeneity of α -synuclein gene duplication in Parkinson's disease. *Ann. Neurol.* **59**, 298–309. (doi:10.1002/ana.20753)
- Nukada, H., Kowa, H., Saito, T., Tasaki, Y. & Miura, S. 1978 A big family of paralysis agitans. *Rinshoshinkeigaku* **18**, 627–634.
- Oliviera, S. A. *et al.* 2003 Parkin mutations and susceptibility alleles in late-onset Parkinson's disease. *Ann. Neurol.* **53**, 624–629. (doi:10.1002/ana.10524)
- Paisan-Ruiz, C. *et al.* 2004 Cloning of the gene containing mutations that cause PARK8-linked Parkinson's disease. *Neuron* **44**, 595–600. (doi:10.1016/j.neuron.2004.10.023)
- Paisan-Ruiz, C., Saenz, A., Lopez de Munain, A., Marti, I., Martinez Gil, A., Marti-Masso, J. F. & Perez-Tur, J. 2005 Familial Parkinson's disease, clinical and genetic analysis of four Basque families. *Ann. Neurol.* **57**, 365–372. (doi:10.1002/ana.20391)
- Pankratz, N. *et al.* 2003a Genome-wide linkage analysis and evidence of gene-by-gene interactions in a sample of 362 multiplex Parkinson disease families. *Hum. Mol. Genet.* **12**, 2599–2608. (doi:10.1093/hmg/ddg270)
- Pankratz, N., Nichols, W. C., Uniacke, S. K., Halter, C., Rudolph, A., Shults, C., Conneally, P. M., Foroud, T. &

- Parkinson Study Group 2003b Significant linkage of Parkinson disease to chromosome 2q36-37. *Am. J. Hum. Genet.* **72**, 1053–1057. (doi:10.1086/374383)
- Parkinson, J. 1817 *An essay on the shaking palsy*, pp. 1–66. London, UK: Sherwood, Neely, and Jones.
- Park, J. *et al.* 2006 Mitochondrial dysfunction in *Drosophila* PINK1 mutants is complemented by Parkin. *Nature* **441**, 1157–1161. (doi:10.1038/nature04788)
- Perez, F. A. & Palmiter, R. D. 2005 Parkin-deficient mice are not a bobust model of Parkinsonism. *Proc. Natl Acad. Sci. USA* **102**, 2174–2179. (doi:10.1073/pnas.0409598102)
- Pesah, Y., Pham, T., Burgesse, H., Middlebrooks, B., Verstreken, P., Zhou, Y., Harding, M., Bellen, H. & Mardon, G. 2004 *Drosophila* parkin mutants have decreased mass and cell size and increased sensitivity to oxygen radical stress. *Development* **131**, 2183–2194. (doi:10.1242/dev.01095)
- Polymeropoulos, M. H. *et al.* 1997 Mutation in the α -synuclein gene identified in families with Parkinson's disease. *Science* **276**, 2045–2047. (doi:10.1126/science.276.5321.2045)
- Ramirez, A. *et al.* 2006 Hereditary Parkinsonism with dementia is caused by mutations in ATP13A2, encoding a lysosomal type 5 P-type ATPase. *Nat. Genet.* **38**, 1184–1191. (doi:10.1038/ng1884)
- Ramsay, R. R., Salach, J. I., Dadgar, J. & Singer, T. P. 1986 Inhibition of mitochondrial NADH dehydrogenase by pyridine derivatives and its possible relation to experimental and idiopathic Parkinsonism. *Biochem. Biophys. Res. Commun.* **135**, 269–275. (doi:10.1016/0006-291X(86)90972-1)
- Ren, Y., Zhao, J. & Feng, J. 2003 Parkin binds to α/β tubulin and increases their ubiquitination and degradation. *J. Neurosci.* **23**, 3316–3324.
- Rideout, H. J., Lang-Rollin, I. & Stefanis, I. 2004 Involvement of macroautophagy in the dissolution of neuronal inclusions. *Int. J. Biochem. Cell. Biol.* **36**, 2551–2562. (doi:10.1016/j.biocel.2004.05.008)
- Rohe, C. F., Montagna, P., Breedveld, G., Cortelli, P., Oostra, B. A. & Bonifati, V. 2004 Homozygous PINK1 C-terminus mutation causing early-onset Parkinsonism. *Ann. Neurol.* **56**, 427–431. (doi:10.1002/ana.20247)
- Rubinshtein, R. C. 2006 The role of intracellular protein-degradation pathways in neurodegeneration. *Nature* **443**, 780–786. (doi:10.1038/nature05291)
- Saigoh, K. *et al.* 1999 Intragenic deletion in the gene encoding ubiquitin carboxy-terminal hydrolase in *gad* mice. *Nat. Genet.* **23**, 47–51.
- Sakata, E. *et al.* 2003 Parkin binds the Rpn10 subunit of 26S proteasomes through its ubiquitin-like domain. *EMBO Rep.* **4**, 301–306. (doi:10.1038/sj.embor.embor764)
- Sato, C. *et al.* 2005 Analysis of the glucocerebrosidase gene in Parkinson's disease. *Mov. Disord.* **20**, 367–370. (doi:10.1002/mds.20319)
- Sato, S. *et al.* 2006a Decline of striatal dopamine release in Parkin-deficient mice shown by *ex vivo* autoradiography. *J. Neurosci. Res.* **84**, 1350–1357. (doi:10.1002/jnr.21032)
- Sato, S., Chiba, T., Sakata, E., Kato, K., Mizuno, Y., Hattori, N. & Tanaka, K. 2006b 14-3-3 η is a novel regulator of parkin ubiquitin-ligase. *EMBO J.* **25**, 211–221. (doi:10.1038/sj.emboj.7600774)
- Schapira, A. H. V., Cooper, J. M., Dexter, D., Jenner, P., Clark, J. B. & Marsden, C. D. 1989 Mitochondrial complex I deficiency in Parkinson's disease. *Lancet* **1**, 1269. (doi:10.1016/S0140-6736(89)92366-0)
- Schlitter, A. M., Kurz, M., Larsen, J. P., Voitalla, D., Mueller, T., Epplen, J. T. & Dekomien, G. 2006 *Parkin* gene variations in late-onset Parkinson's disease, comparison between Norwegian and German cohorts. *Acta Neurol. Scand.* **113**, 9–13. (doi:10.1111/j.1600-0404.2005.00532.x)
- Shimoda-Matsubayashi, S., Matsumine, H., Kobayashi, T., Nakagawa-Hattori, Y., Shimizu, Y. & Mizuno, Y. 1996 Structural dimorphism in the mitochondrial targeting sequence in the human MnSOD gene. A predictive evidence for conformational change to influence mitochondrial transport and a study of allelic association in Parkinson's disease. *Biochem. Biophys. Res. Commun.* **226**, 561–565. (doi:10.1006/bbrc.1996.1394)
- Shimura, H. *et al.* 2001 Ubiquitination of a new form of α -synuclein by parkin from human brain, implication for Parkinson's disease. *Science* **293**, 263–269. (doi:10.1126/science.1060627)
- Shimura, H. *et al.* 2000 Familial Parkinson's disease gene product, Parkin, is a ubiquitin-protein ligase. *Nat. Genet.* **25**, 302–305. (doi:10.1038/77060)
- Shimura-Miura, H., Hattori, N., Kang, D., Miyako, K., Nakabeppu, Y. & Mizuno, Y. 1999 Increased 8-oxo-dGTPase in the mitochondria of substantia nigral neurons in Parkinson's disease. *Ann. Neurol.* **46**, 920–924. (doi:10.1002/1531-8249(199912)46:6<920::AID-ANA17>3.0.CO;2-R)
- Singleton, A. B. *et al.* 2003 α -Synuclein locus triplication causes Parkinson's disease. *Science* **302**, 841. (doi:10.1126/science.1090278)
- Smith, W. W., Pei, Z., Jian, H., Moore, D., Liang, Y., West, A. B., Dawson, V. L., Dawson, T. M. & Ross, C. A. 2005 Leucine-rich repeat kinase 2 (LRRK2) interacts with parkin and mutant LRRK2 induces neuronal degeneration. *Proc. Natl Acad. Sci. USA* **102**, 18 676–18 681. (doi:10.1073/pnas.0508052102)
- Sofic, E., Lange, K. W., Jellinger, K. & Riederer, P. 1992 Reduced and oxidized glutathione in the substantia nigra of patients with Parkinson's disease. *Neurosci. Lett.* **142**, 128–130. (doi:10.1016/0304-3940(92)90355-B)
- Spillantini, M. G., Schmidt, M. L., Lee, A. M. Y., Trojanowski, J. Q., Jakes, R. & Goedert, M. 1997 α -Synuclein in Lewy bodies. *Nature* **388**, 839–840. (doi:10.1038/42166)
- Staropoli, J. F., McDermott, C., Martinat, C., Schulman, B., Demireva, E. & Abeliovich, A. 2003 Parkin is a component of an SCF-like ubiquitin ligase complex and protects postmitotic neurons from kainate excitotoxicity. *Neuron* **37**, 735–749. (doi:10.1016/S0896-6273(03)00084-9)
- Stefanis, I., Larsen, K. E., Rideout, H. J., Sulzer, D. & Greene, I. A. 2001 Expression of A53T but not wild-type α -synuclein in PC12 cells induces alterations of the ubiquitin-dependent degradation system, loss of dopamine release, and autophagic cell death. *J. Neurosci.* **21**, 9549–9560.
- Strauss, K. M. *et al.* 2005 Loss of function mutations in the gene encoding Omi/HtrA in Parkinson's disease. *Hum. Mol. Genet.* **14**, 2099–2111. (doi:10.1093/hmg/ddi215)
- Suzuki, Y., Imai, Y., Nakayama, H., Takahashi, K., Takio, K. & Takahashi, R. 2001 A serine protease, HtrA2, is released from the mitochondria and interacts with XIAP, inducing cell death. *Mol. Cell* **8**, 613–621. (doi:10.1016/S1097-2765(01)00341-0)
- Takahashi, H., Ohama, E., Suzuki, S., Horikawa, Y., Ishikawa, A., Morita, T., Tsuji, S. & Ikuta, F. 1994 Familial juvenile Parkinsonism, clinical and pathologic study in a family. *Neurology* **44**, 437–441.
- Takahashi-Niki, K., Niki, T., Taira, T., Iguchi-Ariga, S. M. & Ariga, H. 2004 Reduced anti-oxidative stress activities of DJ-1 mutants found in Parkinson's disease patients. *Biochem. Biophys. Res. Commun.* **320**, 389–397. (doi:10.1016/j.bbrc.2004.05.187)

- Takanashi, M., Mochizuki, H., Yokomizo, K., Hattori, N., Mori, H., Yamamura, Y. & Mizuno, Y. 2001 Iron accumulation in the substantia nigral of autosomal recessive juvenile Parkinsonism (ARJP). *Parkinsonism Rel. Disord.* **7**, 311–314. (doi:10.1016/S1353-8020(00)00050-X)
- Tan, E. K. et al. 2007 The LRRK2 Gly238Arg variant is associated with Parkinson's disease; genetic and functional evidence. *Hum. Genet.* **120**, 857–863. (doi:10.1007/s00439-006-0268-0)
- Tanaka, K., Suzuki, T., Hattori, N. & Mizuno, Y. 2004 Ubiquitin, proteasome and Parkin. *Biochim. Biophys. Acta* **1695**, 235–247. (doi:10.1016/j.bbamcr.2004.09.026)
- Tayebi, N. et al. 2003 Gaucher disease with Parkinsonism manifestations; does glucocerebrosidase deficiency contribute to a vulnerability to Parkinsonism? *Mol. Genet. Metab.* **79**, 104–109. (doi:10.1016/S1096-7192(03)00071-4)
- Toft, M., Myhre, R., Pielsticker, L., White, L. R., Aasly, J. O. & Farrer, M. J. 2007 PINK1 mutation heterozygosity and the risk of Parkinson's disease. *J. Neurol. Neurosurg. Psych.* **78**, 82–84. (doi:10.1136/jnnp.2006.097840)
- Trétiakoff, C. 1919 Contribution à l'étude de l'anatomie pathologique du locus niger. Thesis, University of Paris, cited in Oppenheimer, D. R. 1984 Diseases of the basal ganglia, cerebellum and motor neurons, In *Greenfield textbook of neuropathology*, 4th edn. (eds J. H. Adams, J. A. N. Corsellis & L. W. Duchon), pp. 699–747. Edward, London: Arnold.
- Tsai, Y. C., Fishman, P. S., Thakor, N. V. & Oyler, G. A. 2003 Parkin facilitates the elimination of expanded polyglutamine proteins and leads to preservation of proteasome function. *J. Biol. Chem.* **278**, 22 044–22 055. (doi:10.1074/jbc.M212235200)
- Um, J. W., Min, D. S., Rhim, H., Kim, J., Palk, S. R. & Chung, K. C. 2006 Parkin ubiquitinates and promotes the degradation of Ran BP2. *J. Biol. Chem.* **281**, 3595–3603. (doi:10.1074/jbc.M504994200)
- Valente, E. M., Bentivoglio, A. R., Dixon, P. H., Ferraris, A., Ialongo, T., Frontali, M., Albanese, A. & Wood, N. W. 2001 Localization of a novel locus for autosomal recessive early-onset Parkinsonisms, PARK6, on human chromosome 1p35–36. *Am. J. Hum. Genet.* **68**, 895–900. (doi:10.1086/319522)
- Valente, E. M. et al. 2004a Hereditary early-onset Parkinson's disease caused by mutations in PINK1. *Science* **304**, 1158–1160. (doi:10.1126/science.1096284)
- Valente, E. M. et al. 2004b PINK1 mutations are associated with sporadic early-onset Parkinsonism. *Ann. Neurol.* **56**, 336–341. (doi:10.1002/ana.20256)
- Ved, R. et al. 2005 Similar patterns of mitochondrial vulnerability and rescue induced by genetic modification of α -synuclein, parkin and DJ-1 in *Cernorhabditis elegans*. *J. Biol. Chem.* **280**, 42 655–42 668. (doi:10.1074/jbc.M505910200)
- Volles, M. J. & Lansbury Jr, P. T. 2002 Vesicle permeabilization by protofibrillar α -synuclein is sensitive to Parkinson's disease-linked mutations and occurs by a pore-like mechanism. *Biochemistry* **41**, 4595–4602. (doi:10.1021/bi0121353)
- von Coelln, R., Thomas, B., Savitt, J. M., Lim, K. L., Sasaki, M., Hess, E. J., Dawson, V. L. & Dawson, T. M. 2004 Loss of locus coeruleus neurons and reduced startle in parkin null mice. *Proc. Natl Acad. Sci. USA* **101**, 10 744–10 749. (doi:10.1073/pnas.0401297101)
- Webb, J. L., Ravikumar, B., Atkins, J., Skepper, J. N. & Rubinsztein, D. C. 2003 α -Synuclein is degraded by both autophagy and the proteasome. *J. Biol. Chem.* **278**, 25 009–25 013. (doi:10.1074/jbc.M300227200)
- Weinreb, P. H., Zhen, W., Poon, A. W., Conway, K. A. & Lansbury Jr, P. T. 1996 NACP, a protein implicated in Alzheimer's disease and learning, is natively unfolded. *Biochemistry* **35**, 13 709–13 715. (doi:10.1021/bi961799n)
- West, A. et al. 2002 Complex relationship between parkin mutations and Parkinson disease. *Am. J. Med. Genet.* **114**, 584–591. (doi:10.1002/ajmg.10525)
- Williams, D. R., Hadeed, A., Al-Din, A. S. M., Wreikat, A. L. & Lees, A. J. 2005 Kufor Rakeb disease, autosomal recessive, levodopa-responsive Parkinsonism with pyramidal degeneration, supranuclear gaze palsy, and dementia. *Mov. Disord.* **20**, 1264–1271. (doi:10.1002/mds.20511)
- Wszolek, Z. K. et al. 1997 German–Canadian Family (Family A) with Parkinsonism, amyotrophy, and dementia, longitudinal observation. *Parkinsonism Rel. Disord.* **3**, 125–136. (doi:10.1016/S1353-8020(97)00013-8)
- Wszolek, Z. K. et al. 2004 Autosomal dominant Parkinsonism associated with variable synuclein and tau pathology. *Neurology* **62**, 1619–1622.
- Yamamura, Y., Sobue, I., Ando, K., Iida, M., Yanagi, T. & Kono, C. 1973 Paralysis agitans of early onset with marked diurnal fluctuation of symptoms. *Neurology* **23**, 239–244.
- Yokota, T., Sugawara, K., Ito, K., Takahashi, R., Ariga, H. & Mizusawa, H. 2003 Down regulation of DJ-1 enhances cell death by oxidative stress, ER stress, and proteasome inhibition. *Biochem. Biophys. Res. Commun.* **312**, 1342–1348. (doi:10.1016/j.bbrc.2003.11.056)
- Yorimitsu, T. & Klionsky, D. J. 2005 Autophagy: molecular machinery for self-eating. *Cell Death Differ* **12**(Suppl 2), 1542–1552.
- Yoritaka, A., Hattori, N., Uchida, K., Tanaka, M., Stadtman, E. R. & Mizuno, Y. 1996 Immunohistochemical detection of 4-hydroxynonenal protein adducts in Parkinson's disease. *Proc. Natl Acad. Sci. USA* **93**, 2696–2701. (doi:10.1073/pnas.93.7.2696)
- Youdim, M. B. H., Ben-Shachar, D. & Riederer, P. 1989 Is Parkinson's disease a progressive siderosis of substantia nigra resulting in iron and melanin induced neurodegeneration? *Acta Neurol. Scand.* **126**, 47–54.
- Zarranz, J. J. et al. 2004 The new mutation, E46K, of α -synuclein causes Parkinson's disease and Lewy body dementia. *Ann. Neurol.* **55**, 164–173. (doi:10.1002/ana.10795)
- Zhang, Y., Gao, J., Chung, K. K., Huang, H., Dawson, V. L. & Dawson, T. M. 2000 Parkin functions as an E2 dependent ubiquitin-protein ligase and promotes the degradation of the synaptic vesicle associated protein, CDCrel-1. *Proc. Natl Acad. Sci. USA* **21**, 13 354–13 359. (doi:10.1073/pnas.240347797)
- Zhong, L., Tan, Y., Zhou, A., Yu, Q. & Zhou, J. 2005 RING finger ubiquitin-protein isopeptide ligase Nrdp1/FLRF regulates parkin stability and activity. *J. Biol. Chem.* **280**, 9425–9430. (doi:10.1074/jbc.M408955200)
- Zimprich, A. et al. 2004 Mutations in LRRK2 cause autosomal-dominant Parkinsonism with pleomorphic pathology. *Neuron* **44**, 601–607. (doi:10.1016/j.neuron.2004.11.005)