Published in final edited form as:

Bioorg Med Chem Lett. 2008 April 15; 18(8): 2555–2557. doi:10.1016/j.bmcl.2008.03.044.

Design of High Affinity Peptide Conjugates with Optimized Fluorescence Quantum Yield as Markers for Small Peptide Transporter PEPT1 (SLC15A1)

Praveen M. Bahadduri, Abhijit Ray, Akash Khandelwal, and Peter W. Swaan*

Department of Pharmaceutical Sciences, School of Pharmacy, University of Maryland, Baltimore, MD 21201, USA

Abstract

We employed a computational approach to design and synthesize a series of fluorescently labeled hPEPT1 substrates. Five Alexa Fluor-350™ labeled peptides were assessed for their *in vitro* inhibitory activity in hPEPT1-transfected CHO cells. At least four labeled peptides show potent inhibitory activity towards hPEPT1-mediated uptake of [³H]-GlySar and three compounds displayed significant cellular uptake specifically mediated by hPEPT1.

Keywords

Alexa Fluor 350™; hPEPT1; Peptide conjugation; Fluorescence; High-throughput screening; CoMFA

The human small peptide transporter, hPEPT1, is a clinically important intestinal solute carrier protein critically involved in the absorption of its natural substrates, food protein-derived di- and tripeptides. Additionally, it mediates the intestinal uptake of numerous drug classes, such as beta-lactam antibiotics, ACE-inhibitors, several antivirals,¹ as well as various prodrugs. For example, valyl prodrugs of the nucleoside antiviral drugs acyclovir and ganciclovir display 3-5 fold and 10-fold higher bioavailability upon oral administration in humans compared to their parent compounds, respectively.^{2, 3} The increased intestinal permeability can be attributed solely to their affinity for intestinal hPEPT1. This transporter's relaxed substrate specificity, relatively high capacity and low affinity ($K_m \sim 1.1$ mM for the prototypical PEPT1 substrate glycylsarcosine (GlySar)) makes it an ideal target for the design of peptidomimetic drugs with increased intestinal permeability and bioavailability.

In the absence of a crystal structure for hPEPT1, the rate at which new substrates and inhibitors can be identified or rationally designed is expectedly slow. Alternative experimental strategies to rapidly screen large compound database for hPEPT1 affinity can be carried out using stably expressed cell lines in a high-throughput screening (HTS) setup. The success of this approach will be determined by assay sensitivity for prototypical substrates or new chemical entities (NCEs) when assessing inhibition or transport via hPEPT1. The most commonly applied assay techniques involve laborious liquid chromatography-mass spectrometry or costly radiolabeled

*Corresponding Author: Tel.: +1 410 706 0103; Fax.: +1 410 706 5017; e-mail: pswaan@rx.umaryland.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

tracer methodologies. However, fluorescence offers a convenient alternative for HTS applications due to its sensitivity and applicability to a wide variety of cell-based systems.

Several other groups have reported fluorescently labeled peptides directed to develop hPEPT1 HTS assays, albeit with limited success. Abe and colleagues⁴ synthesized FITC and coumarin-3-carboxylic acid conjugated peptide analogs that showed affinity for hPEPT1 expressed in Caco2 cells, but were not transported. Other studies reported AMCA (7-amino-4-methylcoumarin-3-acetic acid) peptide conjugates⁵ and peptide analog-FITC esters⁶ with hPEPT1 affinity. However, the fluorescent dyes used in these studies have limited quantum yields and may not be particularly suitable for HTS implementation.⁷ In fact, fluorescein dyes are known to display self-quenching properties at high concentrations, thereby limiting potential assay sensitivity. The present study was designed to develop a metabolically stable fluorogenic substrate for hPEPT1 with superior fluorescence quantum yield at a range of intracellular pH values. Based on our previous knowledge of the hPEPT1 pharmacophore and structure-activity relationship,⁸ we designed five novel Alexa Fluor 350™ (AF)-labeled peptide analogs using computational design, synthesis and *in vitro* analysis of AF labeled substrates designed specifically to target PEPT1.

We built a three-dimensional quantitative structure activity relationship (3D-QSAR) using Comparative Molecular Field Analysis (CoMFA) from the extensive dataset by Gebauer and colleagues⁹. Molecules were built and subsequently energy minimized in SYBYL (v.7.1) using the Tripos force field.¹⁰ The most active compound in the dataset, *o*-(methylphenyl)-D-tyrosyl-L-alanine⁹ was used as a template and all the other molecules were superimposed to it using atom-based alignment. The resulting CoMFA steric and electrostatic contour plots are shown in Figure 1. The model developed has good descriptive ability as evidenced by a r^2 value of 0.989 with six components.

The overall steric contribution (53%) to binding is higher than the electrostatic contribution (47%). The AF analogs (Table 1) using pK_i ($-\log K_i$) (M) as dependent variable and CoMFA descriptors as independent variables in the partial least square (PLS) analysis (data not shown). Out of a set of 50 permutations, the 5 highest scoring compounds were selected for synthesis and *in vitro* testing.

The synthesis of dipeptide conjugated AF (Molecular Probes, Eugene, OR) was carried out by dissolving BocLys(Z)OSu in a mixture of N,N-dimethyl formamide (DMF), 1,4-dioxane, diisopropylamine (DIEA), and an appropriate amino acid or dipeptide (Scheme 1). The progress of the reaction was monitored by mass spectroscopy for the disappearance of a peak at m/z 478.5. The dipeptide is then deprotected by catalytic hydrogenation using 10% Pd/C to remove the carbonyloxybenzyl (Z) group. The deprotected compound was dissolved in DMF, dioxane and diisopropylethylamine followed by the addition of AF carboxylic acid succinimidyl ester to give the fluorescently labeled dipeptide. This was finally deprotected using TFA in dichloromethane to give the desired product. All *in vitro* studies were conducted with stably transfected CHO cell lines expressing hPEPT1 as described previously⁸. Competitive inhibition studies were performed at pH 6.0 for 30 mins using [³H]-glycylsarcosine (GlySar) (1 μ M, 1Ci/mmol) (Moravek, Brea, CA) as a tracer. Initially, parent peptides were assessed for their inhibitory potencies of [³H]-GlySar uptake in hPEPT1-CHO cells. Subsequently, the inhibition of 100 μ M and 1000 μ M AF-labeled peptides (in DMSO, not exceeding 1% final DMSO concentration) on [³H]-GlySar (1 μ M) accumulation in hPEPT1-CHO cells was determined. To determine the cellular uptake of fluorescently labeled peptides and to assess their mutual competitive inhibition of GlySar, the effect of 10mM GlySar (Sigma, St. Louis, MO) on the uptake of 100 μ M AF-labeled dipeptide was assayed in hPEPT1-CHO cells. Cells were washed three times and lysed with 1N NaOH for 1h and neutralized with 10% HCl solution. Total cellular fluorescence was determined using a Spectramax Gemini XS

(Molecular Devices, Sunnyvale, CA). Data were calibrated using fluorescence standard curves and normalized to protein content.

AF-labeled peptides have strong affinity for hPEPT1 as demonstrated by their potent inhibition of [^3H]-GlySar uptake in hPEPT1-transfected CHO cells (Fig. 2). Reduction in hPEPT1 affinity imparted by AF-conjugation is minimal as compared to the activity of the parent peptide (Fig. 2). Of the compounds tested, the LysVal, LysSer and LysAla analogs showed the strongest relative inhibition of [^3H]-GlySar uptake. Interestingly, the cellular uptake of AF-labeled LysVal was significantly greater (2-3 fold) compared to the other peptides (Fig. 3). This would indicate that inhibition data do not correlate with hPEPT1-mediated substrate translocation. Inhibition of AF-labeled peptide uptake by a 100-fold GlySar excess indicates mutual inhibition between GlySar and AF-labeled analogs, indicative of a shared translocation pathway, i.e. hPEPT1 (Fig. 3). Interestingly, the LysVal conjugate of AF displays relatively high cellular uptake while only modestly inhibiting [^3H]-GlySar uptake (~40% at 1 mM), whereas LysAla and LysSer conjugates exerted stronger [^3H]-GlySar uptake inhibition (<~22%) with only modest cellular permeation. This would suggest that AF-LysVal is a good substrate for PEPT1 compared to the other dipeptide conjugates, but further studies are warranted to fully characterize this phenomenon.

Overall, AF offers significant advantages over other reported fluorescently labeled peptides such as a large Stokes shift (100nm for AF vs. 30 nm for FITC) and a relatively high fluorescence quantum yield (0.89 AF-LysVal vs. AMCA reference in water), which may enable the use of these novel conjugates in a HTS setting. In summary, this report describes the design and synthesis of five novel AF-conjugated peptides with sub-mM affinity levels for hPEPT1. Future studies should determine whether cellular uptake can be monitored without cellular disruption and in real-time to facilitate HTS screening.

Acknowledgements

This research was funded by a grant from the National Institutes of Health R03 DK075157 (to P.W.S.) in response to the Molecular Libraries and Imaging Roadmap Initiative.

References and notes

1. Swaan PW, Tukker JJ. J Pharm Sci 1997;86:596. [PubMed: 9145385]
2. Jung D, Dorr A. J Clin Pharmacol 1999;39:800. [PubMed: 10434231]
3. Weller S, Blum MR, Doucette M, Burnette T, Cederberg DM, de Miranda P, Smiley ML. Clin Pharmacol Ther 1993;54:595. [PubMed: 8275615]
4. Abe H, Satoh M, Miyauchi S, Shuto S, Matsuda A, Kamo N. Bioconjug Chem 1999;10:24. [PubMed: 9893960]
5. Groneberg DA, Doring F, Eynott PR, Fischer A, Daniel H. Am J Physiol Gastrointest Liver Physiol 2001;281:G697. [PubMed: 11518682]
6. Landowski CP, Han HK, Lee KD, Amidon GL. Pharm Res 2003;20:1738. [PubMed: 14661916]
7. Leung WY, Trobridge PA, Haugland RP, Haugland RP, Mao F. Bioorganic & medicinal chemistry letters 1999;9:2229. [PubMed: 10465551]
8. Ekins S, Johnston JS, Bahadduri P, D'Souza VM, Ray A, Chang C, Swaan PW. Pharm Res 2005;22:512. [PubMed: 15846457]
9. Gebauer S, Knutter I, Hartrodt B, Brandsch M, Neubert K, Thondorf I. J Med Chem 2003;46:5725. [PubMed: 14667225]
10. Ekins S, Swaan PW. Rev Comp Chem 2004;20

Figure 1.

CoMFA contour map. The green and yellow regions indicate that an increase in steric bulk favors or disfavors binding to hPEPT1, respectively. The increase of negative charge near red region or positive charge near blue region favors binding. The template compound is shown in capped stick model.

Fig. 2. Relative inhibition of 1 μM [³H]-GlySar uptake (% control) in hPEPT1-transfected CHO cells in the presence of fluorescently labeled peptides (n=3 ±s.d.).

Fig. 3. AF-labeled peptide uptake in hPEPT1-CHO cells. Bars indicate cellular uptake of 100μM labeled peptide in the absence (open bars) and presence (closed bars) of 10mM GlySar. (n=3 ± s.d.)

$R = H, CH_3, CH_2OH, CH(CH_3)_2$

Scheme 1.

i. DMF, 1,4-Dioxane, DIEA; ii. 10% Pd/C, EtOAc, MeOH; iii. DMF, 1,4-Dioxane, Alexa Fluor 350 carboxylic acid succinimidyl ester; iv. CH_2Cl_2 , TFA

Table 1

Predicted activities of Alexa Fluor-350™ derivatives from CoMFA analysis.

Compounds	-log <i>K_i</i>	<i>K_i</i> (mM)
AF-LysGly	4.550	0.028
AF-LysVal	4.459	0.035
AF-LysSer	4.590	0.026
AF-LysAla	4.611	0.024
AF-LysGlyGly	4.700	0.019