who was languid and anæmic, with a spleen reaching half way to his umbilicus, and decided excess of leucocytes; he improved under the use of arsenic, and I lost sight of him. I have lately had an opportunity of examining him, and found him quite well. Hæmorrhage, high fever, diarrhoea, vomiting, and œdema, deficient urea, and the presence of acetone may be looked upon as signs of grave prognostic significance. Death may occur from general asthenia, from coma of Kussmaul's type closely resembling that which occurs in diabetes, from cerebral hæmorrhage, or from some intercurrent affection. The tissues *post mortem* may be found to be infiltrated with micro-organisms, suggesting that their resistance to ordinary infections had been greatly reduced.

TREATMENT.

The treatment should be designed to place the patient under the most favourable hygienic conditions—in a healthy bracing climate on a dry soil, with good air, good food, and favourable surroundings—and he should if possible be free from all sources of worry or mental emotion. The most valuable drug is arsenic, which should be commenced in small doses and gradually increased. All the above indications may probably be best fulfilled by sending the patient to undergo a course of arsenical waters at La Bourboule, in France (Department Puy-de-Dôme). Cold douches with the galvanic and faradic currents may possibly be useful. Quinine and iron are often employed, and oxygen inhalations may be tried, but are probably only resorted to in the later and more hopeless stages. The operation of splenectomy is hardly justified from our present pathological standpoint as a rational proceeding; but it has also against it the terrible objection that, out of twenty-four operations, only one patient has recovered. The chief danger seems to be that in leukæmic patients the blood has lost the power by which normal hæmostasis occurs, and the cut surfaces continue to bleed from vessels which offer nothing that the surgeon can tie. Transfusion appears to be useless. Ewart has recommended the inhalation of carbonic acid gas, but no one has confirmed his favourable opinion of this treatment. The enlarged spleen may be supported by a bandage or belt, and the stomach troubles are relieved by the administration of small doses of dilute mineral acid after meals.

AN ADDRESS ON ACUTE DILATATION OF THE HEART IN DIPHTHERIA, INFLUENZA, AND RHEUMATIC FEVER.

Delivered before the Manchester Medical Society.

By D. B. LEES, M.D., M.A., B.Sc., F.R.C.P.,

Senior Physician to the Hospital for Sick Children, Great Ormond Street; Physician to St. Mary's Hospital; Examiner in Medicine for the University of Cambridge and for the Victoria University.

SUDDEN DEATH DURING AND AFTER DIPHTHERIA.

GENTLEMEN,—It is a fact of common knowledge that sudden death, often quite unexpected, is sometimes the result of an attack of diphtheria in a child. Most medical practitioners of ten years' standing have seen or heard of a case in which this has happened. Let me briefly refer to a few instances which I have myself met with, or which have been related to me by trustworthy observers.

CASE I.—When I was a clinical clerk, a young girl was admitted into hospital one afternoon on account of weakness after diphtheria. She sat up in bed and did not seem ill. When I came next morning to take notes of her condition she was dead.

CASE II.—Some years after this, when I was in charge of out-patients at St. Mary's Hospital, I admitted under my own care (by the kindness of Sir William Broadbent) a child suffering from diphtherial paralysis. I saw it in the ward the same afternoon, and perceived no cause for apprehension. When I came to visit it next morning it was dead.

CASE III.—A child of 3 years of age had suffered from diphtheria, but was thought to be convalescent. She was standing by a window when some one entered the room a little abruptly. She turned quickly round and fell dead on the floor.

CASE IV.—A medical man in good practice attended a child for diphtheria. As she did not recover as quickly as he expected he sent her to the seaside for change of air. She fell dead on the sands.

CASES V AND VI.—Another medical man had two little patients suffering from diphtheria. When one of them seemed convalescent he allowed it to get up. The child was dead in an hour. Warned by this, he kept the

second child in bed for a week longer; it was then allowed to rise. Within twenty-four hours this child also died.

CASE VII.—A girl of nearly 11 years old had a severe attack of diphtheria. She looked pale after this and was kept in bed for eight weeks. She was then allowed to rise. In five minutes after rising she was dead.

Occurrences of this kind, so painful both to the surviving relatives and to the practitioner who has given a hopeful prognosis, call for careful study to see if we can detect any indication which will warn us of the coming danger and enable us to guard against it.

DUE TO DEGENERATION OF THE HEART MUSCLE.

On what do they depend? The first explanation that suggests itself is that they are the result of a neuritis of the vagus, causing arrest of the heart. The paralysis of the diaphragm or of the intercostal muscles, which is sometimes produced by diphtheria, may be due to a neuritis of the phrenic or of intercostal nerves, though it is not certain that this is always the explanation. But before we assume that sudden death after diphtheria is caused by a neuritis of the vagus, we must remember that whereas the action of the respiratory muscles is brought about by a stimulus from the respiratory centre in the medulla oblongata, the action of the heart is maintained by the automatic contractions of the cardiac muscle, and that it is by no means certain that a neuritis of the vagus would arrest the working of the heart. We must always bear in mind that the cardiac muscle itself is the prime factor in the circulation, and it seems likely that the fatal syncope which may follow diphtheria is due to a diseased condition of the muscular wall of the heart itself, rather than to disease of the nerves which pass to it. The microscope alone cannot determine this question, for when degeneration of muscular fibre is present it may be impossible to say whether the toxic action was exerted on the nerve-ends, or on the motor end-plates, or on the muscle itself.

That the muscular wall of the heart does undergo serious degeneration in diphtheria has been shown by several observers. Dr. Sidney Martin¹ found that the diphtherial albumoses produced an advanced degree of fatty degeneration in the cardiac muscle, and he was unable to find any degeneration in the vagus nerve. Dr. Mott² found fatty degeneration of the heart muscle in each of four cases of diphtherial paralysis which he examined, but in only one of them was there degeneration of the peripheral nerves. My colleague Dr. Poynton³ has recently published a comparative study of the condition of the heart wall in diphtheria and in rheumatism respectively, and has described very marked degeneration of many of the cardiac muscular fibres, and complete destruction of some parts of them. It is therefore proved that in fatal cases of diphtheria the cardiac muscle is often much degenerated. The conclusion is irresistible, that it must be more or less degenerated in a large proportion of the cases which recover. Can we detect any evidence of this degeneration by clinical investigation? If so, fatal cardiac syncope might be avoided, and many lives might be saved which would otherwise be lost.

The clinical indications which should be sought for are these:

1. Feebleness of the pulse wave.
2. Feebleness and diffusion of the cardiac impulse.
3. Extension of the cardiac dulness to the left.
4. Feebleness of the first sound at the apex, with accentuation of the pulmonary second sound.

These four indications of a weakened left ventricle would all naturally be expected in a heart in which fatty degeneration of its muscle has been produced. They are all present, more or less, in many cases of diphtheria.

5. A fifth sign, which could not have been anticipated, but which is usually present also, is a marked accentuation of the aortic second sound. This is often very decided, yet the radial pulse is not tense, and one can only imagine that the tension in the aorta is raised by a contraction of the splanchnic arterioles through some central vasomotor irritation caused by the toxins. If the vascular tension is much increased at the same time that the ventricle is weakened, the danger of fatal syncope is obviously great.

PHYSICAL EXAMINATION OF THE HEART.

All these physical signs need careful and accurate investigation. It is a matter much to be regretted that the examina-

tion of the heart is often very imperfectly performed, and that a hasty auscultation—easily satisfied if no murmur is detected—is in too many cases the only method employed. But alterations in the normal sounds of the heart may be vastly more important than a murmur; an almost inaudible first sound, or a greatly exaggerated aortic second sound, may be much more alarming. One must go even further, and say that the stethoscope has been far too dominant in the physical examination of the heart (and also of the lungs), and that palpation and percussion are too much neglected and often most inefficiently performed. Especially with regard to percussion, the ordinary method of examination of the heart is quite fallacious and almost useless. What is called the “superficial cardiac dullness” (often the only point investigated by percussion) informs us merely how much of the heart is not covered by lung; it gives us no information as to its actual size. Yet to be able to detect this actual size and to discover any enlargement of the left ventricle or of the right auricle, is often of the most extreme value to the practitioner, for a correct determination of these is in very many cases a matter of vital importance to the patient.

The difficulty felt by many in ascertaining the true size of the heart is largely caused by the word “deep.” The outer margins of the heart are covered by lung, hence it has been thought that the true cardiac dullness can only be detected by a forcible percussion which shall bring out the deeper dullness in contrast with the overlying pulmonary resonance. Unfortunately this forcible percussion brings out far too much pulmonary and gastric resonance from a distance, and usually confuses the result. The trained ear may sometimes be able to detect the change of note at the same point whether the percussion employed be light or forcible, but a light percussion demonstrates the margin of the heart with greatest certainty. The edge of the heart on either side is thick and airless while the overlapping lung is thin, so that light percussion easily reveals the limit of the heart to left and to right, as well as the smaller area of its surface which is not covered by lung.

A finger of the left hand is by far the best pleximeter, and it is one which is always available. The terminal phalanx should be firmly pressed on the spot where percussion is to be practised, and the rest of the finger kept away from the chest wall, or the middle phalanx may be used if the terminal phalanx is slightly hyperextended, so that both it and the first do not press on the chest. If this method be adopted the practitioner will have no difficulty, after a little practice, in defining with very considerable accuracy the limit of the heart both to left and to right. The limits thus discoverable are verified by *post-mortem* examination in fatal cases.

When the extent of the cardiac dullness transversely has been determined by careful light percussion in the fourth right, and the fourth, fifth, and sixth left intercostal spaces, the position of the lateral margins of the heart may easily be defined if it be remembered that they both slope upwards and inwards, and the percussed finger be held in a position parallel to this slope in each case. The right margin above the nipple level rapidly approaches the sternum; but when the right auricle is much dilated, its dullness may be detected in the third space as well as in the fourth. The left margin normally rises to the inner side of the nipple, but in a moderately dilated heart the limit of dullness will be found to pass through the nipple, and where the dilatation is great it may cross the vertical nipple line at one, two, or in extreme cases even three fingerbreadths above the nipple. For further details I may refer to my paper on Acute Dilatation of the Heart in Rheumatic Fever in the *Medico-Chirurgical Transactions* for 1898.

CARDIAC DILATATION IN DIPHtheria.

In a child suffering from diphtheria the cardiac dullness is usually increased towards the left. It is very important to determine carefully the extent of this increase. So long as it does not exceed one fingerbreadth outside the left nipple line, there is, I think, usually no immediate danger. But if the dullness is greater than this, the case should be very carefully watched. If the dullness extends two fingerbreadths to the left of the nipple line, there is urgent peril, and the child must not be allowed to sit up in bed for any reason whatsoever.

I wish to draw special attention to the fact that the increase of dullness is sometimes very rapid: a further extension from one fingerbreadth to two may occur within a few hours. This acute dilatation is frequently accompanied by vomiting, and this symptom is an important danger signal. It has long been recognised that when vomiting occurs in diphtheria the prognosis is bad, but I do not think that it is generally known that the vomiting in this disease is often a sign of an acute cardiac dilatation.

The symptom may be due to other causes, but in every case in which it occurs, especially if along with it there is increased pallor of face and a feeble pulse, the size of the left ventricle should be most carefully investigated afresh, even though percussion had been practised only a few hours previously.

CASE VII.—A little girl under my care at the Hospital for Sick Children, who had suffered from a severe attack of diphtheria six weeks before, and whose cardiac dullness extended one fingerbreadth to the left of the nipple line, was seized with vomiting which was ascribed to some pharyngeal irritation. But her marked pallor attracted attention to the heart, and it was found that the dullness had within twenty-four hours increased to two fingerbreadths outside the nipple line. By great care and watchfulness her life was saved. I have seen other similar cases.

When the shock of the acute dilatation has passed off, or if it have occurred more gradually, the patients may look and feel well; yet their lives are in urgent danger. It is, I believe, cases of this kind which come to an unexpected and tragic end. It is the apparent recovery, while the heart remains enfeebled and dilated, which misleads the practitioner who does not carefully percuss and palpate.

The cardiac dilatation of diphtheria may occur at an early period of the illness, even after only a few days. But a rapid dilatation, or a rapid increase of an earlier dilatation, may take place even after several weeks. In the case already narrated it occurred six weeks after the onset. It is therefore necessary to keep a very careful watch on the condition of a child's heart for at least two months after a severe attack of diphtheria.

I have seen one case in which a permanent dilatation of the heart, apparently the result of diphtheria, caused dyspnoea, dropsy, and such imminent danger to life, that removal of blood by leeches was necessary. But I doubt whether the diphtherial dilatation is often permanent; the less seriously damaged hearts probably again become of normal size, while the worst cases die; for the mortality of “diphtherial paralysis” is certainly much greater than would be inferred from the statements of the textbooks. It must be added that the virulence of diphtheria is more intense in children than in adults. Herein there is a marked contrast to the behaviour of the disease of which I next speak—influenza; for this is often highly dangerous to adults, while it affects children much less frequently and usually less severely.

CARDIAC DILATATION IN INFLUENZA.

In influenza rapid dilatation of the heart frequently occurs, to a greater or less extent, within a day or two after the onset of the disease, and it sometimes causes fatal syncope. If the heart be carefully examined, it will often, though not in all cases, be found distinctly enlarged towards the left, the impulse diffused and weakened, the first sound feeble, and the pulse wave also feeble. Here again, as in diphtheria, if the increase of dullness does not exceed one fingerbreadth to the left of the nipple line, there is probably little danger. But if there is a second fingerbreadth of dullness, there is real danger. And in this disease also the extension of dullness may develop rapidly.

CASE IX.—In one of the first years of the influenza epidemic a young man was admitted into St. Mary's Hospital under my care, suffering from influenza and obviously very ill, with some evidence of pleurisy, but none of pneumonia. The cardiac dullness extended as far as one fingerbreadth outside the nipple line. Fearing possible syncope, I asked the house-physician to direct the night sister to give a hypodermic injection of strychnine immediately if the patient were taken ill during the night. A few hours after I saw him an attack of syncope occurred. The strychnine was administered and the house-physician sent for. On examining the patient he found that the dullness of the left ventricle now extended two fingerbreadths outside the nipple line. This I myself found to be the case the next day. Forty-eight hours after the first attack of syncope a second occurred; the house-physician found that the dullness now extended to three fingerbreadths to the left of the nipple line. Half an hour later he was again summoned, and found the patient dead.

Whether acute dilatation may occur some weeks after the onset, as in diphtheria, I do not know; but it is not unlikely, for many of the sequelæ of influenza do not manifest themselves until weeks or even months after the attack, and it is

quite possible that an extra cardiac failure may be one of these. It is certain that the dilatation caused by influenza may remain as a permanent dilatation, and may give rise to very serious symptoms. Thrombosis is also of frequent occurrence after influenza, probably due in part at least to the enfeeblement of the heart. This thrombosis may in its turn produce pulmonary embolisms resulting in great dyspnoea and hæmoptysis, or even in sudden death from obstruction of the pulmonary artery.

Minor degrees of cardiac dilatation after influenza may cause merely a feeling of incapacity for exertion. Unless the size and strength of the left ventricle be carefully ascertained, such a patient may be considered a hypochondriac, and very injudicious advice may be given to him. Active exertion, unless carefully controlled, may do such a patient much harm by increasing the dilatation. Any prolonged strain—for instance, the effort of cycling uphill—may be very injurious after an attack of influenza.

CASE X.—Some years ago I was asked to see a young lady whose only complaint was that she felt languid and incapable of exertion. She had suffered from influenza some months before. She was very fond of cycling, and had acted as the "hare" in games of "hare and hounds" by cyclists. Her left ventricle extended to two fingerbreadths outside the nipple line. There was no murmur. By careful regulation of her exercise she recovered completely, and the limit of the cardiac dullness is now well inside the nipple line.

CARDIAC DILATATION IN RHEUMATIC FEVER.

In rheumatic fever, even in the most subacute attacks, acute dilatation of the heart seems to be invariably present. Since I first observed its occurrence, in 1894, I have never seen a first attack of this disease, whether in a child or in an adult, in which it was absent. When the rheumatic attack is over, the dilatation lessens and the cardiac dullness may again become of normal extent. It may be well to quote a recent case in illustration of this:

CASE XI.—A woman of 28, who had never previously suffered from rheumatism, was admitted into St. Mary's Hospital under my care on May 29th, 1900, on account of some pain in her limbs, of five days' duration, with slight swelling of both knees and of one ankle, and some oedema of the legs. The temperature was 101° F. There was no murmur, but the cardiac dullness extended $\frac{2}{3}$ fingerbreadths outside the left nipple line. The house-physician prescribed potassium citrate with caffeine and strychnine, but no salicylates. After twenty-four hours the rheumatic symptoms were worse, and the dullness had increased to 3 fingerbreadths to the left of the nipple line. The medicine was then changed to salicylate and sodium bicarbonate. Within thirty-six hours the temperature had fallen to normal, and it remained normal. On June 5th—five days after the change of medicine—the left limit of the cardiac dullness was only $\frac{1}{2}$ fingerbreadth outside the left nipple line. On June 12th it extended only $\frac{1}{2}$ fingerbreadth to the left of the nipple, and on June 15th the border of the dullness was in the nipple line itself. On her discharge from the hospital the cardiac dullness and sounds were normal.

In this instance the rheumatic dilatation was unusually great for a first attack, and the cardiac weakness had produced some oedema of the legs, which is also unusual in a first and apparently slight attack of rheumatism. The latter symptom disappeared after three days' rest in bed, and coincidentally with a marked improvement in the rheumatic symptoms and with the fall of temperature. Both the oedema and the considerable increase of cardiac dullness indicated a severe implication of the cardiac muscle, yet there was no evidence of endocarditis or pericarditis.

In later attacks also an acute dilatation is usually present, for it may almost always be observed, on careful examination, that the dullness of the left ventricle diminishes to some extent—greater or less—as the attack subsides. At all events one may say with certainty that an acute dilatation of the heart is much more common in rheumatism, even in slight attacks, than in either diphtheria or influenza. Yet, though more common, it is far less dangerous. I have already said that an extension of the cardiac dullness to two fingerbreadths outside the left nipple line is an indication of grave danger in a child affected with diphtheria. I will now add that the same amount of increased dullness in a child suffering from rheumatism implies, in itself, no immediate danger of death whatever.

It is a very remarkable fact that the dilatation of rheumatism is so much less dangerous than that of diphtheria or of influenza, in spite of its greater frequency in considerable amount. The difference must be produced by a different effect of the several toxins upon the cardiac muscle. In diphtheria, and apparently in influenza, the muscular fibres of the left ventricle suffer greater destruction; in rheumatism the myocardial changes are less intense, and one can only suppose

that the elasticity of the ventricle is more affected. Dr. Poynton's sections show that, though in the rheumatic heart there is evidence of fatty degeneration of the cardiac muscular fibres, with interstitial foci of small cells and vascular dilatation, yet the destruction of muscle is much less pronounced than in the diphtherial heart. I pass round drawings illustrative of this.

But though, as I have said, an increase of cardiac dullness to two fingerbreadths outside the nipple line in a case of rheumatism involves no danger of sudden death, yet a further extension, occurring rapidly, may cause decided symptoms of collapse. Here again, as in diphtheria, a sudden vomiting is often the danger signal. Pallor, coldness, and general feebleness follow, and the cardiac dullness is found to have increased by an additional fingerbreadth within a few days or even within a few hours. From two fingerbreadths outside the nipple it increases to three, or from three it may extend to four; at the same time the border of the left ventricle reaches a greater height above the nipple, from one to two fingerbreadths, or from two to three. The cardiac impulse becomes more diffused and much weaker. Mitral valvulitis will almost certainly be present, causing an apex systolic murmur, and very frequently a presystolic murmur is present also. There may be in addition evidence of pericarditis or of pericardial adhesion. The implication of the mitral valve, allowing regurgitation into and therefore increased tension within the auricle, has doubtless some share in the great enlargement of the left ventricle which may occur in the later stages of rheumatism in a child. But the damaged condition of the cardiac muscle is certainly the chief cause, and such cases should not be looked upon as simply "mitral regurgitation." In children the dilatation is of much greater importance than the valvular lesions. But it may easily escape the notice of a practitioner who does not constantly practise a careful examination of the heart by palpation and percussion, for it is very remarkable how slight may be the superficial evidence of a rheumatic toxæmia, which yet is able to produce a perceptible enlargement of the heart.

CASE XII.—A little girl under my care, who was convalescent from a rheumatic attack which had dilated her left ventricle to two fingerbreadths outside the nipple line, was found one day to have some bronchial catarrh, with slight pain in her left ankle, and a new small nodule over this joint. On my next visit to the ward I was struck with her pallor and evident illness; the ankle was better, but the left ventricle was now dilated to three fingerbreadths instead of two.

CASE XIII.—In another child the only obvious indication of a rheumatism which perceptibly enlarged his left ventricle was a very slight and transitory patch of erythema on his neck.

The slightest suspicion of rheumatism in a child should therefore lead to careful and repeated examination of the heart. Even in adults, much oftener than is generally recognised, it is fresh rheumatism that kills, breaking down compensation. It is important to notice that at the necropsy of patients who have died from chronic rheumatic heart disease, there is usually evidence of fresh endocarditis on the cardiac valves. And clinically it may often be observed that when a case of mitral stenosis breaks down there is some evidence of fresh rheumatism.

CASE XIV.—A woman recently under my care at St. Mary's Hospital exemplified this. She was moribund on admission, and was found to have mitral stenosis, a much dilated heart, and thrombosis of the right jugular vein. It was thought by all who saw her that she would not live more than a day or two. Venesection three times repeated, once from each arm and once from the left jugular, saved her from the immediate danger; the thrombosis did not extend, and she improved so much that she begged to be allowed to go home to look after her children. But during the five weeks that she lived after her admission there was persistent recurrence of a slight rheumatic arthritis in the joints of her hands—an arthritis which at once yielded to salicylates, but regularly returned when the medicine was omitted. At the necropsy evidence of fresh endocarditis was found, in addition to the mitral stenosis.

I have now shown that diphtheria, influenza, and rheumatic fever are all, in various measures, associated with acute dilatation of the heart. In fatal cases of diphtheria and of rheumatism, myocarditis and fatty degeneration, or even destruction of cardiac muscle are often found; probably similar lesions exist in influenza.

Now, it cannot be questioned that in diphtheria and in influenza the injury to the heart is caused by the deleterious action of the toxin produced by a microbe. In each of these two diseases the microbe is well known. Is it not then in the highest degree probable that rheumatic fever also is a microbial disease, and that the resulting cardiac inflammation is due to its toxin? A study of rheumatism

in childhood compels me to say that it is as certainly microbic in nature as measles or whooping-cough, in neither of which the microbe has yet been demonstrated.

But it is no longer a matter of inference merely that rheumatic fever is a microbic disease, for it has been proved by Dr. Poynton and Dr. Paine of St. Mary's Hospital¹ that a diplococcus can be isolated from the tissues of a patient who has died from rheumatic fever, and also from blood obtained by venesection during life, which can be cultivated in pure culture in a special medium, and which, when injected intravenously into rabbits, can produce in them a transitory arthritis of several joints, pericarditis, valvulitis, cardiac degeneration and dilatation, pleurisy and pneumonia, and yet nowhere any suppuration. By careful staining of their sections they have succeeded in demonstrating these organisms in the substance of the mitral valve, in the tonsils, and in many other organs. They have found them also in that most typically rheumatic lesion, the subcutaneous nodule. Their results will, of course, require confirmation by other observers, but it certainly looks as if the problem of acute rheumatism has at last been solved.

IMPORTANCE OF THE DIAGNOSIS OF CARDIAC DILATATION.

Finally, let me appeal for greater care and accuracy in the examination of the heart by percussion and palpation. It is a matter of great importance to the patient. Dilatation and feebleness of the left ventricle may indicate, as I have shown, urgent danger of death from syncope, and neglect of this indication may cost the patient's life. Dilatation of the right auricle (quite easily detected by percussion in the fourth right interspace), and weakness of the right ventricle (detected by palpation of the epigastric region) are usually accompanied by considerable dyspnoea and often by some lividity. A marked degree of dilatation of the right auricle (from two to three fingerbreadths to the right of the sternal margin in the fourth space) may indicate grave danger of death from asphyxia and call urgently for venesection or leeches.

After a considerable experience as an Examiner in Medicine I am forced to the conclusion that these facts are still very inadequately recognised, for I rarely meet with a candidate who understands the object of percussion of the heart or the proper method of procedure. He usually thinks only of the useless "superficial cardiac dullness," and contents himself with trying to determine a horizontal upper limit (which does not exist), and then palpating the cardiac impulse, as if that were equivalent to the left border of the dullness! The right limit of the heart he usually neglects altogether. And even some physicians of great eminence are apparently not conversant with the fact that the dullness of the right auricle normally extends one fingerbreadth into the fourth right space, and that its border can be quite easily detected by careful light percussion.

Let me assure you, gentlemen, that there is really very little difficulty in the determination of the actual size of the heart, and that a small amount of careful practice will demonstrate to you what an enormous advantage both to your patients and to yourselves will result if you will accustom yourselves to consider this as an essential part of the investigation of every case to which you are called.

REFERENCES.

¹ Goulstonian Lectures, 1892. ² Croonian Lectures, 1900. ³ *Lancet*, May 12th, 1900. ⁴ *Ibid.*, September 22nd and 29th, 1900.

UNIVERSITIES FOR WOMEN IN JAPAN.—The rich family of Mitsui of Tokyo has offered an extensive site in that city for the erection of a university for women, and three other citizens have between them contributed a sum of £24,000 for the cost of the necessary buildings. The work is already in progress, and it is hoped that the new university will be opened in the spring of 1901. It is not likely that there will be any want of students, as in recent years very many young ladies of good family have applied to be admitted to the university courses, especially to the Faculty of Medicine and the Polytechnic School. The latter institution is intended for the training of civil engineers, a circumstance which seems to show that Japan is about to set an example to Europe in opening up a new sphere of labour for the women of the future.

EPIDEMIC ARSENICAL POISONING AMONGST BEER DRINKERS.

BY

NATHAN RAW, M.D., F.R.S. EDIN.,

Medical Superintendent, Mill Road Infirmary, Liverpool;

FRANK H. BARENDT, M.D., F.R.C.S.,

Physician to St. George's Hospital for Diseases of the Skin, Liverpool;

AND

W. B. WARRINGTON, M.D., M.R.C.P.,

Pathologist, Stanley Hospital; Demonstrator of Pathology, University College, Liverpool.

INTRODUCTION BY NATHAN RAW, M.D.

THERE is every appearance that the severe epidemic of arsenical poisoning which was first accurately diagnosed by Dr. Reynolds, is subsiding. It is difficult to accurately determine when the outbreak actually occurred, but the following table, showing the admissions of cases of alcoholic neuritis, is of great interest as giving some idea when the contamination of the beer really began. The symptoms of alcoholic neuritis and arsenical neuritis are, as far as I can determine, very much the same when the lesion is advanced; but the association of pigmentation of the skin renders the diagnosis clear.

Table of Admissions of Cases of Alcoholic Neuritis in 1898 and 1900.

	1898.	1900.
January	3	0
February	1	5
March	1	1
April	0	2
May	3	7
June	2	11
July	4	12
August	1	17
September	3	25
October	2	22
November	3	26
December	3	15 up to Dec. 10th.
Total	26	143

From this table it would appear that the contamination of the beer commenced about June, and remained fairly constant until November, when a further increased contamination took place causing more acute symptoms.

Arsenical melanosis, using the term in its etymological sense to indicate a deepening or blackening of the skin, was accurately described by Nielsen, who records some previous outbreaks of arsenical poisoning. Moreira reports some cases where the skin was diffusely pigmented after taking rat poison, and Barthélemy reports 405 cases of arsenical poisoning at Hyères, which resulted from the mistake of a wine merchant who poured a solution of arsenic into his wine casks. Of these cases the greatest number showed general pigmentation of skin varying from brown to black. The same writer thinks that the great outbreak which occurred on the banks of the Seine in 1828, and again in 1829, causing the loss of 40,000 lives, was due to arsenical poisoning of wine. I have had over 70 cases of neuritis during the present epidemic under treatment with so far only one death. The majority of the cases are slowly improving, some have been discharged recovered, whilst a small minority have advanced motor lesions, with but small hope of permanent recovery. I have only been able to detect arsenic in the urine in 5 cases out of 33 examined, and these were all recent acute arsenical attacks. The symptoms have been so well described before that it is unnecessary to recapitulate them, but they have been undoubtedly caused by the presence of arsenic in the sulphuric acid in the manufacture of invert sugar or glucose used largely in brewing cheap beer and stout. This sugar has been promptly withdrawn, with the result that the cases have very rapidly decreased. The quantity of arsenious acid has in some cases been found to be as much as 3 grs. per pound of sugar, so that it is possible for a glass of beer to contain one-fifth of a grain of arsenic, and certainly some severe lesions have been pro-