aggravation of the obstruction to the bladder and rectum, and the former symptoms did not particularly exceed what is often experienced by healthy women." The os was not felt till an anæsthetic was given and the hand passed into the vagina; it was then dilated with one finger and the child's scrotum recognised. The child was next rotated somewhat with the help of the other hand externally, so that breech and anus were brought within reach. "It was then obvious," writes Oldham, "we had to do with a complete retroflexion at term." A foot was seized, pulled through the os, and the child delivered dead *per vaginam*.

Of the 5 cases referred to by Veit¹¹ as persisting to term, the first is Franke's case,¹² which terminated naturally, the cervix being obliterated, and the os behind the symphysis. The second is one recorded by Merriman,¹³ in which labour was difficult, terminating naturally, the child dead. The third case, also recorded by Merriman, was a primipara who had had only some pain in emptying the bladder during the last two or three months of pregnancy. Escape of the waters followed by pain first attracted Merriman's attention, who then detected retroversion. On the following day the pelvic tumour pressed on the perineum, when a severe attack of convulsions set in, and were said to be arrested by bleeding to 12 ounces and other treatment. The child's movements then ceased. Fifteen days later the os uteri could not be reached, but on the sixteenth day a change occurred, the retroverted segment of the uterus slipping upwards and a fleshy mass coming down into the vagina, when the head could be detected, was perforated, emptied of its contents, and the child extracted. The fourth case is that of Oldham referred to above, and the fifth that of Wyse,¹⁴ in which the feet were seized through the os, the head being pushed up with the fingers in the bowel.

I believe another case is recorded by Stille, the reference to which I am not able to give, where also delivery was effected *per vias naturales*, the os uteri being brought sufficiently into line with the ostium vaginæ, the cervix being obliterated.

The case recorded in this paper differs from these 7 cases in five points: (1) The non-obliteration of the cervix; (2) the impossibility of bringing the cervix into such position as would enable delivery to take place through the vagina, and this even after control of the uterine body had been obtained through an abdominal incision; (3) delivery by Cæsarean section; (4) the presence of an intramural fibroid; and (5) removal of the anterior uterine wall with the fibroid.

NOTE.—A letter of November 17th, 1900, says: "Mrs. — is very well indeed. There has been no return of the menstrual flow nor any periodic disturbance of health suggesting a tendency for its return to recur. The external wound gives her no trouble, though there is still a pin-point superficial opening."

REFERENCES.

- ¹ *Science of Midwifery*, vol. i, p. 263. ² *Textbook of Pregnancy*, translated by Edgar, p. 234. ³ *Manual of Obstetric Accidents, Emergencies, and Operations*, p. 183. ⁴ *Manual of Obstetrics*, vol. i, p. 391. ⁵ *A Manual of Midwifery*, p. 342. ⁶ *Textbook of Midwifery*, Syd. Soc., vol. i, p. 392. ⁷ *Précis d'Obstétrique*, p. 722. ⁸ *Textbook of Obstetrics*, p. 212. ⁹ *Obstetric Medicine and Surgery*, vol. i, p. 346-7. ¹⁰ *Obstetrical Transactions*, vol. i, p. 318. ¹¹ *Volkmann Sammlung, Klin. Vort.*, 1879, No. 170. ¹² *Monatsschrift f. Geburtsh.*, Bd. xxi, s. 161. ¹³ *Dissertation on Retroversion of the Womb*, London, 1810, p. 25. ¹⁴ *Ed. Journal of Medical Science*, Part 4.

A SECOND SUCCESSFUL CASE OF CÆSAREAN HYSTERECTOMY.*

By COLIN CAMPBELL, L.R.C.P.L., M.R.C.S. ENG.,
Saddleworth.

History of Patient.

ON July 25th, 1900, I was called about 1 P.M. to see a woman in labour in a gipsy van. The labour had commenced about "daybreak."

The woman, aged 24, was about 4 feet in height. Her mother said she had been "deformed all her life." It was her second pregnancy, the previous one having been terminated by craniotomy at Halifax "about" a year before.¹

Examination revealed a pseudo-malacosteon rachitic pelvis, also that the child was living. I therefore advised its imme-

diate removal by abdominal section, and that the patient should be at once removed to a room, as in the van the bed was only approachable on one side, and one could not stand nearly erect. (2) That if they declined removal and operation that she should be immediately taken to an infirmary, Oldham and Ashton being respectively five and six miles distant.

After much delay the former alternative was chosen, and at 5 P.M. I was informed that a Good Samaritan had taken her in, making a bedroom of her "parlour," which opened directly on the road. I saw the patient again at 6 P.M. with my partner, Dr. David Price, equipped for any emergency.

Selection of Operation.

In such a case, supposing the child to be living, the alternatives before the practitioner are:

(1) Killing the child by the brutal "operation" of craniotomy.

(2) Ordinary Cæsarean section.

(3) Porro's operation.

(4) Symphysiotomy.

(1) As the child was still living we rejected the first alternative. Personally I can see little difference between plunging a perforator into the brain of a living fetus *in utero*, and plunging it into the brain of a child hidden under the bed-clothes.

(2) Except in the able hands of Murdoch Cameron, or Leopold, Cæsarean section has still a terribly high mortality, and successful cases leave the patient liable to further pregnancies—and "sections"! This is not ideal.

(4) As to the merits or demerits of the Singultian operation we did not enter, rejecting it at once in favour of

(3) A modified Porro's operation, in which both child and uterus are removed, and further pregnancies (and operations) effectually prevented. In Porro's operation the procedure is, shortly, as follows:

To cut down first through the linea alba, next through the anterior wall of the uterus, "avoiding if possible the placenta," next the fetus is removed. "Hæmorrhage from the uterine wound may be checked by grasping the cut edges with pressure forceps, having large transverse blades. The uterus is drawn through the abdominal wall, the intestines being kept back by an assistant, and covered with large flat sponges. And then a Keberlé's *serre-naud* is passed round the lower part of the uterus." This is tightened, and the stump fixed in the abdominal incision by means of transverse pins.

The modification of this operation adopted by the late Mr. Lawson Tait appears to me to be of more practical importance than the original design. He so simplified the procedure as to render it practicable, at a moment's notice, by any surgeon who has at hand a knife, a piece of stout rubber tubing, and two knitting needles; by his method it is also a bloodless operation.

In Porro's operation an assistant (as well as an anæsthetist) is required to check the hæmorrhage and work the sponges; large flat sponges should be at hand, also a supply of transverse-bladed forceps, and that somewhat cumbersome instrument, the *serre-naud*. The control of the hæmorrhage from both sides of the uterine incision, as well as from a possibly wounded placenta, suggests considerable difficulty.

Mr. Tait ligatured the uterus before opening it. This was done by passing a loop of strong rubber tubing over the fundus, then drawing the tubing down to the lower part, and tightening it as an Esmarch's bandage. Thus performed, neither assistants nor unusual instruments are required.

In anticipating a similar, and also successful, operation, my friend, Dr. John Sutcliffe (now of Cheadle Royal), suggested that the loop of tubing which was intended to pass over the fundus would be more manageable if stiffened, and this was found to be so; for whilst it was impossible to get the limp tubing over the fundus through an incision which extended upwards only as far as the umbilicus, it was easily done when a piece of gutta-percha-covered wire was first passed through.

The operation was performed under the most unfavourable conditions; the room was small, dark, and crowded with furniture; no nurse or handy woman was available, only a neighbour who had once "seen an operation." The patient had become irritable and restless, her pains were continuous, pulse 128, temperature 100°. My partner's whole time was occupied in looking after the anæsthesia. The abdominal

* Read at a meeting of the Lancashire and Cheshire Branch of the British Medical Association.

incision was 4 inches in length downwards from the umbilicus; through this the tube, stiffened as described, was passed over the fundus, and then drawn down as low as possible, the wire drawn out, and the tube tightened and knotted. Then the uterus was opened without hæmorrhage, and the left hand introduced to push aside the placenta and find a foot. This was found with some difficulty, and a living infant delivered. Next the uterus with placenta enclosed were drawn out, coils of gauze being introduced to prevent intestines following, and the only assistance that was required by the operator was the "neighbours" to hold up the uterus whilst it was being transfixed with guarded pins and amputated. Four sutures closed the abdominal wound, the "stump" filling quite one-third of its length.

After-History.

On the third day the temperature reached 102°; then, after a dose of Epsom salts, it became normal. The stump came away on August 21st. I exhibit it to show the effect of the elastic ligature. I regret I am unable to show the patient and infant.

The woman was walking about by the fifth week, and they then left the district. I have had the help of the police and the "King of the Gipsies," but the scent was lost at Blackburn.

In conclusion, I would express the hope that two successes so easily obtained may (in the words of Dr. Herman) help "to relegate craniotomy, the induction of premature labour, and symphysiotomy, alike to the past."

NOTES AND REFERENCES.

¹ They were very uncertain about everything. Two "doctors" were said to have attended, but neither of the names given as at Halifax are in the *Medical Directory*. ² Galabin. ³ *Lancet*, 1890. ⁴ In the previous case reported in the *Lancet*, I described the operation as "Tait-Porro's operation." As I find this is not generally understood, I now adopt Galabin's name, "Cæsarean hysterectomy."

A NOTE ON THE TREATMENT OF PUERPERAL ECLAMPSIA.

By R. P. RANKEN LYLE, M.D., B.A.O.,

Lecturer in Midwifery and Diseases of Women, University of Durham
College of Medicine, Newcastle.

IN appreciation of Dr. FitzGerald's article (in the *BRITISH MEDICAL JOURNAL*, November 24th, p. 1496) on the treatment of this condition; but more as a criticism on the article by Dr. Jardine, I would like to point out that Dr. Jardine makes a great mistake in considering "diuretic infusions" the treatment of puerperal eclampsia. It can only be considered a small, and by no means the essential part of the treatment.

There are three great principles in the treatment of this condition:

1. The purifying of the blood.
2. To control the convulsions.
3. The emptying of the uterus.

Diuretic infusion is a valuable addition to the usual treatment adopted for purifying the blood (purgatives, enemata, diaphoretics, vapour bath, diuretics, etc.), but it has no immediate effect in controlling the convulsions, nor has it any effect on the action of the uterine muscle.

In order to control the convulsions in eclampsia, it is necessary to allay the irritability of the cerebro-spinal system; morphine, chloral, veratrum viride, and chloroform have all been used. Chloroform is undoubtedly bad, as its action is very temporary, and very depressing to the patient; the action of chloral and veratrum viride is more lasting, but they are both cardiac depressants; while morphine, judiciously given, is quite free from any disadvantage, and has the following advantages:

1. It controls the convulsions by allaying the irritability of the cerebro-spinal system.
2. It prevents excess of waste products being thrown into the blood.
3. It does not weaken the patient.
4. It does not injure the child.
5. It has no effect on the kidney.
6. When the patient is under its influence labour often commences, and quickly terminates without causing more convulsions.

In my own experience at the Rotunda Hospital, of eight consecutive cases of puerperal eclampsia, all of them treated

by morphine, only one died. In this case only did the morphine appear to have no effect, the convulsions continuing frequent and severe, and the temperature rapidly rising to 106.6° F. The liver was very large and extensively diseased.

Now, with reference to the third great principle in the treatment—the emptying of the uterus. As soon as the patient enters the second stage of labour she should be delivered by the forceps, but any mechanical interference (induction of premature labour, mechanical dilatation of the cervix, version, or *accouchement forcé*) during the first stage of labour is extremely unsuccessful treatment, as is evidenced not only by his own statistics (all the maternal deaths having occurred under this heading), but by the experience of many authorities.

Should labour not supervene under the morphine, the morphine will generally control the convulsions for the time being, and the patient, if carefully treated, will probably go on to full term without any recurrence of the symptoms of eclampsia.

A NOTE ON THE OCCURRENCE OF ABORTION.*

By J. B. HELLIER, M.D.

Honorary Obstetric Physician, Leeds Infirmary.

IT is a question of some interest to determine what proportion of married women have one or more miscarriages, but never carry the function of utero-gestation to full term. One is not infrequently asked under such circumstances by a woman disappointed of maternity what her chances are of bearing a living child. Some data for answering this question are afforded by the following statistics. I have had the opportunity of collecting records concerning 1,800 women who attended as out-patients at a gynaecological clinic. The observations were made while I was on the honorary staff of the Hospital for Women and Children at Leeds; and by the courtesy of my late colleagues I was enabled to have reference made to their records as well as to my own. The out-patient papers contained a statement of the number of children and abortions in each case. It is quite true that a woman cannot always be relied upon to state accurately the number of abortions she has had. She has probably miscarried as many times as she says, perhaps more. This uncertainty qualifies the value of the figures here given, but does not seriously invalidate the one point on which I draw conclusions from them. The patients being gynaecological, and having, *ipso facto*, in most cases some pelvic disorder, the tendency amongst them to miscarry would be above the average for all married women. They were drawn principally from the poorer working classes in a manufacturing centre, and were almost all engaged in household or other manual labour. The cases of unmarried women were neglected in collecting these statistics.

The 1,800 married women had given birth to 6,974 children, an average of 3.87 each. They had had, according to their own statement, 1,288 abortions. This amounts to about one abortion to every five and a-half children. The number of women in the 1,800 who had had abortion but no children was 58. Deducting from the 1,800 184 women who had never been pregnant, 1,616 remain. So that of 1,616 women who had been pregnant, 58 had had one or more abortions, but had never gone to term. Thus it follows that amongst a group of women where the frequency of abortion might be supposed to approach the maximum, 96.5 per cent. of those who became pregnant did sooner or later bear one or more children at term.

I think this statement of probabilities is likely to afford consolation to many of our patients.

A SUBJECT-PURVEYING SOCIETY.—There appears to be a dearth of dissecting material in Brussels. The scarcity is said to be increasing, and to be so great as to threaten a dearth not only in dissecting rooms but in operative surgery. This state of things has led Drs. Aburnin and G. Daniel to suggest the foundation of a new society, to consist of persons who will undertake to leave their bodies at the disposal of the universities to be used in furthering medical education and the advance of medical science.

* Read before the North of England Obstetrical and Gynaecological Society.