its development and recurrence; so that when you have really cured a patient of a malignant tumour you leave him still more predisposed to cancer than the average man; much the same as in tuberculosis when you have cured a tuberculous joint your patient is unfortunately still predisposed to some other manifestation of the disease. I quite believe in the curative power of the knife for cancer locally. Every time a tumour is entirely removed it is cured. Recurrences of the original disease are only the natural development of traces left behind, and unless the disease is taken in hand at a very early stage traces usually are left behind. But, unfortunately, when you have removed all, and cured your patient of one kind of cancer, he is still more liable than others to be attacked again.

I don't wish to dishearten you, least of all to prevent you from urging on your patients the only remedial measure we have to offer them, for it seems to me no reasonable person can doubt that benefits result from the surgical treatment of cancer; but we must not shut our eyes to facts if we wish to advance in knowledge, and this constitutional predisposition—of very varying intensity—is one of the facts of malignant disease. I could illustrate this opinion by quoting several cases, among which the following is a striking example:

In 1894 I saw a female patient, from whom I removed a small epithelioma of the lip. There was no local recurrence, but three months later a submental gland enlarged, and was excised. It was infiltrated with epithelioma. Again, three months later, we had a similar experience with a submaxillary gland. Then the patient was cured. These two glands, which cropped up respectively at three and six months after the first operation, were, of course, parts of the original disease, and might have all been excised at once had there been any means of telling how far it was disseminated at the time of the first operation. In 1898 I excised from this same patient a malignant tumour of the breast of the ordinary mammary type, and recently she died from further extension of this disease. Other similar cases occurring to my mind at this moment are of carcinoma of the left breast, followed three years later by a sarcoma of the right breast, one of epithelioma of the lip, followed by pigmented sarcoma of the sole of the foot two years after cure, and two more cases of epithelioma of the lip, in one of which an epithelioma subsequently developed in the tongue, and in the other in the scar of an old ulcer of the leg. Hence, with experiences of this kind it would be quite consistent to allow that our patient might well have suffered in the first instance from rodent ulcer, and that an epithelioma might subsequently have developed in the scar.

DIAGNOSIS OF THE CASE.

I pass over the possibility of this disease having been epithelioma all through, because, as I have already stated, I do not think such a view to be a possibility. This would be the simplest and most plausible explanation if it could be accepted, but it cannot. The duration of the case is too long for epithelioma under almost all circumstances; but when this disease has been continuously present as in our patient for many years without treatment or interruption of any sort, and is still slowly progressing without secondary infection, the view that it is epithelioma becomes untenable. Indeed, I may tell you now, what I was not aware of when we discussed this case in the wards, that the microscope negatives any diagnosis of epithelioma, for the structure proves to be that of a glandular type of growth.

What, then, are we to call this tumour? It certainly has not the least naked-eye resemblance to rodent ulcer; though in its position, in its long history, in its freedom from secondary infection, and in its histological appearance eight years ago and at the present time we recognise the characters of rodent ulcer. Yet the fact that it lacks erosion is so striking a deviation from the type, that we must hesitate to accept it as the same disease.

CANCER OF THE SWEAT GLANDS.

As I have said, rodent ulcer is a carcinoma of one of the appendages of the skin, and so differs from epithelioma, which is a carcinoma of the rete mucosum. We must remember that the appendages differ as much from each other as any

one from the rete, and it seems most probable that cancer of the sweat glands would differ as much from cancer of the sebaceous glands as would the latter from cancer of the rete. Hence my view of this case is that it is one of cancer of the sweat glands, which, while it possesses some of the clinical features of cases of cancer of the sebaceous glands, or ordinary rodent ulcer, tends to grow into a solid tumour instead of excavating the face, as would happen when the characteristic erosion of the latter occurred. I have several times seen small tumours of the face more or less ulcerated on the surface, and having a very glandular type of structure, which I regarded as rodent ulcers, and assumed that the usual erosion would have occurred later if they had not been excised. In the light of this case I am inclined to change that view, and to believe that we have two varieties of carcinoma of the skin appendages, both of which are characterised by continuous slow growth; but in only one of which, true rodent ulcer, or carcinoma of the sebaceous glands, does ulceration keep pace with growth. The two varieties of adenoma of the skin glands are fairly well recognised, and are quite distinct from each other, which is a strong argument in favour of their being also two varieties of carcinoma.

TREATMENT.

Well, gentlemen, we have been dealing with the diagnosis of a rare and peculiar tumour of the face; but my object has been less to attract your attention to this particular case than to try to interest you in the pathology and diagnosis of two rather common but very important affections—rodent ulcer and epithelioma. In a clinical lecture one generally likes to occupy a part of the time in discussing treatment; but, after all, the diagnosis and the recognition of the true nature of the complaint are of the first importance, and often, as in this instance, when you have come to a correct diagnosis the treatment may be described in very few words. In the case of rodent ulcer it is only thorough, very thorough, microscopically thorough, and prompt excision. No waiting for a diagnosis; no tinkering with caustics and cauteries; uncompromising, complete, and immediate removal. To have a reasonable chance of success the operation must always be undertaken early. Everything, then, depends upon your diagnosis, and as the diagnosis of malignant tumours becomes more prompt and correct and their individual pathology better understood, so will their surgical treatment become more successful. In the case of our patient the opportunity for cure has of course long gone by. For him "tinkering" is all that is left to be done. Already the removal of some tender outgrowths has made him comparatively comfortable again, and he is going home, no doubt to return shortly with some other trouble as a fresh nerve or other important structure becomes involved in the growth, and adds a further distress which we must do our best to relieve.

ON RINGWORM INFECTION IN MAN AND ANIMALS.

(With Special Plate.)

By J. L. BUNCH, D.Sc., M.D.,

Research Scholar of the British Medical Association.
(From the Pathological Laboratory, University College.)

SINCE the researches of Gruby on the subject of ringworm, communicated first to the Académie des Sciences in the years 1841 to 1845, the subject has been investigated by numerous observers both in England and abroad, and our knowledge of the subject, both clinically and pathologically, has advanced very considerably. This advance is to a great extent due to the brilliant researches of Sabouraud, who since the year 1892 has contributed several valuable papers on the subject. It is now generally held that ringworm in human beings is usually due to two classes of fungus, the "microsporon" and the "trichophyton," which may be distinguished to a certain extent clinically and microscopically, but still better culturally. A less common form of ringworm is due to a fungus which culturally gives a faviform growth, but the lesions of which do not resemble favus, but are either circinate or a well-marked kerion. The trichophyton class includes two forms: the ectothrix, both large and small-

spored, and the less common (in England) endothrix, while microscopically an intermediate variety can be distinguished, in which the fungus lies both within and without the hair. The question of etiology of these different forms has always been one of extreme difficulty, for the patients are in most cases children, who, either at school or at play, may very easily have associated with other children suffering from ringworm; and, seeing how very numerous the cases of ringworm in London are, the exact source of contagion becomes almost impossible to trace. In the case of adults affected with trichophytosis, the source of infection may sometimes be determinable with considerable exactitude, and of course in children, also, this is sometimes possible.

During the past three years in the skin department of the Middlesex Hospital by the kindness of Dr. Pringle, and more recently at the German Hospital, I have been enabled to investigate the question of the etiology of a large number of ringworm cases, and it is with the question of ringworm infection from animals that I now wish to deal. Even to veterinary surgeons the determination of the diagnosis of ringworm in animals seems to present difficulties, and I have more than once found that cases thus certified were in reality cases of scabies or pediculosis. After careful search, however, I have been fortunate enough to meet with a few cases of ringworm in human beings, and also in animals with which they had respectively been thrown into intimate contact, to isolate and grow the fungus on culture media both from the human being and from the animal, and to demonstrate the similarity between the cultures from the two sources. Cases of ringworm have from time to time been described as having an animal origin, but the source of contagion has only rarely been ascertained, and, indeed, in some of these cases the probability of the patients having contracted the disease from animals appears problematical.

Inasmuch as ringworm cultures obtained from man can be inoculated into animals, and will in them give rise to characteristic lesions in which the fungus can again be demonstrated and reinoculated, it might, I suppose, be argued that in these cases there is a possibility of the horse or dog having contracted the disease from the human being instead of *vice versa*, but this possibility can be practically eliminated by comparing the dates at which the first symptoms were noticed in the two cases.

The first case was that of a girl, aged 3½, in whom had been noticed a week previously a small ringed patch on the right forearm, just above the wrist, and a few days later a second patch on the right shoulder. When seen, the first-mentioned lesion was circinate, measuring 1 inch in diameter, with a slightly scaly surface and a red raised edge, which in places showed some minute vesicles containing clear serum, and the redness extended somewhat beyond the vesicular border, fading off gradually into the surrounding skin. The lesion on the shoulder was also circinate, but smaller; the surface was dry and somewhat scaly, and there was no vesiculation. Microscopic examination showed chains of large squarish spores both inside and outside the hairs, and in preparations treated with 6.8 per cent. liquor potassæ a coarse segmented mycelium was seen in the intrafollicular portion of the hair shaft, terminating in a fringe above the bulb, and also outside the hair in the root sheath and in the connective tissue of the follicle. Scrapings from the lesions showed also chains of spores crossing between the epithelial cells, and a more abundant fine branching mycelium, segmented at somewhat longer intervals, and showing in potash preparations as bright glistening threads, which were, however, less highly refractile than the square-shaped spores. In places, this mycelium gave off short lateral offshoots, which swelled out at the extremity into small oval bodies; and from these, again, there occasionally sprang a still finer and apparently quite recently developed mycelial thread. Similar mycelial appearances I have also observed in ectothrix preparations from other cases, and also in microsporion cases.

Cultures from minute portions of the affected hairs and from the scales, grown on French proof agar, the materials for which were obtained from Paris, so that the cultures might be accurately compared with those of Sabouraud, showed a network of coarse aerial hyphæ, radiating from the centre, white in colour, and terminating at the edge in tapering, somewhat pointed, processes.

I learnt that the child's father was a coachman, and that there was a horse in his stable which had been certified by a veterinary surgeon to have ringworm, and that the child had been brought into contact with the horse. On paying a visit to the stable I found that there were two patches on the horse, one on the neck and one on the head, both scaly, and covered with broken and twisted hairs but showing no signs of vesiculation. Microscopic examination of these hairs showed chains of large spores outside the hairs, just as in the child, and cultures on the same medium proved to be almost identical with those obtained from the child's scalp. (Figs. 1 and 2).

The second case is also one in which infection was derived from a horse, but the species of fungus was here quite different. The patient was an ostler, aged 19, with a circular scaly patch, about an inch in diameter, on the right side of his neck. This patch was said to have been present a week, and it began as a small spot, which spread gradually at the edges while clearing in the centre. When seen the edge was red and somewhat raised, with a few small scattered papules here and there. Examination of scales from the lesion showed a network of abundant, fine, branching mycelium, irregularly septate. Fine lanugo hairs showed numerous round or oval small spores, and on careful focussing mycelial threads could be made out running chiefly in the direction of the long axis of the hair. The microscopic appearances, therefore, did not point to any species of trichophyton, but I was loth to make a diagnosis of microsporion until cultures had been made. These, however, left no room for doubt. The fungus grew with extreme rapidity, appearing first as a white downy growth, which rapidly spread over the surface of the medium until in sixteen days it had acquired the appearance shown in photograph No. 3. This culture closely resembles one of microsporion obtained directly from a horse by Bodin.

On inquiry, I found that the ostler had been looking after a young horse with some "pimples" on its nose, and on going to the mews I found that the animal had some indistinctly ringed vesicles above the right nostril, and the hairs around appeared to be irregular and broken. Microscopically they showed a sheath of small spores outside the hair, and the hair itself contained fine mycelium. Cultures gave a rapidly-growing white downy mass (Fig. 4) similar to that obtained from the lesion of the patient. Microscopic examination of the aerial hyphæ from the culture showed an unusual appearance, inasmuch as the organs of reproduction proved to be lateral bud-like growths, from 2 to 3 µ in maximum length, situated directly in the mycelial threads.

The third case was that of a boy, aged 3, with kerion of the scalp. There was a patch, about ¾ in. in diameter, on the left parietal region, crusted with yellowish scabs, which had been present for a fortnight. The hairs were broken, and could be very easily removed. They mostly came away without the bulb, and the hair shaft showed a typical felting with a mosaic of small spores, while inside the hair was mycelium, terminating in a fringe above the region of the bulb. Cultures showed alternate concentric rings of white and brown growth, with a delicate filiform border and a central raised prominence. (Fig. 5). Hairs from a terrier with which the child had often played proved to be infiltrated with typical microsporion, and cultures from these hairs resembled closely those obtained from the child's scalp. (Fig. 6, six weeks' growth.)

A fourth case is rather peculiar, in that the patient was only 4 weeks old when brought to the hospital, and had already a well-marked patch of tinea circinata, strictly localised and circular, on the left cheek. The surface of the lesion was red and somewhat scaly, with some vesicles and a few pustules at the spreading edge. Abundant large-spored fungus was found in scrapings and in the root sheath of the hair, and on cultivation it gave a somewhat coarse growth of radiating aerial hyphæ of a dull whitish colour. A cat in the house, with which the child played, had a patch on the neck, from which the hairs had almost all come out, and on getting the mother to bring the animal to the hospital I found that the hairs over this patch were extensively affected with ectothrix fungus. Cultures showed a beautiful white growth of very coarse hyphæ, which grew much more rapidly than that from the lesion on the child, and was more downy in appearance. The medium on which the fungus from the cat was grown was of exactly the same composition as that used for the

Fig. 1.—Culture from patient.

Fig. 2.—Culture from horse.

Fig. 3.—Culture from patient.

Fig. 4.—Culture from horse.

Fig. 5.—Culture from patient.

Fig. 6.—Culture from terrier.

Fig. 7. Culture from patient.

Fig. 8.—Culture from cat.

Fig. 9.—Culture from patient.

Fig. 10.—Culture from cat.

Fig. 11.—Culture from patient.

Fig. 12.—Culture from cat.

Fig. 13.—Culture from patient.

Fig. 14.—Culture from canary.

Fig. 15.—Culture from patient.

Fig. 16.—Culture from calf.

fungus from the child, but it had been freshly prepared, whereas the other had been kept a short time and had, perhaps, got a little dry. This may account for the greater luxuriance of the growth in the second culture. (Figs. 7 and 8.)

The fifth case was also one in which the source of infection had been a cat. The patient, a youth, aged 16, had several well-marked patches of *tinea circinata* on the left arm and trunk, two of which were slightly vesicular at the edge, but the rest scaly. A large-spored trichophyton was found both within and without the hairs of the lesions, and on cultivation gave a somewhat coarse white growth very similar to what has already been described by other observers as being derived from the cat. (Fig. 9.) From a patch on the patient's cat, which had been noticed to be "mangy," a very similar culture was obtained. (Fig. 10.)

The next case was rather exceptionally interesting, in that the fungus which gave rise to the infection was an endothrix, and an endothrix is supposed to have an animal origin but seldom. Indeed, this species of fungus can only be inoculated into animals with difficulty, but it can be inoculated on the guinea-pig, the rabbit, and the cat, and this being so, there can be no reason why human beings should not become infected from animals. The patient in whom this infection occurred was a boy, aged 9, who came under observation for two more or less circinate, smooth, almost bald patches on the scalp, which had been present for more than a month. The hairs were said to have come out rapidly at first, and the patches were never markedly scaly. When seen, there were scattered over the affected areas some stumps of hair, dark in colour, projecting a few millimetres above the surface of the scalp, and swollen slightly at the far extremity. Within these hairs were chains of large spores running chiefly in the direction of the long axis of the hair, but no fungus could be found outside the hairs, which came away easily, leaving the root sheath behind. The surface of these patches was so smooth that it was very difficult to scrape any of the superficial

epithelium away, but in none of the *débris* thus removed could any fungus be found. Cultures gave an irregularly plicated growth, somewhat powdery, and with a tendency to crack on the surface. (Fig. 11.) Hairs from a cat in the house also contained an endothrix fungus, but the surface of these lesions on the cat was not smooth, and was covered with fine scales. Cultures from these lesions on the cat proved to be very similar to those obtained from the boy. (Fig. 12.)

A case in which the nature of the fungus was open to some question was that of a girl aged 13, with a patch of *tinea circinata* on the forearm, the surface of the lesion being scaly and nowhere vesicular, though at the margin there were some red raised papules. Both in the scales and in the hairs was present a mycelium, rather delicate, and divided at irregular intervals by transverse septa, branching freely in its course. Cultures gave a dull, opaque, whitish growth, spreading somewhat irregularly from the centre, and with a faviform appearance (Fig. 13). The growth was quite unlike anything which I had obtained from either trichophyton or microsporon infections, but the clinical appearances differed in no way from an ordinary *tinea circinata*, and presented no possible similarity to favus. On inquiry, I found that there was a pet canary, whose feathers had been coming out in patches, and microscopic examination of these feathers showed an invasion by a fungus similar to that already demonstrated in the patient. The cultures, too, from the bird were faviform in appearance, and only differed in that the spreading edge was slightly more thickened. (Fig. 14.)

The last case was one of ringworm of the beard in a man aged 29. On the right side of the face were two small scabbed pustular lesions, over which the hairs were broken off within a short distance of the surface of the skin, and were fixed but loosely in the follicles. There was also a single red ringed patch with a pustular edge on the glabrous skin of the same side of the neck. When the hairs were pulled out, the root sheath often came away at the same time, and on microscopic examination this was found to be packed with chains

of large glistening spores, and at the mouth of the follicle a collarette of mycelium was in some instances present. But also within the hairs there was some fungus present, though there was no apparent stripping up of the cuticle. Cultures gave a whitish downy growth with a central prominence, and a fine striated edge (Fig. 15), but in many of the original tube inoculations the growths were contaminated by staphylococci, and it was only with difficulty that a pure culture was obtained.

The patient, who was a farm labourer, stated that one of the calves had ringworm, and brought me up some hairs from one of the patches. I did not see the lesions on the calf myself. These hairs also contained an endo-ectothrix fungus, somewhat closely packed, so that the chain formation was not very evident. Cultures from these hairs also gave a white aërial growth with a striated edge. (Fig. 16.)

These cases comprise, then, cases of ringworm in which the infection was derived from animals of various kinds, but I need hardly say that, in order to obtain even such a small number as eight, I have during the past three years been obliged to examine a considerable number of animals which had not ringworm, and to pay many fruitless visits to stables, mews, and other unsavoury spots. In some cases there was good reason to presume from the clinical and cultural appearances that the fungus had an animal origin, but I was unable to locate the animal. And this is not surprising, for the number of cases having such an origin must always be a very small percentage of the whole, and it must be remembered that one such case in a child is capable of serving as a source of infection to many more.

Valuable though microscopic examination is in enabling a diagnosis to be made, greater reliance must always be placed upon cultures, which either confirm the conclusions derived from the microscope or serve to differentiate those cases where otherwise a differential diagnosis could not be made. Such, for instance, are those cases where the spores are so closely packed that their shape necessarily appears round, and the microscope is quite unable to distinguish between microsporon and ectothrix.¹ Cultures, however, soon clear up the diagnosis.

The prognosis in the above cases varied very considerably, for though the inflammatory process was in most cases acute, yet it often yielded readily to treatment, in fact, some of the apparently milder cases proved in the end the most resistant.

Investigation of the organs of fructification in the trichophyton cases showed that they were either clusters of spores or conidia placed laterally on the mycelial hyphæ, or chlamydospores. Rarely intramycelial endoconidia from 10μ to 12μ in maximum diameter were found. The clusters of spores were usually borne close to the end of a mycelial thread, sometimes in considerable numbers, so as to form, as it were, a terminal inflorescence. The conidia were from 2μ to 4μ in length, and were borne either directly on the mycelium or on short unbranched pedicles. The chlamydospores or multilocular conidia were pyriform in shape, or somewhat fusiform, with a long diameter of from 25μ to 40μ , and enclosed in a capsule which did not stain with acid fuchsin. The contents were a granular protoplasmic material, divided into two or more loculi by means of transverse septa. These multilocular conidia are borne, like the unilocular conidia, either laterally on the mycelium, or at the end of short pedicles. In some cultures they are very numerous; in others they do not appear until the culture has grown for some time, and then only in small numbers as compared with the inflorescences of groups of spores. Chlamydospores are often numerous in old cultures, in which degenerated white downy growths appear, and here they seem to appear at the expense of the unilocular conidia. In the case of the microsporon derived from the horse, reproductive organs in the form of chlamydospores were also found, divided by one or more transverse septa, and containing granular protoplasm. At the end of some of the mycelial hyphæ occurred conidia somewhat cylindrical in shape, with occasionally transverse partitions dividing them up into loculi. Other reproductive organs found in these cultures were conidia borne laterally on the hyphæ in the form of bud-like processes, and measuring about 3μ in maximum diameter.

I have to thank Dr. Pringle for placing cases at my disposal

and for much valuable assistance in the research, and Dr. Sidney Martin for some kindly criticism and suggestions.

REFERENCE.

¹ Vide Fox and Blaxall, *Journ. Derm.*, February, 1898.

REMARKS

ON

FINSEN'S LIGHT TREATMENT OF LUPUS AND RODENT ULCER.

BY

MALCOLM MORRIS,

F.R.C.S. Edin.,

Surgeon, Skin Department,
St. Mary's Hospital.

and

S. ERNEST DORE,

M.B. Cantab.,

Clinical Assistant, Skin Department,
St. Mary's Hospital.

THE treatment of lupus vulgaris and other diseases of the skin by means of concentrated light rays has now become established as a method which may fairly claim to rank as second to none in importance and utility. Although we cannot as yet make any definite statements as to permanent cure, the results we have obtained have fully justified our expectations, and, backed by the larger experience in Copenhagen, we are confident in hoping that light used in this way will give sufferers from lupus and other intractable diseases of the skin not only better results but more permanent benefit than any forms of treatment hitherto employed.

SOME POINTS IN THE TECHNIQUE.

A detailed description of the apparatus and of the method of application of the rays has already been given by Dr. Bie (in the *BRITISH MEDICAL JOURNAL*), and others, so that it is only necessary here to give a few brief practical details which experience has taught us. The appliances we have in use are in every respect similar to those used by Professor Finsen of Copenhagen. It must be mentioned, however, that we have used electric light only, as sunlight, which is less efficacious on account of the relatively smaller proportion of violet and ultra-violet rays, is almost impracticable in this country.

Current.—We have found a current of about 75 ampères and about 60 volts sufficient to produce a good reaction in the majority of cases, and have preferred to lessen the time of exposure rather than diminish the strength of the current and intensity of the light.

Intensity of Light.—In order to get a strong light it is most important to see that (1) the lenses, especially the bottom one, is clean and bright; (2) the water is clear and free from floating particles; the bottom lens should be thoroughly cleaned and polished every day, and this is greatly facilitated by having it detachable as in the newer pattern. The water should be changed every day. The use of ordinary tap water (which is used in the compressing glasses) has been found to give as good results as distilled, and its theoretical disadvantages are more than compensated for by its practical utility; that is to say, it can be changed, and the lenses cleaned daily instead of every three or four days. If these practical details are attended to a good and—other things being equal—uniform reaction will be obtained.

The Focus.—The area treated is, as a rule, kept well within the focus of the light, but a smaller focus, if it can be borne, has a greater effect. We have found it convenient to use thumbscrews instead of the ordinary ones for focussing. It is very important that the tubes should be kept in a straight line with the light rays so that none of these are lost by impinging on the side of the telescope, we have found a new adjustment which gives a larger range of movement of service in this respect.

Screens.—It is also essential that the rays should fall perpendicularly upon the compressing glass, by the use of cardboard or metal screens fixed to the end of the tubes, the slightest deviation of the glass will throw light upon the screen, and this can then be rectified. This has been found very useful, especially when the attendant is inexperienced in the treatment.

Pressure.—We have continued to use elastics to fix on the