I have no doubt that our practice in this matter in renal disease has a more scientific basis than in many other instances. To limit the input of proteid material so as to bring the work of the diseased kidney within its capacity is to act on scientific lines, but we know so little of the more remote stages of nitrogenous metabolism that beyond this very general statement of our object we cannot safely go. Yet there is no disease in which the tyranny of diet has been so recklessly practised. To live for weeks and weeks on a purely milk diet is not, I am sure, in many cases, necessary, and in some cases it is distinctly harmful, and there is no clear evidence that it meets the scientific indication. Some personal experiments made by Dr. Engelbert Taylor in the Pepper Laboratory of Clinical Medicine¹ showed that the total output of nitrogen was greater on a purely milk diet than on an ordinary mixed diet. I only quote this to show that it can be demonstrated that so far as the urea output is concerned there is no advantage in a purely milk diet. Dr. Hale White has shown the same thing clinically in chronic Bright's disease. My own practice in the matter of diet is simplicity itself, and I believe in the great majority of cases it is advantageous for the patient. My rule is that if the bowels are acting freely the patient may live on such ordinary mixed diet, including meat, as he has an appetite for and can digest. Patients who are confined to bed and suffering from uræmic vomiting or diarrhoea, and are therefore getting towards the end of the disease, cannot of course have any appetite, and the difficulty is to contrive food of any kind for them.

In 1897 there was in the Leeds Infirmary for the whole year a man suffering from large white kidney (that was the clinical diagnosis). Dropsy was his chief and almost only distress. His abdomen was tapped for ascites no fewer than eleven times. At the end of twelve months his dropsy had entirely disappeared, and he was practically well so far as the symptoms were concerned. He returned to his work and continued at it for six months, then came back to the infirmary with dropsy as before, and was tapped again several times. During the whole time he was in the infirmary he lived on ordinary diet, and I may say he also lived upon the "white mixture," for that was his only medicine.

The distinctions which we draw between the different meats in dieting cases of renal disease and gout are to my mind ridiculous. A patient may eat chicken and fish, and not mutton; mutton and not beef, and so on. I am sure there is no reason for this. Pork which is white, and in the case of Alexis St. Martin, was shown to be the most digestible of meats, is usually entirely forbidden. Van Noorden² mentions the case of a patient with chronic parenchymatous nephritis, who took half-a-pound of poultry daily for five days, excreted the same amount of nitrogen, and a trifle more albumen than he did in the next five days, in which, instead of poultry he took an equivalent amount of nitrogen in beef. Water and other fluid sustenance should be freely taken. Alcohol is perhaps as well avoided if cardiac failure does not require it. Climatic treatment, when at the disposal of the patient, is an important aid, but I do not propose to discuss it in detail.

DRUGS IN CHRONIC BRIGHT'S DISEASE.

Of drugs, Cushny, in his *Pharmacology and Therapeutics*, says, "The period of constructive pharmacology has scarcely dawned; at present its chief function is destructive and critical, and it is due to the latter fact that the patient with Bright's disease in the next few years will have a much more endurable life."³ Still, of the many drugs which are in use in the treatment of Bright's disease, some are of great value. We have not, and we cannot have, any drug or other thing which can replace the kidney destroyed. We might have, and may have some day, some drug or something which will arrest the process of Bright's disease as distinguished from its results. Our therapeutic indications are at present entirely symptomatic. I have no great belief in what is known as the general treatment of Bright's disease by drugs, beyond the systematic use of purgatives to keep the bowels acting freely. Beyond that no drugs are necessary except for the treatment of special symptoms as they arise. I have no great faith in the so-called vicious actions of organs. The proper way out of the body for urea is the kidney, and neither the intestine nor the skin can be made to take its place efficiently. Diuretics such as digitalis are frequently of the greatest use,

and almost as frequently fail. Diaphoretic drugs—and of these pilocarpin is the only really active one—are sometimes unpleasant, not to say dangerous in their effects; and the hot bath and hot-air bath do all that can be done, and are strictly under our control. Vaso-dilators have very little, if any, diuretic action in chronic Bright's disease, but in dyspnoea, headache, and vomiting frequently give great relief. The nervous and respiratory disturbances of the more severe degree of renal toxæmia, which usually portend a not far distant death, exercise our therapeutic resources to the utmost. Bleeding, purging, and sweating are our usual methods of dealing with convulsions and coma, and I do not propose to refer to their use in detail.

The old practical rule—that in renal disease morphine or opium should not be given—is still one to be observed without hesitation in most cases; but there not infrequently arise respiratory and mental disturbances of the most distressing and painful kind, which morphine alone relieves, and in which it can be given with perfect safety. It is a very remarkable fact, and it has been frequently observed, that the manifestations of the uræmic state are singularly mutually exclusive. For instance, great dropsy may exist without any single toxæmic manifestation. (I think dropsy may in some sense be regarded as a toxæmic manifestation.) Albuminuric retinitis, causing absolute blindness, I have seen to be the only manifestation of ill-health apart from the urine. One patient becomes drowsy and comatose and dies; another becomes wildly maniacal—in both cases without a trace of dropsy. Dyspnoea, uræmic asthma of the toxæmic type—that is, unassociated with disease in the heart, lungs, or pleura—exists quite alone, sometimes for weeks together, a constant source of sleeplessness of the most distressing kind. It is in this wakeful, painful, non-soporific form of uræmia that morphine is of the greatest advantage, and may, when given with care, be used with perfect safety. I constantly use it in the treatment of this distressing dyspnoea and wakefulness, and no other drug relieves so completely and safely. The contraindication is that which common sense would suggest—namely, the presence of or tendency to coma. By stupor and coma produced by his own poisons the patient's sufferings ultimately come to an end, and there is no need for us then to administer a soporific in the shape of morphine.

We are much indebted to Dr. Stephen Mackenzie for pointing out the indications for breaking the old-established clinical rule—when they are clear we need not hesitate, when they are not we should never venture.

REFERENCES.

- ¹ *Amer. Jour. of Med. Sci.*, August, 1899, p. 141. ² *Munch. med. Woch.* No. 7, 1899, p. 214; and *Boston Medical and Surgical Journal*, November 16th, 1899, p. 498. ³ *Boston Medical and Surgical Journal*, November 16th, 1899, p. 498.

ON GENERALISED INFECTION IN GONORRHOEA.*

By ARTHUR H. WARD, F.R.C.S.,
Surgeon to the London Lock Hospital.

THE thesis which I have to develop, and shall hope to justify step by step, is as follows. The gonococcus in its process of growth in the human body produces an irritating toxin. This toxin is the direct cause of all the symptoms of the disease. In all cases it is absorbed into the system, where its presence causes systemic degenerations of varying degrees of severity. Gonorrhoea is thus a general toxæmic affection; but the microbes which form the toxin are generally localised on or around a mucous tract. The microbic invasion may extend to the organs communicating with the infected tract, or it may penetrate into the tissues, either by direct extension, as in the invasion of the peritoneum through the uterus and Fallopian tubes, or by a process of growth through the mucous membrane affected. Thence the infection may invade the cellular tissues, the lymphatics and glands, and the vascular system. This invasion is rendered possible by the action of the absorbed toxin upon the leucocytes, which is of a paralysing nature, and prevents the encapsulation of the microbes by these cells. Having reached the circulation the

* Report presented to the International Medical Congress held in Paris, August, 1900.

gonococci may invade the heart and endocardium, or may be carried to the peripheral capillaries. In these they become stranded and grow, producing more toxin, which sets up local inflammations. The microbes invade the joints, and are found in the synovial sacs, and also in the pleura and pericardium. They are probably present in the analogous inflammations of the tendons and periosteae. The invasion of the organism is favoured by all too energetic measures directed to the local infection, since they depress the local powers of resistance, and by abrading or lacerating the mucous surface, may directly open the door to the invasion. General treatment must vary according to the general conditions, and will differ when these are referred to toxæmia alone, or to toxæmia complicated by metastases. Local treatment is always required, and should always be free from instrumental, mechanical, or chemical violence.

THE TOXIN OF THE GONOCOCCUS AND ITS ABSORPTION.

The existence of this toxin has been demonstrated by J. de Christmas¹ who found that when separated from the culture by filtration, and injected into rabbits, it produced fever, wasting, anæmia, and death. Sterile abscesses were formed. When injected into the human urethra, suppurative urethritis followed in four hours. The epithelium was attacked. Gonococci were absent from the discharge; the attacked conferred no immunity. Toxins obtained from cultures of other cocci did not excite a similar urethritis. Again, Laitinen,² using sterilised cultures of this microbe, found that local and general reactions followed their injection into guinea-pigs. The animals died with signs of toxic poisoning; precipitation showed the presence of toxins. From these facts we may, I think, conclude that the acute local inflammations caused by the growth of the gonococcus on mucous membranes are directly due to the toxins produced; and that their absorption into the system is the direct cause of the *malaise*, rise of temperature, and loss of weight, more or less evident in all cases of acute gonorrhœa, and often marked in cases in which a local extension of the infection to the epididymis or other tract has taken place. In such a case with a larger area of microbic growth, a greater quantity of toxin must be developed and absorbed. I hold, indeed, that the local inflammation is direct evidence of toxic absorption, and that the fluid part of the discharge is the osmotic equivalent, and the pus cells the chemotactic equivalent, of the irritant absorbed. This view explains the clinical fact that when epididymitis takes place the urethral discharge diminishes. In this way, from the urethra the toxin passes into the capillaries by osmosis, and an equivalent of serum escapes. At the same time the cells of the capillary wall are irritated and leucocytes adhere to it; they thus come within the sphere of influence of the toxin, and by chemotaxis are caused to traverse the cell wall, and travel to the surface of the mucous membrane. But when the infection reaches the epididymis much toxin is produced there and is absorbed; this raises the toxin tension in the blood, and diminishes the tendency to osmosis and chemotaxis from the urethra, hence the discharge becomes less. When the infection of the epididymis subsides, the original conditions return, and the urethral discharge reappears.

But in exceptional cases the general toxæmia may have much more severe results. In prolonged gonorrhœa, in persons debilitated by drink, starvation, or disease, marked anæmia, a toxic degeneration of the blood cells, is found. Neisser,³ under systemic intoxications, mentions cachexia, nervous affections, optic neuritis, and erythematous, hæmorrhagic, and hyperkeratitic skin diseases. Allard⁴ records a case in point, in which multiple neuritis followed a badly-treated case of gonorrhœa in fifteen days. There was high-stepping gait, with muscular wasting of the legs and fibrillar tremor. Also diminished sensibility, with great pain on pressure, along the sciatic and anterior crural nerves. The knee-jerks were lost, the leg muscles showed incomplete reaction of degeneration. The local infection yielded to treatment, and slow improvement followed the use of electricity. Could there be a more complete clinical picture of a toxic degeneration in a neurotic subject?

Laeleneff⁵ states that the toxin of the gonococcus exercises a most destructive action on cellular protoplasm. He draws no distinction, however, between systemic degenerations due

to toxic absorption and the inflammatory changes or secondary fibrosis caused by a local growth of the microbe, whether by direct extension or by invasion of the organism and metastasis. He considers gonorrhœa to be a general infective disease, but I venture to think that, were this so, the evidence of general infection would be common instead of rare. The toxin is generalised in every case, it is a general toxæmic disease; but, beyond causing varying degrees of *malaise*, fever, anæmia, and cachexia, generally of small moment, this toxæmia rarely produces the alarming series of degenerations set down in his paper; such, at least, is my experience with the sturdy and well-fed English race.

THE INVASION OF THE TISSUES BY THE GONOCOCCUS.

Beyond the local extension of the infection, from the mucous tract of origin to contiguous passages, gland ducts, and the organs in connection with them, the interior of the body may be invaded by direct extension in the female. The peritoneum is reached through the uterus and Fallopian tubes. Cushing⁶ relates two such cases in which the presence of the gonococcus was demonstrated. He considers that his cases offer a convincing proof that diffuse peritonitis may be due to this cause. It is either rare or unrecognised, he says, and may depend on a specially receptive condition of the membrane or on the virulence of the organism. I venture to think that the idea of a special susceptibility of the leucocytes to the paralyzing effect of the toxin absorbed is the rational view, since the gonococcus, by continually passing from one organism to another, should, I suggest, attain a fixed virulence comparable to that acquired by the microbe of rabies when passed through a series of rabbits. Muscatello⁷ also quotes a case in point—that of a woman with old gonorrhœa persisting more or less for twenty-five years. She was admitted to hospital suffering from pyosalpinx and parametritis, with severe pain and fever. The pus was evacuated *per vaginam*. Three days afterwards general peritonitis set in, and the patient died. The pus, both in the tube and in the peritoneal cavity, contained gonococci.

These cases occupy an intermediate position between the local extensions of the infection and the invasion of the tissues through the epithelia. On this question Brewer⁸ states that the investigations of Bumm and others have shown that the gonococci very quickly penetrate the epithelial covering of the mucous membrane, and collect in great numbers in the lymph spaces of the upper connective tissue layer. He quotes Halsted's theory to the effect that although large numbers of microbes do penetrate the tissues, their development takes place at or near the surface of the mucous membrane, and consequently within the reach of injected fluids. Halsted thus tries to account for the clinical fact that the use of antiseptic injections undoubtedly does exert a marked effect on the disease, even when the gonococci are crowding the connective tissue spaces. To my mind it is incredible that this invasion can take place by anything but active microbic growth. I should explain the action of injections by saying that they wash away the toxin in the urethra and check its production there, and thus diminish the amount absorbed. Without its assistance the gonococci already in the tissues are unable to withstand the onslaught of the phagocytes, and are soon destroyed.

THE ACTION OF THE TOXIN ON THE LEUCOCYTES.

With regard to this inhibition of the phagocytes by toxin, Drobny's research⁹ is one of great interest. By a special method of staining he confirmed the view that the gonococci are often extracellular—that is to say, are found free as well as in the pus cells. When they are free the course of the urethritis is more severe and complicated than when they are within the cells. Out of 77 cases, 27 had complications, lymphangitis, balanitis, prostatitis, and rheumatism, and in all these 27 cases free gonococci were found. Drobny thinks it possible that the extracellular position of the cocci may be due to paralysis of the leucocytes by toxins, thereby rendering them incapable of surrounding the cocci. He thus confirms the observations of Podrez made in 1885.

This very important observation indicates, I think, the reason why, in exceptional cases, the gonococcus can make its way into the circulation, and so reach the cardiac valves

and more remote situations. In an individual whose phagocytes were exceptionally susceptible to toxic influences, the urethral infection would be imperfectly restrained, and thus become very acute. Much toxin would be formed and absorbed. The invasion of the tissues by the microbe would also be very active, and the resistance of the guardian wall of inflammatory cells imperfect. The microbes would thus grow into the lymphatics and small veins, and by one route or the other gain the circulation. Here the toxin would have preceded them and already inhibited the leucocytes generally; so the microbes would be enabled to grow on the cardiac valves should they offer a suitable nidus, or might be carried to the peripheral circulation, and there be stranded like sticks and leaves in a back-water, in the looped terminal capillaries of the synovial membranes, and the slender vessels of the periosteal tissues and tendons. The observation of Heiman¹⁰ is of interest in supporting this view. He examined 4 cases of gonorrhoeal arthritis for the microbe, and found it in two of them. One of these was that of a newborn infant whose arthritis of the wrist occurred two weeks after gonorrhoeal ophthalmia. The embryonic cells and tissues of a child may well be more susceptible to toxic poisoning than those of an adult. Further, this case shows the possibility of a general infection by the gonococcus, arising from a local focus other than the urethra. We have so far been able to trace the microbic invasion from a mucous surface through the membrane into the subcutaneous tissues and lymph spaces.

According to Brewer,⁸ the lymphatics and corresponding glands may next be involved. Lymphangitis is more common, and the inflamed vessels can be felt as large cords. There is but slight tendency to suppuration when the glands are affected; when this does occur it is probably due to a mixed infection. There seems no reason to doubt that the lymphatic system may thus be invaded.

THE GONOCOCCUS IN THE CIRCULATION.

The demonstration of the gonococcus in the blood itself is the next link in the chain of causation. This is furnished by an observation of Panichi,¹¹ who records two cases of gonorrhoeal rheumatism, in which good cultures were obtained from the blood. In both the disease was of old standing, and the urethral discharge contained numerous gonococci. Agar and human serum formed the culture medium. The microbes were not found in the blood either extracellularly or intracellularly. This I suggest may have been due either to the microbes existing in the blood in spore form, or in consequence of their comparative scarcity. Clinical experience shows that, as a rule, the symptoms of gonorrhoeal rheumatism are not acute. This suggests the idea that the microbes reach their nidus in sparse showers; and, consequently, produce but little toxin, and so set up only a chronic form of inflammation. Such a sparse invasion of the blood might easily escape microscopic detection, although cultures had been successfully obtained.

A much more acute process takes place when the microbes find a nidus in the heart, and the toxæmic symptoms are of corresponding severity in such cases. Ghon and Schlagenhauser¹² record a case in point; that of a young girl who had been suffering for four weeks with pains in the limbs, and other symptoms attributed to influenza. Four days before admission she began to suffer from rigors, which continued on admission to hospital. She was found to be suffering from acute gonorrhoea. Six days later acute pain in the right foot was followed by gangrene. Systolic murmurs were noted. The patient became rapidly worse, and died on the fourteenth day. At the necropsy ulcerative endocarditis with myocardial abscesses was found. The gangrene was caused by femoral embolism. There was an abscess in the peritoneal covering of the posterior surface of the uterus. Gonococci were found in the heart lesions, in the retro-uterine abscess, and in the cervical canal urethra, and vagina. The microbe found in the heart was tested by cultivation, and eventually by introduction into the human urethra. It was not present in the femoral embolism. The case was notable, in that many of the signs of malignant or pyæmic endocarditis were absent, such as splenic enlargement and septic emboli. There were no typical joint affections. The point of entry of the infection into the blood or lymphatic systems was not determined.

THE LOCAL EFFECTS OF THE TOXIN DEVELOPED IN METASTASES.

The presence of the gonococcus having been thus demonstrated in the blood and on the endocardium, there is no difficulty in understanding that it may occasionally be carried to many regions other than the usual points selected. Laeleneff, in the paper already quoted, mentions a case of pleurisy with effusion which contained the specific microbe. Here, again, is a terminal circulation favouring the stranding of the micro-organism. I am not aware that the gonococcus has been demonstrated as present in tendons, periosteum, or in the sclerotic coat of the eye, but in the presence of clinical signs similar to those caused by the microbes in the synovial sacs, there seems no reason to doubt that they are also present in the regions named. I hold that it is obvious that any localised inflammation or sclerosis must be due to toxin locally formed by microbic metastases; while degenerative lesions are due to the general toxæmia, whether absorption takes place from a mucous tract or from such a focus together with secondary metastatic foci. To dermatologists the degenerations indicated by erythematous, hæmorrhagic, and hyperkeratotic conditions of the skin are of most interest. These I suggest must always be due to general toxæmia, whether metastases coexist or not.

The importance of the action of the toxin on the leucocytes as favouring the invasion of the body, is indicated by the research of de Christmas already quoted. He showed that the gonococcus when injected into human beings and guinea-pigs, does not give rise to the specific disease, but dies in from twenty-four to forty-eight hours. Hence we may conclude, that for the successful invasion of the body, the gonococcus requires its way to be paved by the absorption of its toxin, produced locally. This explains the clinical fact of the necessity of attacking the local disease, and stopping the absorption of toxin from that source, in all cases of gonorrhoeal metastases. The toxin so absorbed will have been the cause of the original failure of the leucocytes to bar the invaders out; and with its withdrawal the inhibition of the leucocytes will be lessened, and the destruction of the microbes localised in the joints facilitated. I submit that the thesis I have brought forward receives much support from the foregoing considerations.

THE QUESTION OF PREVENTION AND CURE.

The treatment of the metastases of the gonococcus by drugs is so unsatisfactory, that perhaps the less said about it the better. Since the microbes are undoubtedly present locally an internal treatment by a drug of the germicide class is indicated. My belief is that quinine holds out the best hope of benefit, but if this fails it is well to try mercury, arsenic, or the salicylates. When, on the other hand, toxic degenerations are threatening, a course of treatment directed to the oxidation and elimination of the toxin is indicated. Potassium iodide internally, with quantities of water, with the idea of washing out the soluble poison, together with baths, massage, exercise, and change of air, with a view to its attenuation by the oxidation promoted by such measures will be of most service; but it is to the possibility of preventing these disastrous invasions that I wish to strongly draw attention. A good many years ago I saw a case which first opened my eyes to the dangers of too energetic measures directed to the destruction of the local infection. Briefly it was as follows:

A healthy young man of 20 was the subject of acute retention from spasm of the urethra. He was suffering from acute gonorrhoea. A catheter was passed and tied in; the profuse discharge saturated the tapes; the instrument was withdrawn on the third day. On the fifth day attempts were made to pass a metal instrument, which failed owing to spasm; slight bleeding resulted. A soft instrument was subsequently tied in. On the thirteenth day No. 10 English scale was easily passed under ether. On the twentieth day a hammer-toe was amputated: the wound sloughed, and cellulitis of the foot supervened; pus was evacuated by an incision. On the thirty-first day the foot was improved and the temperature normal. The patient passed No. 10 daily for himself. On the forty-sixth day he went to a convalescent home. Three days later he developed *malaise*, and a temperature of 103° F., with abdominal pain and a mitral systolic murmur. On the fifth day the temperature was 105° F. The heart sounds were heard all over the chest. The temperature remained very irregular till the seventeenth day, when he was better, and got up. There was a purulent urethral discharge. On the twenty-second day he had frequent chills, with a temperature of 104°. On the twenty-fifth day, left epididymitis occurred. On the twenty-seventh day the temperature again touched 105°. There was again a marked systolic *bruit* with vomiting and

emaciation. He again improved, but on the thirty-first day rigors again appeared. The epididymitis subsided, and the urethral discharge returned. He again improved, and got up. On the fortieth day he relapsed, with temperature of 104° , precordial pain and tumultuous action of the heart. After this slow improvement set in; the patient was markedly anæmic. He was sent to the seaside on the fifty-sixth day, suffering from gleet. The treatment was with quinine.

This very instructive case I hold to have been one of general infection by the gonococcus. The instrumentation of the urethra gave every facility for the invasion of the tissues, both by depressing the local resisting power, and by epithelial abrasion. It may be contended, of course, that the subsequent conditions were septic and secondary to the cellulitis of the foot, but this itself may have been due to gonococcal infection. Further, the cellulitis had ceased some eighteen days before the general infection developed, while the continued activity of the gonococcus was shown by the urethritis and epididymitis. The general symptoms also were more those of a gonococcal infection of the heart, than of a septic endocarditis. Anyway, this case strongly suggested to me the danger of favouring the generalisation of a gonorrhœa by applying to the urethra any measures calculated to injure or abrade the mucous surface, and so weaken its resisting power. Since then I have never permitted any instrument to be passed in the acute stage, nor have I employed high pressure injections or abortive antiseptics. Having realised the danger of causing the generalisation of the infection, I do not think such energetic measures wise, since through them a patient may be seriously imperilled or crippled for life. Consequently I believe, I have seen very few cases of gonorrhœal rheumatism in my out-patient practice. I cannot remember ever having seen a case of grave cardiac disease, or of the profound general toxic degenerations quoted by other observers. I have examined the records of 2,000 of my cases, and find 5 cases of gonorrhœal rheumatism on admission, and 7 occurring during treatment. Two others entered as "rheumatism" and "pain in the ankle" were considered to be true rheumatism, and were not specially treated. The remaining 12 were: Affection of knee 3, ankle 1, instep 1, plantar fascia 1, arthritis and scleritis 1, pains in joints 1, gonorrhœal rheumatism 4. This works out at 0.6 per cent. No prolonged or severe case occurred, but such a condition arising in an out-patient would probably lead him to seek admission to a hospital or infirmary, and the case would be missed.

The opportunity for error in statistics, drawn from the fluctuating population of the out-patient department, is very considerable, and only the broadest conclusions can safely be drawn from them. As the result of the policy of non-intervention, in the shape of energetic measures such as instrumentation, high-pressure injections, and strong abortive antiseptics, the fact remains, that only 12 in 2,000 cases of generalised infection occurred, and of these 5 were present when first seen. I may make a large allowance for error, and still contend that these results are better than those quoted by Besnier and Julien of 2 per cent., and Gresolle 2.8 per cent., or the statistics of Gricolle quoted by Neisser, in which he found 68 cases in 4,423, 1 in 63, or 1.5 per cent.

I venture to echo the words of Neisser, who writes: "It is surely beyond a doubt, that the frequency of complications depends on the method of treatment." I submit that the cases and observations I have brought forward clearly indicate that any local injury or irritation must favour the growth of the microbe, and increase the production of toxin, which is always absorbed. That this toxic absorption directly depresses the resisting power of the phagocytes, and favours the invasion of the organism by the gonococcus. In view of the extreme obstinacy of the generalised infection to treatment, we cannot afford to neglect any precaution calculated to minimise the danger of generalisation; and therefore it is of great importance to avoid any local injury of the mucous tract involved. Even in chronic gleet I have seen epididymitis once follow a too rapid dilatation of a stricture, and once too energetic use of the endoscope; and epididymitis is a frequent cause of sterility, which is no small matter. But with due care, my experience is that more vigorous treatment is both necessary and effectual in the chronic forms of gonorrhœa.

I have gratefully seized this opportunity of drawing the attention of the profession to the distinction between the

always generalised toxæmia of gonorrhœa and the rarer metastases, and the indications for the corresponding methods of treatment. If at the same time I have made evident the desirability of gentle local measures during the acute stage of gonorrhœa my labours will not have been in vain.

REFERENCES.

- 1 J. de Christmas, *Ann. de l'Inst. Pasteur*, No. 8, August 25th, 1897.
- 2 Laitinen, *Centrabl. f. Bakt.*, No. 20, 1898.
- 3 Neisser, *New York Med. News*, January 13th, 1900.
- 4 Allard, *Arch. d'Elect. Méd.*, Jun. 15th, 1898.
- 5 Laeleneff, *Vratch*, No. 4, 1898.
- 6 Cushing, *Bull. Johns Hopkins Hosp.*, 1899, p. 75.
- 7 Muscatello, *Il Policlinico*, August 15th, 1899.
- 8 Brewer, *Morrow's Syst. Gen.-Ur. Dis.*, vi, p. 154.
- 9 Drobny, *Arch. für Dermat. und Syph.*, October, 1898.
- 10 Helman, *New York Med. Rec.*, January 15th, 1898.
- 11 Panichi, *Seltmann Med.*, August 26th, 1899.
- 12 Ghon and Schlagenhauser, *Wien. klin. Woch.*, No. 24, 1898.

A CASE OF CONGENITAL HEPATIC CIRRHOSIS WITH OBLITERATIVE CHOLANGITIS (CONGENITAL OBLITERATION OF THE BILE DUCTS).

BY

H. D. ROLLESTON, and	LOUIS B. HAYNE,
M.D., F.R.C.P.,	M.D., B.C. Cantab.,
Physician and Lecturer on Pathology,	Formerly House-Physician,
St. George's Hospital; Senior	Victoria Hospital for
Physician to Out-Patients,	Children, and to
Victoria Hospital for	St. George's
Children.	Hospital.

The following case was under our observation in the Victoria Hospital for Children, Chelsea:

History.

A male child, aged 6 months, had been jaundiced since birth; it was treated as an out-patient with mercury and chalk, magnesium sulphate, and podophyllin. The jaundice varied from time to time, and the child's nutrition was fairly preserved. A fortnight before death the jaundice became more marked, and on May 28th, 1897, the child was admitted under Dr. Ridge Jones, to whom we are indebted for permission to publish the case.

Condition on Admission.

The child then had universal, but not extreme, jaundice. There was some erythema in the left axilla. The liver was much enlarged, and came down to the anterior superior spine of the ilium. The spleen was also enlarged, and projected three fingerbreadths below the costal arch. There was no ascites. The urine was bile-stained, and the motions clay-coloured.

On June 5th the temperature went up to 102° , and the child died after bringing up blood from the lungs.

The child was the first-born, and presented no signs of congenital syphilis.

Necropsy.

The necropsy was performed by Dr. Hayne. The body was thin, and all the organs and tissues were bile-stained. The œsophagus was normal, and free from staining by blood. The pleuræ were healthy. The trachea contained blood. Blood was found to have been aspirated into both lungs, which showed emphysematous bullæ and some small caseous masses, the latter were chiefly close to the surface of the lung. Microscopically these caseous areas showed broncho-pneumonia, with early caseation, but no definite evidence of tuberculosis. One of the tracheal glands was enlarged and caseous.

Condition of Liver.

The liver weighed 12 oz., or nearly twice the normal weight. Holt gives 7.5 oz. as the normal weight of the liver for a child of 6 months, while according to Birch-Hirschfeld 6 oz. is the average weight. It was much enlarged, yellow in colour, and manifestly cirrhotic. The common bile duct was small, and as its lower half was transformed into a slender fibrous cord considerable difficulty was experienced in finding it and in distinguishing it from the hepatic artery. The gall bladder was small, thickened, collapsed, and buried in adhesions; the cystic duct was represented by a thin fibrous cord. There were enlarged glands in the portal fissure suggesting the condition found in hypertrophic biliary cirrhosis. The liver cut like a cirrhotic liver and showed fibrosis.

Microscopic Examination.

Microscopic sections were taken from various parts of the liver, fibrosis was everywhere present, but the appearances varied in different situations. Where the fibrosis was least marked there was monolobular cirrhosis, the fibrous tissue being old and including a large number of newly-formed bile ducts; in other situations, where the cirrhosis was multilobular, there were areas of extensive fibrosis, including compressed liver cells and small bile ducts. Some of the latter were dilated and contained plugs of inspissated bile. The cirrhosis was everywhere old and not progressive. The liver cells inside the lobules were in a very fair state of nutrition, the trabecular arrangement was disturbed and the columns of liver cells were separated from each other by spaces, which, however, appear empty.

The appearances are, therefore, those of a mixed monolobular and multilobular cirrhosis. Sections of the common bile duct near its obliteration show very considerable fibrotic thickening of its walls, with complete alteration of its normal