A CLINICAL LECTURE

ON

SOME CASES ILLUSTRATING THE SURGERY OF THE LARGE INTESTINE.

Delivered at the Bristol General Hospital,

By CHARLES A. MORTON, F.R.C.S.,

Professor of Surgery in University College, Bristol; Surgeon to the General Hospital and the Hospital for Children and Women.

GENTLEMEN,—I propose in this lecture to bring under your notice some cases illustrating the surgery of the large intestine which have been in this hospital; and the first case to which I would call your attention is one of

VENTRIFIXATION OF THE SIGMOID FLEXURE.

This method of treating prolapse of all the coats of the rectum in an adult occurred to me as one likely to be successful, and I then looked up the subject and found it had been practised in several cases, and I propose to give you an account of the method and its results—results which encouraged me to adopt it.

This method of operating consists in opening the abdomen, drawing up the prolapsed bowel from within, and fixing it in the higher position. To the operation the name of colopexy has been given, but it seems to me the same name which is used to describe the fixation of the uterus for prolapse—ventrifixation—might be employed. The first operation of this kind was performed by Verneuil in 1889. He opened the peritoneal cavity, drew up the prolapsed rectum by traction on the sigmoid flexure, and fixed the latter to the wound in the abdominal wall by a silk suture which passed through several appendices epiploicæ. The appendices did not hold well, and a relapse occurred, but to a less extent than before operation. In 1890 Professor Macleod (of Calcutta) performed a somewhat similar operation, but incurred an unnecessary risk by transfixing the bowel. He fixed the colon to the abdominal wall by pins passed through this structure and then into the interior of the bowel, and then again through the abdominal wall, and thus incurred considerable risk of faecal leakage into the peritoneal cavity at the needle punctures. He did not open the peritoneal cavity, but guided the needles through the colon (or upper rectum) by his hand pushed up inside the bowel through the greatly relaxed sphincter ani. His patient had severe abdominal symptoms on the ninth and tenth day; but recovered. The case was reported too soon to judge of the ultimate result.

Before Verneuil's operation, in 1889, Jaennel, a French surgeon, opened the peritoneal cavity and drew up and fixed the colon; but he also made an artificial anus, and this proceeding seems to me quite unnecessary. Between 1890 and 1894 several Swedish, French, and Austrian surgeons performed the operation of ventrifixation, and in 1894 an English surgeon, Caddy, of Calcutta, did so with excellent result. In his case the meso-rectum was attached to the muscles and peritoneum of the abdominal wall. Several cases have also been recorded by American and Continental surgeons since 1894, and in the edition of Mr. Allingham's book on *Diseases of the Rectum*, published in 1896, he records one, in which severe prolapse in a woman was greatly reduced by this operation, though half an inch of bowel wall still protruded. This protrusion was successfully treated by the application of the cautery. The prolapse was eight inches in length before colopexy was performed. The mesentery was fixed to the "abdominal parietes in the neighbourhood of the incision in the abdominal wall." Dr. Bryant, of New York, who himself performed ventrifixation—but also formed an artificial anus—has published² collected records of 29 cases, and it is to his paper—which is by far the best account of the operation which I have been able to find—that I am indebted for some of the information I am able to give you in this lecture, and I propose now to tell you of the results in the cases which he collected.

The 3 cases in which an artificial anus was made I shall exclude, for I feel strongly that the formation of an artificial anus is not a satisfactory method of treating even severe rectal prolapse. This will reduce the number of cases to 26.

In all these cases the prolapse seems to have been readily reduced by traction on the sigmoid flexure. There was no death from the operation. Indeed, the risk of the operation is obviously almost *nil*, with due care as to the details of our antiseptic method of operating. The important question to decide is, What is the tendency to relapse? Only 13 cases are reported later than three months after operation, and these 13 alone seem to me of any value in deciding as to the frequency of relapse. Of these 13, 6 were reported six to ten months after operation, and of these Bryant says 3 had no recurrence, 1 a slight recurrence, and 2 a considerable recurrence; 7 cases were reported a year or more after operation, and of these 6 were cured. Another very important point which Bryant has worked out in connection with these cases is the attachment of the colon. The portion of colon sutured has varied considerably. Sometimes the peritoneum of the sigmoid flexure, or the band on it, has been used, sometimes the meso-colon, and, again, in a few cases the appendices epiploicæ. The attachment to the abdominal parietes has also varied greatly. The parietal peritoneum was often used; but the "abdominal wall" is also spoken of in distinction to the parietal peritoneum, and I fail to understand to what part the attachment was made in these cases. An attachment to the wound is also mentioned separately. In 1 case the colon was fixed to the lumbar aponeurosis, but this case relapsed. The method is not described. Bryant thinks the records do not establish the utility of any special method.

It seems to me that it is better not to pass the sutures into the wall of the bowel for fear of accidentally entering the lumen, but rather to suture the meso-sigmoid. The appendices epiploicæ alone have not proved reliable. With regard to the ventral attachment, I should be inclined to use the peritoneum in the neighbourhood of the wound, as easy to reach, but I should not fix the colon itself in the wound, for Bryant's own case shows that this is likely to lead to hernia there.

The results which I have just given you encouraged me to undertake the operation in the case of which I will presently give you the details. It was one of extensive prolapse of all the coats of the rectum. There was no evidence of any undue relaxation of the sphincter, and no tumour within the bowel, dragging it down, and hence it seemed to me that the great object in treatment should be to prevent the tendency of the bowel to fall out through the anus, from a deficiency in its attachment in the pelvis or abdomen. In other words, what was wanted was a better fixation for the rectum within the peritoneal cavity. As I have just pointed out to you, Dr. Bryant's collection of cases showed that the operation involved no risk, and indeed unless the sutures actually penetrated the interior of the bowel it was difficult to see how there could be any risk. Resection of the prolapsed portion of bowel was a much more serious operation. I did not feel at all sure linear cauterisation would succeed in this case; and it seemed to me that methods which narrow the anal orifice may lead to a condition of stricture. Moreover the operation proposed by Verneuil³ of drawing up the prolapsed rectum, and fixing it in the coccygeal region, only deals with the lower end, and in this and similar cases the whole rectum falls down too much, and ought, I think, to be fixed higher up than in the coccygeal region. I will now give you the details of the case.

Ventrifixation of Sigmoid Flexure for Prolapse of Rectum in an Adult.

Mrs. N. I., aged 24, was sent to me at this hospital by Dr. Flood in May last.

History.—For two years she had suffered from prolapse of the rectum. It commenced during the latter part of pregnancy, and was increasing at the time I saw her. There had been only very slight bleeding from the bowel at times.

Condition on Admission.—The prolapse occurred whenever she moved about, and a T bandage would not keep it up. The whole thickness of the rectal wall protruded for 3 inches. Strictly speaking it was an intussusception, for there was a sulcus half an inch in depth between the stretched anus and the protruding portion of the bowel. There were no piles, and no tumour was present in the rectum. It could be easily returned, and then the anal sphincter was found to contract well. She said it sometimes came down as large as her fist.

Operation.—On May 25th, when she was under ether, the prolapse continued. I took a line from the anterior superior spine to the umbilicus, and a point on this line, 1 inch from the anterior superior spine, and from this point made an incision to another 1 inch above Poupart's ligament just where the femoral artery lies beneath. The incision was therefore parallel with Poupart's ligament and about an inch above it. I thought this incision would give ready access

to the lower part of the sigmoid flexure. I divided the abdominal wall as if operating for recurrent appendicitis, so as to weaken it as little as possible. The external oblique was divided in the course of its fibres, and the internal oblique and transversalis at right angles to this, that is, in the direction of their fibres. The peritoneum was divided in the line of the skin incision. The sigmoid flexure, loop after loop, was drawn up out of the pelvis, and then as the lower loop was drawn on the prolapse disappeared. The upper part of the rectum was now on the stretch, and with the bowel in this position the meso-sigmoid at its lowest part was stitched with fine silk to the parietal peritoneum at the top of the wound in the following manner: Two silk threads were passed through the meso-sigmoid, at a distance of about an inch. By pulling on these a fold of meso-sigmoid was drawn forwards, and at the same time the bowel pushed back under the abdominal wall. The upper thread was then passed through the two edges of parietal peritoneum in the upper angle of the wound and tied, thus fixing the meso-sigmoid, and at the same time closing the upper angle of the wound. The lower thread was tied in the same way about the middle of the wound. Two or three other sutures were then passed through the edges of the parietal peritoneum, and the fold of meso-sigmoid in the centre, and tied. The lower part of the opening in the parietal peritoneum was closed by a few more sutures, and then the muscular layers of the abdominal wall carefully sutured together.

After-History.—She made an uninterrupted recovery, and when she left the hospital there was no sign of the prolapse. I examined her again a few days ago, and I found no more prolapse than one does in the prolapse of one moderate-sized internal pile; but she said that for some weeks there had been rather more prolapse than this during defæcation. The case could not therefore be regarded as cured, but as very greatly benefited by the operation.

Fig. 1.—Carcinoma of the cæcum; the excised portion of bowel. The small portion of bowel at the top of the specimen is the ileum, the large portion at the bottom the ascending colon. The cæcum is infiltrated by nodular growth. An inch measure is photographed by the side of the specimen.

EXCISION OF CÆCUM FOR MALIGNANT DISEASE.

The second case about which I shall speak to you to-day is of interest on account of the difficulty it presented in diagnosis, and the encouraging results of operation. It is sometimes very difficult to distinguish between a distended gall bladder and a movable kidney, but it is not often that a tumour of the cæcum gives rise to the same difficulty in diagnosis. This tumour could be pushed well back into the loin, and when the patient was under the anæsthetic, just before the operation was commenced, I could push it right under the umbilicus. It did not feel like a kidney in shape, but such free mobility was very suggestive of movable kidney, especially the way in which it seemed to fall back into the loin. Inflation of the colon with hydrogen gas would not have aided us in diagnosis, for the production of greater resonance in the position of the swelling would have seemed to indicate an overlying colon, and would therefore have favoured the view that the swelling was renal, whereas the increased resonance would have been within the swelling itself. This case shows the good results that may be obtained by excision of such a large mass of malignant growth within the abdomen, together with several affected glands.

A Case in which a Malignant Tumour of the Cæcum Simulated Movable Kidney: Excision of the Cæcum and Glands in Mesentery: No Recurrence Two Years after Operation.

The patient was a woman, aged 49.

Symptoms and Diagnosis.—Her symptoms were pain in the right loin and right side of the abdomen with occasional vomiting, but no symptoms of intestinal obstruction were present. The swelling was hard, not tender, and lay between the right anterior superior spine and the umbilicus, but rather nearer the former than the latter. It was about the size of the kidney, but not quite that shape, being broader at the top than below. It was not thought possible to make out the colon lying over it, and therefore it was not very resonant, if resonant at all.

Operation and After-History.—I operated in November, 1898, and first explored the loin to determine if it was a movable kidney. It was so movable that one of my colleagues even suggested that it must be a floating kidney with a meso-nephron. I will not give you details of the operation and after-treatment, further than to say that some glands in the meso-cæcum had to be removed with the cæcum, and that the ileum was united to the ascending colon with a Murphy's button. The growth in the cæcum was very extensive, as the photograph of the specimen shows* (Fig. 1). Microscopic section shows the structure of carcinoma. The patient is now (two years after the operation) in good health, and has no sign of recurrence. She is, indeed, far too fat, for she has actually doubled her weight since the operation.

OBSTRUCTION FROM MALIGNANT GROWTH OF COLON.

I will now call your attention to a common result of malignant growth in the large bowel, and the case is a good example of what should, if possible, be done in a condition of this kind. Excision of the growth and immediate union of the bowel is not a proceeding to be undertaken in cases of obstruction, for the patient is not in a condition to stand such an operation, and moreover the passage of enormous quantities of retained fæces through the intestinal junction will add greatly to the risk of leakage. It is wiser not even to excise the growth until the obstruction has been relieved. The less we do in the way of operation the better until the bowels have freely emptied themselves, and the patient's general condition has improved.

A Case of Intestinal Obstruction from Malignant Growth in the Colon: Colotomy and Fixation of the Tumour outside the Peritoneum: Subsequent removal of Tumour.

Mrs. M. M., aged 51, was seen by me in consultation with Dr. Maddison on June 1st, 1898.

History and Symptoms.—The symptoms of chronic intestinal obstruction had been present; the abdomen was much distended, and occasionally one particular coil would distend and harden with peristalsis. No growth could be felt, but from her age and the history malignant growth seemed the most probable cause of the obstruction.

Operations.—She came into the hospital on June 2nd, and on June 3rd I opened the abdomen over the descending colon, and found a constricting growth there. This I brought outside the abdomen, and fixed there with silk sutures. The following day I opened the protruding bowel above the growth, and thus relieved the obstruction; a fortnight later I excised the growth. No anæsthetic was required for this excision of the tumour. Two months later I closed the artificial anus by separating the ends of the bowel from the surrounding parts and sewing them together. At this time I discovered a nodule of growth on the left lobe of the liver. She left the hospital in October, and had not any serious difficulty with the bowels until December, and then obstruction at the bowel juncture was present and I had to form an artificial anus again. I believe this was due to kinking at the seat of junction. She never had obstruction after this, but the growth eventually recurred around the artificial anus, and she died in the spring of this year, one year and nine months after the removal of the primary growth.

REFERENCES AND NOTE.

¹ *Lancet*, 1890, vol. ii, p. 117. ² *Annals of Surgery*, 1897, vol. xxvi, p. 165. ³ *Bull. et Mem. de la Soc. Chir. de Paris*, t. xv, p. 754; and *Annals of Surgery*, 1891, vol. xiii, p. 218. ⁴ This specimen was shown at a meeting of the Bristol Medico-Chirurgical Society in the spring of 1899.

CHOLECYSTECTOMY: PARTIAL HEPATECTOMY AND PYLORECTOMY: RECOVERY.

By BERTRAM C. STEVENS, M.B., M.R.C.S.,
Leeds.

THE following case of cholelithiasis associated with malignant disease involving the liver, gall bladder, and pylorus, and complicated by an abscess in the abdominal wall, is, I think, of sufficient interest to report separately, and not in a group of cases. The case was under the care of Mr. Mayo Robson, whom I desire to thank for permission to record it.

Mrs. S., aged 54, was sent by her medical attendant, Dr. Fry, to consult Mr. Mayo Robson on July 28th, 1900.

History.—The previous history was rather indefinite, as the patient did not think it necessary to seek medical advice before April, 1900. She considered that she had been in excellent health until November, 1898, then for six months she suffered every month or six weeks from what was probably hepatic colic, though she said she was never jaundiced after the attacks. She also had pain at the epigastrium like indigestion. She had had on several occasions previously attacks of so-called spasms.