of length mentioned by Mr. Clarkson. It was supplied with blood by an anastomosis of vessels from the coats of the ileum and communicated freely with the ileum through a large opening. Arising as it did from the lower and anterior aspect of the gut there was no mesenteric attachment, and before it had adhered to the parietal peritoneum in the crural canal, it must have been freely movable in the abdominal cavity. By an unfortunate mistake the specimen was not available for microscopic examination, but though it was of rather smaller calibre than the ileum, its walls were only slightly thinner, and its appearance generally was similar to that of a piece of small intestine.

2. The chief danger of these persistent diverticula is usually considered to be the risk to the other abdominal contents of strangulation, on account of the presence of a fibrous band within the abdominal cavity. Here, however, it was the diverticulum itself which had established pathological relationships, producing strangulation of itself, and secondary inflammatory results on the contiguous bowel and peritoneum, which gave all the signs of complete obstruction of the bowel in its continuity.

3. It is interesting from a surgical point of view that this, evidently the most freely movable of the abdominal contents, should have been forced into the canal by the intra-abdominal pressure, and that, having become adherent at its extremity, a volvulus, apparently secondary in point of time, should have formed.

4. It is also curious to think that this patient had lived a long and active life, without any inconvenience arising from this irresponsible appendage in his abdominal cavity, only to succumb at last to results arising from its persistence.

CASE OF CANCER OF THE LARYNX WITH A LONG COURSE.

By E. DONALDSON, B.A.T.C.D., L.R.C.S.I.,
Surgeon Londonderry Eye, Ear, and Throat Hospital.

I AM induced to publish the main facts of this case on account of its long duration, nine years and one month, and because of the great difficulty experienced in diagnosis during the earlier part of the illness. For a long time a benign growth was diagnosed, and the disease was allowed to run its course unmodified by the more radical methods of treatment. The fact I wish to emphasise is that a warty laryngeal growth in an elderly person may present symptoms of a benign tumour for about eight years, and in the end turn out to be malignant.

History. Miss S. first complained of hoarseness in December, 1888. In the spring of 1889 she was under the care of a well-known surgeon, skilled in the use of the laryngoscope, for seven weeks. He found a sessile warty growth at the anterior part of the left vocal cord, which he removed with forceps, and had examined microscopically. The pieces removed were found to be of a non-malignant nature. The surgeon on June 28th, 1889, made a sketch of the larynx before and after treatment. There was only a slight scar visible on the anterior part of the left vocal cord after the treatment. She then lost her hoarseness and remained well until November, 1890, when the hoarseness returned and occasional slight difficulty in breathing was felt.

State on Examination.—I saw her for the first time on December 3rd, 1890. She was then 58 years old. I found that the wart had returned on the anterior part of the left vocal cord; it was pale in colour, sessile, and occupied nearly one-third of the cord. From this on she remained hoarse, and did not again completely recover her voice. In November, 1892, I found a growth on the right cord opposite that on the left. I could not say that there was any marked immobility of the cords. From time to time the growths were touched with ethylate of sodium, absolute alcohol, and chloride of zinc solution.

Progress.—Progress was very slow indeed, and there was little difference from month to month. In 1896 she was again under the care of the surgeon who originally treated her, and he was obliged on November 3rd of that year to perform tracheotomy on account of difficulty of breathing. After this operation she again came under my care, wearing a tracheotomy tube. The condition of the larynx did not improve, but became progressively worse. At the end of December, 1896, the left ary-epiglottic fold had become much swollen. There was a mere chink between the growths on the cords. I was now certain that the disease was malignant, and I communicated with the surgeon under whose care she had been. He wrote to say that he agreed with the diagnosis, but thought that extirpation of the larynx would not prolong life. In February, 1897, I found both ary-epiglottic folds swollen. She wrote on a slate what she wanted, and could not make any attempt at speaking. Swallowing was decidedly painful. She wrote that she could take a big drink better than a sip, and could swallow best with the head raised. She liked "to bolt" her milk. In May, 1897, there was a slight discharge of blood through the tube and round it; wasting and loss of strength were noticeable, but anæmia was not at all marked. For a short time before she died milk sometimes

came through the tracheotomy tube. She emaciated rapidly towards the end, and died of exhaustion on January 22nd, 1898.

Pathological Examination.—After death I removed the larynx for microscopical examination. There was very little swelling of the neck at the seat of disease. Opposite the tracheotomy wound there was a sloughy opening into the œsophagus. The larynx was so disorganised as to be unrecognisable. Round the tracheotomy opening the larynx and the tissues were in a sloughy, fœtid condition.

The microscope proved that the growth was an epithelioma. In regard to whether it was of this nature from the beginning, or became malignant some time during the nine odd years of its existence, I leave each one to form his own opinion.

FURUNCULOSIS OF THE EXTERNAL AUDITORY CANAL SIMULATING MASTOID PERIOSTITIS.

By JAMES GALBRAITH CONNAL, M.B., F.F.P.S.G.,

Lecturer on Aural Surgery, Anderson's College Medical School; Assistant Surgeon, Glasgow Ear Hospital, etc.

THE leading local symptoms of furunculosis of the external auditory canal—namely, pain in the ear, dullness of hearing, tinnitus, sometimes giddiness, and occasionally a scanty discharge from the ear—are well known. Of the constitutional symptoms a moderate degree of febrile reaction is common. There is, however, another type of case where, in addition to the above-mentioned symptoms, we have an œdematous swelling over the mastoid process, which is confusing and apt to lead to errors in diagnosis. The error is pardonable, for the clinical picture is that of a mastoid periostitis; so much so, that apart from a local examination of the external auditory canal a correct diagnosis and prognosis is hardly possible.

Fig. 1.—Boy, 8 years of age, with furuncle in the right external auditory canal, swelling over mastoid displacing the auricle downwards and forwards.

Anatomical Considerations.—To rightly understand this condition the anatomical structure of the outer ear must be borne