NOTE ON THE PATHOLOGY OF TOXIC AMBLYOPIAS.

By J. HERBERT PARSONS, B.S., B.Sc., F.R.C.S.,
Research Scholar of the British Medical Association.

It has been known since 1880¹ that the papillo-macular bundles of the optic nerves are degenerated in cases of tobacco amblyopia. These fibres are the axis-cylinder processes of the ganglion cells of the macular region of the retina, and hence changes in these cells are probably the cause of the degeneration. Sachs and de Schweinitz failed to find any microscopic evidences of such changes; but their observations were made before the introduction of Nissl's method of staining the intracellular granules. No case of tobacco amblyopia has yet been submitted to this test; but, on the one hand, the effect of nicotine upon the central nervous system has been investigated, and, on the other hand, changes have been described in the ganglion cells of the retina in other forms of toxic amblyopia.

To the second group belong those cases of experimental amblyopia produced by the administration of quinine, flin mas, carbon disulphide, etc., to animals. Distinct changes are found in the ganglion cells and nuclear layers of the retina, which Ward Holden² attributes largely to diminution in the blood supply due to vaso-constriction of the vessels, produced in the cases recorded by him by the administration of large doses of quinine to dogs. This and other forms of toxic amblyopia have been very carefully reinvestigated recently by Birch-Hirschfeld,³ who confirms the general results, though with some modifications.

The effect of nicotine upon nerve cells has received much attention from the experimental point of view, since Langley⁴ discovered that it paralysed them. The effect of the administration of nicotine to animals has also been investigated histologically in the cells of the central nervous system, with positive results, by Vas and Pándi.⁵

The paralyzing effect of the drug is obtained whether it is injected into the circulation or whether applied to the cells locally. It seemed to me to be worth while to investigate any changes produced in the cells by local application.

For this purpose the superior cervical ganglia of cats, which had been anaesthetised with ether, were removed immediately after death, and immersed for various periods from 1 to 3 hours in aqueous or normal saline solutions of nicotine of 1 per cent., 2.5 per cent., or 5 per cent. strength. Thin sections of the ganglia were compared with others of normal ganglia prepared and stained by identically the same process—namely, fixation in saturated corrosive sublimate solution, imbedding in paraffin, and staining with toluidine blue.

After being immersed for one hour in 1 per cent. solution of nicotine, which would be fully sufficient to paralyse the cells, they cannot be distinguished from normal cells. After more prolonged action or immersion in stronger solutions various changes are brought about which need not be referred to here.⁶

This supports the conclusion arrived at by Langley⁷ that the drug acts, not essentially upon the nerve cell, but upon the branchings (synapse) of the pre-ganglionic fibres around the cell. That there may be changes in the cell itself, which the delicate Nissl method is not sufficiently delicate to show, is, of course, possible.

The results are of some interest from the point of view of toxic amblyopia, for it would seem that if there are changes in the ganglion cells of the macula, they are a secondary result of changes in the fibres forming networks around them, that is, in the terminal fibrils of the inner granules, which form a network in the inner molecular layer. These, in turn, may be the result of changes in the outer molecular layer, in the fibrils of the cone fibres.

Warrington⁸ has shown that cutting off the afferent impulses to a nerve-cell is an even more potent cause of degeneration of the Nissl granules than cutting its own axis-cylinder process.

The action of nicotine may reasonably be supposed to be twofold: (1) Vascular, causing vaso-constriction of the arterioles, which would explain the selection of the sparsely-supplied macular region; (2) paralytic, upon the synapses either of the cone fibres, or of the inner granules, or of both.

BIBLIOGRAPHY.

- ¹ Samelsohn, *Graefe's Archiv*, 1882, Bd. xxviii, s. 1. ² Ward A. Holden, *Trans. Amer. Ophth. Soc.*, 1898. ³ Birch-Hirschfeld, *Graefe's Archiv*, 1900, Bd. 1. ⁴ Langley and Dickinson, *Proc. Roy. Soc.*, 1880, xli, p. 425. ⁵ Pándi, *Ungar Archiv. f. Med.*, ii, 1893, s. 257. ⁶ See *Proc. Phys. Soc., Journ. of Phys.*, xxvi, p. 38. ⁷ Langley, in Schäfer's *Textbook of Physiology*, ii, p. 621. ⁸ Warrington, *Journ. of Phys.*, xxiii, p. 114.

PARALYSIS OF THE CERVICAL SYMPATHETIC.

With Special Plate

By PURVES STEWART, M.A., M.D., M.R.C.P.,
Assistant Physician to the Westminster Hospital; late Physician to the
Imperial Yeomanry Field Hospital.

LESIONS of the cervical sympathetic are probably not so infrequent as the comparatively scanty references to them in current literature might lead us to expect. Many cases escape recognition because their signs and symptoms are not so obvious as in the paralysis of an ordinary cranial or spinal nerve.

The following case is unique, not so much because it exemplifies to a marked degree nearly all the classical symptoms of paralysis of the cervical sympathetic, but because it also demonstrates with great clearness the precise limits of one symptom in particular—anidrosis—with reference to which a considerable indefiniteness of statement has hitherto prevailed.

Fig. 1.—Area of anidrosis=right side of head, neck, and upper part of thorax, together with right upper extremity, bounded by thick black line. Area of cutaneous analgesia=shaded area of right upper extremity, back and front.

For the opportunity of studying the case I am indebted to the courtesy of Dr. W. H. Farmer, who, like myself, was a civilian medical officer *en route* home from South Africa on board the hospital ship *Simla*. Since our departure from the Cape the weather had been of tropical heat, and it had been observed that a certain patient sweated only on one side of his face.

History.

The patient, a man 30 years of age, is a soldier who was wounded in the left side of his neck two months and a-half previously by a Mauser bullet. At the time he was hit he was lying on his face, firing at the enemy. The bullet entered the left side of his neck an inch and a-half below the mastoid process, and came out through the seventh right interspace in the posterior axillary line, lodging in his bandolier. The bullet track thus passed in front of the vertebral column. At the time of the injury he felt an "electric shock" all over his body, but especially in his right upper extremity. The latter became at once totally paralysed. He also commenced to spit blood, and the hæmoptysis continued fairly copiously for three days, together with some difficulty in swallowing, the later being apparently referable to an injury of the œsophagus. On the fourth day his temperature became normal, and in three weeks the hæmoptysis entirely ceased.

The right upper limb, which was at first totally paralysed,

gradually recovered power, the return of motor function being preceded by stabbing pain in the ulnar side of the wrist. The motor power began to return about four weeks after the injury, commencing in the fingers, and gradually spreading up to the shoulder. He can now move the limb freely at all joints. He does not complain of any feeling of numbness (*vide infra*).

Ever since the injury, the vision in his right eye has been slightly impaired. On fixing a distant object, it appears to become slightly blurred, as if a haze came over the eye. He noticed, also, that he does not sweat on the right side of his face and neck, or in the right upper limb. He does not notice any difference in the amount of moisture in the two sides of his mouth.

Fig. 2.

Condition on Examination.

March 2, 1901. The patient is an intelligent man, of good physique. The entrance and exit wounds are visible, as already described, on the left side of the neck and in the right axillary region. There is no atrophy of any of the muscles of the right shoulder, arm, forearm, or hand. He can perform all movements of the limb at all joints, but less powerfully than on the left side. Opposition of the thumb is especially feeble. He cannot bring it to within half an inch of the little finger. The movements of the head and neck, of the left upper extremity, and of both lower limbs are powerful. The external ocular muscles, together with those of the jaws, face, palate, and tongue are normal. The diaphragm and intercostals also act normally. The knee-jerks are normal and equal, and there is no ankle clonus. The patient can feel the

lightest touches everywhere on the face, trunk, and limbs, but there is an area of slight analgesia along the inner side of the whole right upper extremity, from the axilla to the fingers. On the dorsum of the limb the analgesic area extends quite to the outer border of the arm and forearm; on the front its boundary runs down the centre of the limb. It splits the ring finger longitudinally, both on palmar and dorsal aspects. (See Figs. 2 and 3.)

Cervical sympathetic: The right pupil is smaller than the left. (See Fig. 4, Special Plate.) It contracts normally to light and on convergence; it does not dilate when shaded. The effect of instillation of cocaine was unfortunately not tried. The cilio-spinal reflex is present on the left side only. The right palpebral fissure is smaller than the left. (See Figs. 2 and 4.)

Fig. 3.

There is slight enophthalmos on the right side, the eyeball being a little sunken back.

The weather being exceedingly hot, he sweats profusely, except in an area on the right side of the face, scalp, neck, trunk, and the whole of the right upper limb. In those parts the skin remains dry. This area of anidrosis is bounded exactly by the middle line, and is well demonstrated by puffing powdered charcoal on the face and trunk. The charcoal sticks to the skin on the sweating side and blows off on the dry side, so that it is possible to photograph its boundaries. (See Figs. 1, 2, 3, 4.)

REMARKS.

The lesion in this case was evidently one which originally affected the whole brachial plexus on the right side, as evi-

denced by the total paralysis of the limb at the time of injury. The practically complete recovery of power, however (save in the opponens pollicis), together with the absence of muscular atrophy, shows that the damage to the greater part of the plexus did not amount to more than a contusion. A few fibres, however, appear to have been permanently interrupted, since there is slight analgesia of the areas to which the posterior roots of the eighth cervical and the first and second thoracic nerves are distributed, together with slight weakness of the small muscles of the thumb, which are innervated by the first thoracic anterior nerve root. The paralysis is therefore an example of the Klumpke type of partial plexus palsy—that is to say, it involves the lowest roots of the brachial plexus.

The most striking evidence, however, of injury to the lowest roots of the plexus consists in the total paralysis of the cervical sympathetic, which, as is well known, receives its oculo-pupillary fibres from the spinal cord through the rami communicantes of the first, and possibly also of the second, thoracic segment. The cervical sympathetic supplies the dilator pupillæ muscle, the non-striated part of the levator palpebræ, and the orbital muscle of Müller behind the globe. Hence in paralysis of this nerve we have narrowing of the pupil on the affected side, together with its absence of dilatation when shaded, or when the skin of the side of the neck is pinched (cilio-spinal reflex). There is also a narrowing of the palpebral fissure, a form of pseudo-ptosis due to paralysis of the non-striated muscle of the eyelid, since the upper lid can still be voluntarily elevated to its full extent by the striped portion of the levator palpebræ muscle. The sinking back of the eye (enophthalmos) is ascribed to paralysis of the orbital muscle of Müller.

In this case there was an area of anidrosis owing to paralysis of the sweat fibres of the cervical sympathetic. It was not due to a mere vasomotor condition, since subsequent hypodermic administration of pilocarpin (which stimulates sweat fibres without any action on vaso-dilator nerves) failed to produce sweating on the affected side, although it induced copious sweating on the healthy side.

The precise area of anidrosis is important, and has not, so far as I know, been hitherto recorded in any previous case. It should be observed that it is sharply limited by the middle line, that it extends as low as the third rib in front and the third dorsal spine behind, and that it includes the whole upper extremity on the affected side.

MEMORANDA:

MEDICAL, SURGICAL, OBSTETRICAL, THERAPEUTICAL, PATHOLOGICAL, ETC.

QUININE AS A PROPHYLACTIC TO INFLUENZA.

JUST as successful results as those published by Dr. Clayton in the BRITISH MEDICAL JOURNAL of May 25th regarding the prevention of malaria by the daily administration of small doses of quinine await any practitioner who will use the same drug as a preventive of influenza.

During the earlier years of my professional life, whenever influenza was prevalent I was surely attacked by it. Altogether, I had five attacks in four years. The next occasion I knew the disease to be active I commenced taking 2 grs. of quinine each morning with my breakfast, and have so done each succeeding period that the disease was prevalent, with the result that I have never since been attacked.

In the same way, whenever those I have advised to follow the same treatment have assiduously carried it out they have remained free from the disease, even although they had been previously sufferers from it. Many patients will object to the treatment, saying that quinine always sets up such unpleasant symptoms in them; but just as many drugs given in ordinary doses may produce symptoms of poisoning—and I have seen dangerous results from 1/200 gr. of atropine—so it is with quinine; in such cases, however, a minimum dose may be found insufficient to produce distress, yet enough to protect against the influenza poison.

Marple.

W. HABGOOD.

A CASE OF TETANUS: RECOVERY UNDER ANTITOXIN.

As the tetanus antitoxin is still more or less on its trial, the following case seems worthy of record.

B. C., aged 6 years, was admitted to the Prince Alfred Hospital, Sydney, on January 16th, 1899. He trod on a sharp piece of bone on New Year's Day. The wound suppurated, but healed after poulticing. On January 12th the parents noticed a "fixed" look about his face, and he complained of stiffness of the jaw. Next day the stiffness had extended to the neck and legs, and there were spasms of the face.

On admission the wound was found to be quite healed. The patient had trismus and stiffness of the legs. He was put on potassium bromide. Two days later he became rapidly worse. He could only separate the jaws by 1/4 inch; there was almost constant spasm of the muscles of the neck, arms, legs, and abdomen, and well-marked opisthotonos. The facial muscles were constantly twitching. The bromide was increased to 10 grs. every three hours, but as this had no effect, 5 c.cm. of antitoxin were injected, followed three hours later by 10 c.cm. There was a distinct diminution in the severity of the paroxysms as a result. Next day two more injections of 10 c.cm. were made, and the patient improved markedly. Slight spasms, especially of the jaw and face, continued for the next four or five days, and the injections were repeated; in all, 85 c.cm. was used. The patient was discharged quite recovered.

In this case the immediate improvement of the patient after the first two injections of the antitoxin was unmistakable.

Parramatta, N.S.W.

E. CUTHBERT HALL, M.B., Ch.M.

SUPRARENAL EXTRACT IN PROSTATIC HÆMORRHAGE.

UNDER the above heading in the BRITISH MEDICAL JOURNAL of May 25th, Dr. W. Habgood reports a case in which he used suprarenal extract for prostatic hæmorrhage, internally, for a man of 75 years, but was unable to continue its use because of its toxic effects. Last year I attended a similar case in a man of 80, who suffered somewhat from cardiac debility. I used an aqueous solution of suprarenal extract (5 grs. to 5j) locally, by injecting a few drops through an indiarubber catheter to the prostate, three times a day, with completely satisfactory results as regards the hæmorrhage, and without any deleterious general effects.

WALTER W. HEELAS, M.R.C.S., L.R.C.P.Lond.

Wolverhampton.

THE SORE THROAT OF MEASLES.

FACTS with regard to the spread of infection are so difficult to get that the following as to measles seem worthy of record.

A young man in service left his place not feeling well, and made for his home. He travelled by train to a town, where he saw two sisters, who were also in service. He slept in the town, and next day walked home eight miles. He felt so ill that he went straight to bed. He complained of sore throat, and his mother saw white patches on looking into his throat. Next day the rash of measles appeared.

It was on January 8th he reached his home, and on January 9th that the rash appeared—that is, on the fifth day of the disease. The two sisters whom he had seen on the third day of the disease both took measles and spread it to others. His brothers and sisters at home also took the disease. They began to be ill on January 15th—that is, seven days after the first exposure—and on January 19th, the eleventh day from exposure, the rash appeared. His home was at an isolated farmhouse, and there was no case of measles in the town where his sisters lived or in the vicinity of his home, but the children at the farm where he was hired had measles, and he took his meals in the house.

I saw in April a case of sore throat which, from the large surface covered by white membrane, was supposed to be diphtheria; but on the fifth day from the beginning of the illness the rash of measles appeared, showing the true nature of the case. The association of white membrane in such cases deserves to be noticed and studied more than it has hitherto been.

Denholm, N.B.

JOHN HADDON, M.D.

Fig. 4.—Paralysis of right cervical sympathetic from bullet wound of brachial plexus.