good growths generally speaking. There are some individuals who give a scantier growth than others, but not one instance has been found when the blood has completely destroyed the growth (Figs. 1 and 3).

Fig. 4.—Twenty-four-hours drop culture from the serum of a bubonic plague patient, who eventually recovered: second day of bubo. *B. pestis*: swollen and irregular. Beautiful involution forms: three seen in various stages as processes sticking out of the leucocytes.

The value of serum reaction above given is of scientific interest, but is of doubtful value, I think, from a diagnostic point of view, inasmuch as, while the observation takes at least twenty hours to give a pronounced opinion, the clinical features of the disease develop with such rapidity as to leave no doubt of the nature of the disease. Yet in a suspected or a convalescent new arrival a positive reaction—that is, absence of bacillus *pestis* in drop cultures—is of value in giving a definite information which may enable one to take preventive measures.

Fig. 5.—Twenty-four-hours drop culture of twenty-days' plague convalescent's serum. Absence of *B. pestis*. Two involution forms seen.

The practical value of the reaction in prognosis is also of doubtful nature, as the clinical course of the disease is, I think, in itself sufficient to guide the physician, yet a good or bad reaction is of interest to watch in the laboratory *in vitro* side by side with the clinical course of the disease *in vivo*.

While the practical value of the above observations is, as I have just indicated, of a doubtful nature from a clinical point of view, its application seems to give ample scope to work out the various problems, and the foremost is that of studying the reaction in Haffkinized subjects at the various stages, before and after the inoculation, and already a few observations have been made by me in conjunction with another worker, but as the material is not yet quite sufficient for publication we are not in a position to give our result just now.

The curative serums are also being tested, but owing to various samples not being within easy reach, I shall not give the result obtained from only one or two samples.

REFERENCE.

- ¹ Plague Com. Rept., 3rd para., p. 68.

THE Chelsea Hospital for Women has received a donation of £500 from H. M. E., who founded the Emergency Fund. The donation is in favour of that fund.

UNRECOGNIZED INFLUENZA.*

BY

THE LATE J. W. WASHBOURN, and J. W. H. EYRE,

C.M.G., M.D.,

M.D.,

Physician to Guy's Hospital and
the London Fever Hospital.

Bacteriologist to Charing Cross Hos-
pital, and Lecturer on Bacteriology
in the Medical School.

As is well known, influenza frequently escapes recognition: this is true not only in the mild cases but even in the fatal ones. In making observations upon broncho-pneumonia we have had occasion to examine a number of lungs, both of children and adults, and we have frequently found the influenza bacillus in cases in which the disease was not suspected. In all, we have examined the lungs from 12 cases, and from 4 of these we have isolated and subsequently identified the *B. influenzae*; in 4 others we have observed bacilli which were morphologically identical with the influenza bacillus, and also resembled that organism in such cultivations as were obtained, but which were not isolated and subjected to those tests necessary to establish their identity absolutely. These cases may be briefly described as follows:

CASE I (*B. P.*, 1) was that of a child 3 months old, who died of broncho-pneumonia. There were large areas of broncho-pneumonia in both lungs, especially at the right base. By inoculating a rabbit into the peritoneal cavity an impure cultivation was obtained from the blood; from this *B. influenzae* was isolated and subjected to the usual tests.

CASE II (*B. P.*, 2), a child 8 years of age, who died of broncho-pneumonia; *post mortem* scattered areas of broncho-pneumonia were observed in the right lung. Microscopical examination of the lung juice showed capsulated diplococci and some fine bacilli. A rabbit which was inoculated into the peritoneal cavity died in forty-eight hours, and from the heart blood a pure cultivation of the *B. influenzae* was obtained.

CASE III (*B. P.*, 3) was a lad 10 years old, who died of broncho-pneumonia following middle-ear suppuration associated with several small abscesses in the adjacent portions of the temporo-sphenoidal lobe. *Post mortem* both lungs contained numerous areas of broncho-pneumonia, some of which were necrotic and foul-smelling. Microscopical examination of the lung juice showed the presence of short, slender bacilli, resembling those of influenza, short chains of streptococci, and a spore-bearing bacillus. By culture methods an impure growth was obtained of the two bacilli, and subsequently the *B. influenzae* was isolated and identified.

CASE IV (*B. P.*, 4).—A male, aged 20 years, was admitted on November 28th into Guy's Hospital and died on December 6th. He had been invalided for three consecutive days on two occasions during the previous five months for cold and headache. The present illness began nine days before admission, that is to say, on November 19th, with headache and shivering. The headache continued, and on the 24th he had to give up work. On that evening and on the 26th and 27th he vomited several times. On admission the temperature was 101° F., pulse 80, respirations 24. He was delirious and irritable, his tongue dry and brown, and he complained of severe frontal headache. The respiratory and circulatory systems were normal but the pulse was slow in relation to the temperature; there was no paralysis, and the knee-jerks were present; the pupils reacted; there was optic neuritis in both eyes. The patient gradually became comatose, the temperature keeping at a fairly constant level of about 103° F., and the pulse varying between 80 and 90. On December 4th he looked dusky, and rales were heard at both bases. On December 5th the respirations were increased to 44; he died on the 6th. At the necropsy the bone in the pituitary fossa was found to be bare, and the pituitary body itself was difficult; there was some clotting in the lateral sinus. The brain was hardened in formalin and subsequently carefully examined but nothing abnormal was observed. There were extensive areas of broncho-pneumonia in both lungs; in the juice of the lung streptococci, diplococci, and fine bacilli were found. Two rabbits inoculated from the lung juice died in two and five days respectively; nothing was found in the heart blood of either. An impure cultivation was obtained from the lung juice, and from this the *B. influenzae* was subsequently isolated.

CASE V (*B. P.*, 5).—An infant who died from broncho-pneumonia following upon measles; in both lungs there were patches of broncho-pneumonia with pleurisy. The microscopical examination of the lung juice showed small bacilli like those of influenza, and cocci like pneumococci. By means of cultivation upon blood agar an impure cultivation of pneumococci and influenza bacilli was obtained.

CASE VI (*B. P.*, 6).—An infant a year and a half old, died of broncho-pneumonia; there was a consolidated area at the base of the upper lobe at the right side, with scattered areas through the middle and lower lobes. A microscopical examination of the lung juice showed the presence of capsulated diplococci and minute bacilli like those of influenza. By means of cultivations from the gland and from the lung an impure cultivation was obtained, and among the bacilli we found those resembling *B. influenzae*.

CASE VII (*B. P.*, 7).—A child, aged 4 years. *Post mortem*, diffuse broncho-pneumonia was present; in the lung juice there were a few cocci, and many small bacilli looking like the influenza bacillus. By means of cultivation we obtained an impure growth of staphylococci and bacilli like the *B. influenzae*.

CASE VIII (*B. P.*, 8).—An infant 4 months old; diffuse areas of broncho-pneumonia scattered throughout the lung were noted at the necropsy.

*The publication of this paper has been delayed for some months owing to the unhappily fatal illness of Dr. Washbourn.

Microscopical examination of the lung juice showed fine bacilli and a few diplococci; cultivations yielded an impure growth of what appeared to be the *B. influenzae*. A rabbit which was inoculated into the peritoneal cavity with 0.2 c.cm. of the lung juice died seven days later, but culture media inoculated from the heart blood, etc., remained sterile.

A point of great interest bearing upon the pathogenesis of the *B. influenzae* was observed during the progress of this work. Pfeiffer, who originally described the organism, stated that it was pathogenic for rabbits when injected intravenously, such inoculation being followed by paralysis of the hind limbs and in some cases death, but in no instance did he obtain effects resembling septicaemic infection. Cantani has also shown that it is possible to infect rabbits by means of intravenous injection, and his experiments bear witness to the powerful effects of the toxin elaborated by the bacillus. In two of our cases (*B. P.*, 1 and *B. P.*, 10) the intraperitoneal injection of material containing the *B. influenzae* was followed by death in about forty-eight hours from septicaemia, and the influenza bacillus was obtained direct from the heart blood, in one case associated with staphylococcus albus, from which it was easily separated; in the other in a state of purity. Its presence was also demonstrated microscopically in smear preparations from the spleen pulp. On the other hand, intraperitoneal inoculation of even large doses of early generations of the *B. influenzae* subcultivated upon blood agar failed to induce general infection and death.

ON "DYSPHASIA" OR APHASIA AS AN INITIAL SYMPTOM OF TUBERCULOUS MENINGITIS.

By M. MCINTYRE SINCLAIR, M.D. GLASG., D.P.H. CAMB.,
Assistant Medical Officer, Cotswold Sanatorium for Tuberculosis, Stroud;
late Resident Physician, Western Infirmary, Glasgow; etc.

TUBERCULOUS meningitis is a disease which, as every clinical student knows, is protean in its manner of onset. In reference to this affection, Bristowe,¹ in his work on medicine, writes: "Sometimes the first indications of disease are furnished by dullness, strangeness or wildness of manner, by impairment of memory or defect of speech." The pathology of the disease, moreover, is such that the occurrence of an aphasia as a primary symptom might at first sight be regarded as no unlikely contingency, yet while I have no doubt many such cases must at one time and another have been under observation, I find the records of them remarkably few. I have, however, in the course of my reading been able to find at least two histories similar to the one which I wish to record, illustrating the occurrence of a definite aphasia as the initial symptom of tuberculous meningitis. One of these is recorded by Schutz in the *Prager med. Woch.* for 1881, and the other by Déjerine in the *Rev. de Méd.* for 1885. The latter is further interesting in that, like my own case, the aphasia was ushered in by several attacks of a transitory character; but I will give a summary of both cases after first detailing the one which came under my own notice.

CASE I.—(PERSONAL OBSERVATION)

Transitory Aphasia occurring as a prelude to Tuberculous Meningitis.

J. B., aged 28, came under my notice in October, 1899, suffering from a chronic type of phthisis pulmonalis. He had occasional attacks of hæmoptysis, but for the first nine months he was under observation always recovered sufficiently to undertake light work between the exacerbations of his illness. Latterly the wasting and hectic fever became more marked. On December 7th, 1899, I was called suddenly to attend him for another attack of hæmoptysis somewhat severe in character. Under treatment he rallied, but remained somewhat blanched in appearance. In the course of a few days' time all traces of blood had left the sputum, and on December 11th he had every promise of recovering, temporarily at least, from the immediate dangers that threatened him. Three days later, however (December 14th), I received another urgent message from his wife stating her husband was apparently going out of his mind, and the history elicited on arrival was somewhat as follows: Mr. B. shortly after awaking that morning had become strange in his manner, and spoken in a peculiar way; his wife at first attributed this to delirium such as he had had previously, but when she could not understand her husband's speech he became more incoherent and excited than ever. At last it struck her that he wanted to know what time it was—a common question of his in the morning—and so apparently he did, for on holding the clock in front of him he looked at it, nodded assent, and quietly went off to sleep again. He had a similar incoherent attack on awakening later in the forenoon.

I saw him at 11 a.m. and found him sitting up in bed, and first appearances did not indicate anything unusual apart from his previous history. I was surprised to find that he could answer my questions regarding his general health, as to how he had slept, etc., in quite a natural tone and manner. We continued talking for some minutes, as I purposely wished by so doing to

bring out any speech defect such as his wife had indicated. In a short time the flaw became apparent. He had kept up an intelligent conversation for I should think three or four minutes, when his words became indistinct and blurred, running into each other and getting so inarticulate that, failing to understand him, I turned to his wife, and looked questioning at her. She, being more accustomed to his talk, could still make out the drift of his conversation, while I was completely puzzled, but in a few moments more his talk had degenerated into such an unintelligible jargon, that not even his own wife could make out a single word. Seeing from our looks and remarks that we failed to comprehend him, Mr. B. endeavoured all the more to say what was on his mind, and in his anxiety to do so came out with the jargon in such an eager, persevering, and excited manner, that I at once saw how his wife had imagined he was going off his head earlier in the day. I now told him we could not make out his speech, but that as he seemed anxious to tell us something, perhaps he could write it down. He perfectly understood what we were saying, and he took pencil and paper in a way that suggested he had a very easy task to perform. This, however, was what he wrote:

Fig. 1.

Looking puzzled, he commenced a fresh line more slowly, again the result:

Fig. 2.

at which he looked in a helpless kind of way, and then handed it over to me to see if I could make anything of it. He then started off again with his mumbling attempt at speech. I allowed him to go on for a few minutes, and then turning over a blank sheet of paper so that he could not see what he had written, I again made an experiment with his writing, and told him to write down what he wished to tell us. This time he wrote in a more deliberate manner, but with a very similar result:

Fig. 3.

and then, looking puzzled, gave up further attempts. It was plainly a case of what Professor Gairdner had termed "intoxication" with a group of letters, and evidently nothing else was to be got out of him. As the patient was by this time somewhat exhausted I desisted from further examination, and ordered absolute rest till I saw him again. His wife reported later in the day that he had remained quiet for about an hour after I left; by that time he had recovered his speech just as he did after the previous attacks. He now voluntarily commenced a conversation with his wife, remarking that the doctor had been seeing him and that he knew he had been unable to express his meaning properly at the time. He also knew that he had attempted to write something down, and asked to see the paper on which he had written. His remarks therefore proved that he was fully alive to all that had taken place at the time of his aphasic attack. Dr. Clarke saw the patient with me later in the day, our observations confirmed those of the morning's visit, and after speaking correctly for a few minutes his speech again quickly degenerated into mere "gibberish." We also determined the fact that when he spoke quite well (for example, at the commencement of our conversation) he wrote quite well. His hearing was good, and there was no "word-deafness" nor "word-blindness."

On the following day, December 15th, I again saw him, and there now was a very material improvement. He conversed with me for a very prolonged period without any flaw becoming apparent. I also showed him his writing of yesterday. He said he recognized the paper he had written on—otherwise he could hardly believe he had written it. He seemed highly amused at the composition but could give no explanation of what the letters stood for. He remembered perfectly well what he wished to tell me at the time his speech failed him; it was in regard to the details of a previous illness which, however, had no material bearing on the present attack. It is clear, therefore, that in his aphasic attacks he knew what he wanted to say but was unable to say it.

His condition now was decidedly improved as compared with that of the preceding day, there was no mental flaw, no headache and no pyrexia. Indeed it looked as if the patient were going to make an, at least, temporary recovery, but fortunately no prognosis was given.

I say "fortunately" because on the same evening insidious symptoms made their appearance, then more alarming ones, and within a week the man was dead from a clear attack of tuberculous meningitis. First he began to complain of severe headache, then his memory at times became defective; like amnesia generally, this first exhibited itself in the non-remembrance of proper names, that of his only child "Dolly" having to be constantly recalled to him.

I need not describe in detail the further progress of the case, the disturbance of vision, the alterations in pulse and tem-