A Clinical Lecture

ON

THE TREATMENT OF PNEUMONIA.

Delivered at St. Bartholomew's Hospital,

By SIR DYCE DUCKWORTH, M.D., F.R.C.P.,
Physician to the Hospital, and Lecturer on Clinical Medicine.

GENTLEMEN,—I am going to speak to you about the treatment of pneumonia, or rather the treatment of cases suffering from pneumonia. I have often called your attention to the fact that we do not treat diseases, but sick people who suffer from diseases.

The treatment of pneumonia is a topic which has excited a great deal of attention. There has been in the past a large amount of writing, and much temper excited, about the treatment of pneumonia. Various theories have been held for many years with regard to the treatment of the disease, but the questions raised on the point are more set at rest now than they were at the beginning of my student days. At that time pneumonia was generally looked upon as a local inflammation of the lung; patients, having caught cold and been exposed in some way, suffered from a local inflammation. The idea was then to get rid of this inflammation, to destroy its influence, so that the patient might again recover his wonted health. In accordance with that view—I am now talking of a century ago and up to within the last forty years—this result was to be achieved by such treatment as bleeding, repeated again and again, and the use of mercury, calomel more especially, and a very low diet. In that way the disease was said to be jugulated or strangled. That method of treatment was called "heroic," for naturally it was found to take the vigour out of anybody, however strong he might be, and unfortunately under it a large number of people died. It was thought not to be a bad result if perhaps one person in three recovered from pneumonia, which was at that time considered to be a very deadly disease. Gradually the knowledge of the subject increased, and it has continued to do so, until we have been enabled to learn what the natural history of pneumonia is. In the Charity Hospital at Vienna it was found that pneumonia, treated with rest and careful dieting, did much better than when submitted to bleeding and powerful drugs, and the percentage of recoveries was very satisfactory. These results opened the eyes of physicians in this country and elsewhere who were not tied down by tradition, and they ventured to give up the so-called "heroic" treatment of pneumonia. The modern treatment of the disease was practised in Vienna by Skoda and Dietl more particularly, and it was studied there by Dr. George Balfour, of Edinburgh, who was much struck by the extraordinary results of it as compared with that which he had been accustomed to see. He made known in this country the principles upon which the new treatment was based, and they were adopted by him and also by Hughes Bennett, who was a great clinical teacher in his time in Edinburgh. These physicians, together with Todd, in London, and others, confuted the prevalent opinion that the disease was to be strangled by bleeding and drugs, and they adopted what they called a supporting line of treatment, and thereby reduced the mortality in pneumonia from 1 in 3 cases to 1 in 26, certainly a very great improvement.

These changes were not adopted without a good deal of friction. The older physicians could not understand the better results that followed the new treatment, and they could only explain them by asserting that pneumonia had changed its type, and that it was not so severe or dangerous a disease as it previously was. They declared that in their day the patients were stronger and had harder pulses. That was as much as to say that the change of type indicated a diminution of human energy and life force. But that idea was contradicted by many facts. Veterinary surgeons began to practise upon the same principles, and gave up the heroic treatment in regard to horses. But no one could assert that horses had degenerated, for they were as good as ever they were, if not better. In the same way no one could say that men had become inferior in stamina and strength. It was

absurd to say that the race was degenerating, or that the disease presented a less sthenic type than it used to do.

But it is not too much to say that pneumonia may present different types from time to time. You will meet with more severe forms of pneumonia at one time than at another. That was recognized, for instance, during the prevalence of the recent influenzal epidemics. The type of pneumonia was then sometimes very severe. Again, certain epidemics appear to intensify the type of pneumonia, so that whatever be the line of treatment, more people are likely to succumb when the disease is more virulent than when it is less so.

It so happened that I was house-physician in Edinburgh to Hughes Bennett for some time, and had charge of many of his patients, and it occurs to me now that the cases which I saw then were certainly not so severe and deadly as those which I have been seeing within the last few years. The type of pneumonia just now is a little more severe, I think, and seems to prove more often fatal than at the time when I was in Edinburgh, but I think this increased severity in pneumonia is to be explained by the fact that at times the disease assumes a more virulent type than at others. This may explain the fallacy of statistics drawn from conditions which are not constant or fairly comparable.

The natural history of pneumonia shows that it is an acute disease, that it is truly an inflammation of the lung, but that it is something more. Although it is a local inflammation it is a specific fever. That is perfectly certain; everything points to that. Nothing points to it more certainly than the fact that, as we now believe, pneumonia is in most instances due to the influence of a toxin which is generated by two or three specific microbes which evidently are at the bottom of the whole process. The commonest microbe found in pneumonia is Fraenkel's diplococcus. In other cases the streptococcus and the staphylococcus have been found, and hence people sometimes speak of streptococcus and staphylococcus pneumonia. But the morbid anatomy is practically the same in the deadhouse whichever organism is present. The character of the infecting microbe itself is a matter for skilled bacteriologists to determine. Therefore I advise you for clinical purposes to study pneumonia, as all other diseases, not so much with regard to its microscopical element, but at the bedside, looking at the patient, taking notice of his state, observing what you see, finding out the individual degree of vitality, and doing everything you can, as a result of that knowledge, to help him through his illness. In other places you can go into the details as to what organisms are associated with the disease.

What is to be done for the patient who presents the early symptoms of pneumonia? These consist in a feeling of weariness, a bad frontal headache, and shivering, with which the disease very often starts. The patient is too ill for his work and goes to bed. There may be nothing yet in his condition which points to inflammation of the lungs. The temperature rises to about 103°, and you examine the lungs, but you perhaps find nothing noteworthy there. You do not yet know what is the matter, and you may perhaps think of typhoid fever or expect some exanthem to come out. This condition of things may go on for a day or two without any localizing signs in the lung. But the experienced eye will detect symptoms worthy of note even at that early stage. A flushed face, rapid breathing, a quick pulse, a high temperature, in the absence of other things which cannot be detected, should certainly make you think of acute pneumonia in such a case. And so you examine the patient day after day until you find something definite.

The signs which indicate a gradual progressive consolidation of the lung are an insidious increase in the frequency of breathing, a high state of fever, the pulse and respiration reaching usually the proportion of one to two—that is to say, instead of there being one respiration to four or five beats of the pulse, the proportion is one to two respectively. The amount of respiration is in relation to the amount of lung involved by the disease.

Then there is the short cough and the expectoration of blood-stained mucus or rusty sputa, sticky and tenacious. When that is present the diagnosis is practically certain. The patient's state is one of fever; he has a pungent skin, and on the third or fourth day, sometimes before, there is apt to come out a rash on the face, nose, or on the cheek or

chin, patches of herpes. This is exceedingly common in pneumonia; you will promptly conclude that a case is one of pneumonia when you find a febrile state of the patient with rapid breathing and herpes. This last-mentioned symptom is of some prognostic significance, because it is found in the majority of cases that herpes on the lips or face is a favourable symptom. So that in any given case one is rather pleased to meet with herpes labialis than not. Some of the worst cases show no herpes; but some grave cases may have it and the patients may die, therefore herpes labialis is not necessarily a favourable indication. On the whole, however, more cases with herpes labialis end promptly and favourably. This symptom does not occur in enteric fever. You may also regard it as of favourable omen in a given case if there be, on examining the blood, some degree of leucocytosis.

The late Sir William Gull used to say that the essential thing for a pneumonia patient was to put him in a warm bed. He did not much favour any drugs, but he did favour good nursing. It is certainly an important part of the treatment to put these patients to bed. A warm bed and good nursing are essential parts of the treatment. Then, again, a symptom of the disease consists in loss of appetite, so that there is no object in forcing food. They are likely to do very well on what is called a fever diet, consisting mainly of milk and beef-tea. If a patient can take that the great probability is that he will do well unless some urgent symptoms set in. The high temperature of 102° or 103° appears to be a necessary part of the disease and one way out of it; the patient cannot get well without fever, therefore you do not take any means to check it as long as it keeps within limits. But the temperature may rise to a higher degree than that in pneumonia, just as it may in typhoid fever. It may reach 104° , 105° , or 106° , and anything over 105° you may call hyperpyrexia, and this requires attention, as it is a dangerous condition. You must take means to reduce that temperature, and this is best done by sponging with ice-water and placing a cradle under the bed coverings. If that is not enough you should suspend small buckets of ice from the cradle, and so keep the patient in a cooling atmosphere. If that is insufficient you will employ Leiter's tubes in which to run ice-water over the head, or put ice in bags between the axillae and the legs. If that also fail you may give the patient some drug, of which the best is quinine. There are many modern drugs in repute for reducing the temperature, but some are apt to be mischievous, such as antipyrin, antifebrin, and phenacetin. Do not use these. Give the patient 5-grain doses of quinine every two, three, or four hours until the temperature gives way. We do not find it necessary to put the patient in cold baths, which is a severe and exhausting process still carried out largely in Germany and America. That is unnecessary, and our results are quite as good as those obtained by physicians who employ cold baths as antipyretic agents.

Next, as to internal remedies. The common line of practice is to prescribe saline medicines, such as potassium citrate and ammonium acetate. These constitute an excellent fever mixture, and appear to promote the action of the kidneys and the skin. Such medicines given every four or six hours are certainly helpful. The late Dr. Andrew of this hospital was of opinion that quinine was perhaps the most useful drug to use in pneumonia, and his practice was to give 2 or 3 gr. of quinine every six hours. It is an admirable drug, and I have thus used it for many years. You may add after a day or two some compound tincture of bark to the saline medicine— $\frac{1}{2}$ to 1 drachm to each dose.

Then you must examine the lungs from day to day without exhausting the patient, to see what progress the disease is making. Very often it is confined to one lung. Sometimes it is the upper lobe that is attacked, and this is called apex pneumonia, and has often excited more attention and anxiety than basal pneumonia, the reason being that it is sometimes followed by tuberculosis. Hence apical pneumonia would naturally excite more anxiety if it occurred in a member of a consumptive family. There is another point of interest in apex pneumonia; for some reason, which we do not yet understand, the patient is more apt in this case to be delirious. In regard to such patients the nurse should be warned of the possibility of that happening, especially in the night-time. The pneumonia may involve both lungs, and then you have what is called double pneumonia. The

involvement of both lungs is more serious than when only one is affected, the danger of the disease being in relation to the amount of lung involved. Sometimes the pneumonia spreads very rapidly, and sometimes it wanders about from one part of the lung to another, instead of going on continuously and regularly, and this is called migrating pneumonia.

The natural history of the disease shows that the inflammatory process comes to an end about the sixth or seventh day. You should therefore study the course of the temperature every day in a case of pneumonia. About the fifth day the temperature usually falls to some extent, and then the inexperienced observer may hope that the disease is coming to an end, but this hope is often dashed by the fact that next day the temperature goes up as high as ever. Perhaps on the seventh day the temperature comes down more decidedly than before, and remains normal or perhaps subnormal. The first fall is called the precrisis or false crisis, which often precedes the true crisis. Though the patient by this time is greatly relieved he may have other symptoms—profuse sweating and diarrhoea. These are both means of excretion, and ought not to be interfered with because they are probably carrying peccant matters out of the system—a process which Nature in its wisdom understands better than we do.

When you come to examine the patient's lungs after the critical fall of temperature and the departure of the fever, you are astonished to find that there is apparently very little progress being made; in fact, you find all the signs of consolidation in the lung, all the physical signs of dullness on percussion, bronchial breathing, and *redux* crepitation.

Although the patient has no fever and is feeling much better, yet the specific febrile state which has come to an end has left the lung hampered and engaged in absorptive processes in order that it may clear up, but this clearance goes on favourably if the patient is carefully treated. The lung is, however, still in a delicate condition.

During the actual course of pneumonia the urine is of a febrile character, scanty and high coloured, and presenting this other peculiarity—that the chlorides gradually pass out of it as the pneumonia proceeds. You test this fact in a simple way by adding pure nitric acid to the urine with some nitrate of silver. In a healthy person's urine that brings down a thick white, curdy deposit. But in a pneumonic patient, as the disease progresses this cloudiness and thickness diminish until you get at last an entire absence of chlorides. The chlorides have passed out of the urine, and have been retained in the pulmonary exudation. When the exudation breaks up during convalescence we find the chlorides in the urine again, having been excreted by the kidneys. There is nothing else remarkable about the urine excepting that in severe cases you may find a little albumen, constituting what is called symptomatic albuminuria. This does not indicate kidney disease at all; it passes away during convalescence, and you need have no fear of any permanent renal changes resulting from it.

Supposing the temperature does not fall by crisis on the seventh day, but goes on to the eighth or tenth or twelfth day, you wonder why the crisis has not come. In that case you will have to examine the chest carefully day by day to see what is going on, and you may fairly suspect by the eleventh or twelfth day that there is probably some further disorder or complication. Either the patient is in a very low condition and the disease is not resolving properly or else there is some pleural effusion, and it is very likely that in such a case the patient is suffering from an empyema. Empyema is generally preceded by pneumonia.

If after the seventh day the dullness and the bronchial breathing remain and the temperature keeps up, this indicates that the pneumonia is not resolving and that something else has supervened, and if you put a needle into any dull area of the chest you will probably find pus there due to an empyema. So that if the temperature keeps up to the end of the second week you had better reconsider your diagnosis, and see whether pneumonia really represents all the morbid process which is present in the case.

Sometimes you find pneumonia to be abortive—that is, it does not last so long as the seventh day. The patient is very ill, and perhaps there are herpes and some slight indications of lung disease, some slight dullness, and a little rise of

temperature; but the crisis may come sharply on the second or fourth day and all the symptoms pass off. That is a mild abortive type met with sometimes in children, sometimes in adults.

I mentioned two symptoms in connexion with the nervous system. First of all as to patients who become delirious at night through exhaustion. That, of course, is more apt to be seen in elderly patients and those who have been intemperate in alcohol. Pneumonia in the drunkard is almost certainly a deadly disease. Double pneumonia in the drunkard is absolutely fatal; there is no chance for him. Hughes Bennett taught that acute pneumonia in a previously healthy young man or young woman is usually recovered from; but the case is very different if you have an aged patient or one degenerated by abuse of alcohol or any other chronic disease. It should be noted that the average age of the patients in Bennett's favourable statistics was 30½ years.

Influenzal pneumonia is a very severe form, especially fatal in elderly people. It was so in the last epidemics of influenza. A more important nervous symptom is that of insomnia. The patient cannot sleep, and that, towards the crisis, about the sixth or seventh day, is a very grave matter. It is most important to secure sleep, as the nervous system is exhausted and the patient is worn out with the laboured breathing and high fever.

This opens up the debated question as to the use of opium in pneumonia. There are some authorities who maintain that the use of opium in pneumonia is very beneficial. Others aver that the effect of opium is to stop the excretion from the bronchial tubes, to check the power of coughing and the expectoration, and to stupefy the patient to such an extent that he does not feel the need of coughing; a further risk arising from the condition of the kidneys, which may be unsound, and unable, therefore, to do an adequate amount of excretory work. It does not do to be too dogmatic, and say that in all cases of pneumonia the use of opium is good or bad. There are some cases in which it is beneficial—those cases in which there is no reason to suppose that the kidneys are involved. A very good plan is to give morphine in small doses with compound spirits of ether; 15 minims of liquor morphinae, with 1 drachm of compound spirits of ether (Hoffmann's anodyne) is an excellent soporific, and will make the majority of these patients sleep. If that is not sufficient, the dose should be repeated. The morphine and ether sustain the action of the heart, and also promote sleep.

Another drug which is used is musk, about which I will say more presently.

The next serious thing in pneumonia that we have to fear, more especially after the age of 40, is failure of heart power. In the majority of patients the heart is unable to stand the strain put upon it by the disease. Where does the strain come from? You can imagine the breathing space in the lung being so diminished that there is comparatively little left. The exudation which fills the air sacs presses upon the small vessels, and so prevents the circulation going on freely, and therefore the blood is thrown back upon the right side of the heart and causes engorgement there. Thence it is thrown back into the venous system and this becomes engorged. The result is that the patient becomes cyanosed, and sometimes even jaundiced. This is a very dangerous state of matters; not many hearts can bear that strain for long, and if the organ is originally weak, owing to age or degeneration, the strain will prove too much for it and the patient will die. This is a condition which may be relieved by blood-letting; a small bleeding to 8 or 12 oz. in such a case may be very useful. Dry pleurisy is a constant accompaniment or part of pneumonia, and you may give relief to the pain of it by the application of two or three leeches.

If the heart begins to falter and the pulse becomes irregular and small—a condition which you may expect in elderly people—you may avert further failure by several measures. First of all you give oxygen. Do not put that off until too late; it is often brought in as the last resource. You had better employ it too soon than too late. If the patient inhales oxygen at intervals for ten minutes at a time that may help him on materially. Again, if the heart begins to fail, you may give great relief by injecting 4 or 8 minims of liquor strychninae under the skin. This has a marked power of

stirring up the cardiac centres in the bulb, and by this means many people have been saved from death. I have not found digitalis as useful as strychnine. Now is the time also to use musk. There is only one objection to it, its costliness. We do not use more than a grain or two at a time, but we ought to use more. The old physicians used 9 gr. to 12 gr. of musk made up in the form of a mixture with syrup and mucilage, and I cannot doubt that they got better results than we do with the small doses. When musk is really wanted it is well worth the money. I once had evidence of the value of musk in pneumonia from a Sister in John Ward, a most observant and careful nurse. She used to say to the house-physician, "I think the time has come for musk, sir." She had noticed the good effects of it. We generally combine Hoffmann's anodyne with tincture of musk, and repeat the doses at short intervals.

Of the secondary pneumonias I shall not say much. The pneumonia which accompanies rheumatic fever I do not think is as severe or as frequent nowadays as it used to be. Treatment by salicylates may perhaps account for this. In this form we find everything classical, excepting that the patient does not spit up rusty sputa.

Pneumonia may be dependent on gout, and in other conditions may be quite unconnected with the pneumococcus or staphylococcus infections. It is quite certain that pneumonia may occur as a form of visceral gout and replace articular manifestations of the latter.

Note once more that the treatment of pneumonia is strictly the treatment of the patient suffering from it. It is a simple disease in children, it is not a dangerous disease in the young or in those who have been previously healthy; it is more dangerous to those who have reached forty years of age; it is extremely dangerous in the old, and absolutely fatal in the drunkard.

You might say that the conduct of a case is very simple and demands little more than the practice of common sense. A great deal may be done for this condition by trying to anticipate what may occur. With regard to the use of stimulants there is no routine practice. You should give them in pneumonia just as in other diseases; if they are not wanted you do not give them. Many cases of pneumonia in the young are best treated without any stimulants at all. But if there be engorgement of the side of the heart, a high temperature, a rapid pulse, the first sound of the heart barely audible at the aortic area (Stokes's indication), stimulants should be given. In the majority of cases 2 oz. to 4 oz. of brandy are sufficient in the course of each day. I have known patients to require even 12 oz. of brandy, and they could not have done without it. Your hands are not tied, therefore, in the matter of administering stimulants. It is sometimes more important to give the patient as much pure water to drink as he desires. This simple matter is too often omitted from the treatment of all febrile conditions.

We may begin our treatment with a mercurial purge, and probably we shall not require any other aperient throughout the case. The subsequent treatment consists of a simple saline, to which some cinchona may be added later on; diet of milk and beef-tea, and, above all, watchful and careful nursing. In winter time successions of hot poultices may be applied to the affected side. In hot weather these may be replaced by cotton wool sprinkled with spirit of camphor. I have no liking for a steam-laden atmosphere in cases of pneumonia if the sick room is properly ventilated. Study the temperature chart, the condition of the heart, and anticipate failure of heart power, more particularly in cases where the venous circulation is engorged.

Pneumonia is to be met with in most parts of the world. Wherever you go, therefore, it is important that you should be prepared to act with skill, and with such hope of success as your knowledge of the disease will afford.

VACCINATION OF SCHOOL CHILDREN.—The majority of the cities and towns of Connecticut have adopted a regulation requiring the superintendent of public or private schools to exclude from attendance any child who has not been vaccinated, unless evidence is produced from a physician that it would not be prudent on account of sickness for such child to be vaccinated. The regulation has been approved by the State Board of Health.