remained, but, notwithstanding this, the final result was good.

From the observations quoted we gather that even after medico-mechanical treatment, as well as after suture of the bones, the final result was, with few exceptions, very gratifying. The one unsatisfactory result of non-operative treatment was traceable to a wrong reading of indications, since, as I have already stated, according to my present principles, suture of the patella should have been performed.

Of the 17 cases of primary suture of the patella of which the final result was known, only 1 showed a bad result. The reason of this was the cutting through of the suture, with considerable separation of the fragments. But neither can this unsatisfactory result be laid to the charge of the method as such, for by a reputed suture which was neglected a better result would doubtless have been obtained. Moreover, in this case the technique was so far deficient that the suture was carried out with catgut by one of my assistants contrary to our usual principles.

The 8 cases of late suture of the patella also gave satisfactory results with one exception. In the unfavourable case, however, a traumatic neurosis with hysterical paralysis lay at the root of the bad result; another circumstance for which the mode of operation cannot be made answerable.

We see, then, that both methods—the operative and the non-operative—give equally good results with a right interpretation of indications and a correct technique. These methods do no stand in opposition to each other as alternatives, but each supplements the other, and has its own definite, sharply-marked-off indications.

Dr. Schmidt, the assistant physician of my hospital, will give a detailed account of our results, with reproductions of the x-ray photographs, in Bruns's *Beiträgen zur klinischen Chirurgie*.

REFERENCE.

- ¹ Lister, A New Operation for Fracture of the Patella, *BRITISH MEDICAL JOURNAL*, 1877, vol. ii, p. 850.

The Bradshaw Lecture

ON

INFECTIVE ARTHRITIS.

Delivered before the Royal College of Surgeons on December 10th,

BY HOWARD MARSH, F.R.C.S.,

Surgeon to St. Bartholomew's Hospital.

MR. PRESIDENT, LADIES AND GENTLEMEN,—In my first words I must convey to you, Mr. President, my thanks for conferring upon me the honour of giving the Bradshaw Lecture this year. Within the walls of the Royal College of Surgeons of England, where Hunter laid the foundation and considerably advanced the development of the grandest museum of its kind in the world, where his spirit has ever since been present to inspire his followers; and where his work has been continued by Cooper, Lawrence, Owen, Paget, and our present distinguished curator, Professor Stewart; where our library grows year by year larger, and, as we believe, more and more useful for study and reference; and where at our examinations we have in the name of English surgery highly important duties to discharge, to be invited to lecture is an honour which he who is worthy of it cannot but warmly appreciate. Nor certainly for a moment can he forget the responsibilities which the office involves.

LORD LISTER.

But before I pass to the lecture itself, there is a circumstance to which Mr. President I think you would wish me to allude. On December 9th, 1852, just fifty years ago yesterday, a certain candidate passed his examination and became a Fellow of the College. He came from Essex, and his name was Joseph Lister.

What that name has since become, not only in the annals of English surgery, but among the greatest benefactors of the human race we all remember with admiration and with gratitude. The poet Gray alludes in trenchant phrase to those who shut the gates of mercy on mankind. Lord Lister has done more than any living man to throw them open. No matter in

what century the roll of fame is called, the name of Lister can never henceforth be omitted from it.

Lord Lister was never the man to seek recognition or reward, yet honours have flowed in upon him from every direction, and surely his cup became full when a few months ago he stood by and saw his principles turned to account in one of the most dramatic incidents of history, and when by appealing to them, Sir Frederick Treves, with conspicuous skill and conspicuous fortitude, averted the imminent peril which beset our King. The profound joy with which his subjects in every part of the world hailed the King's recovery was, in our profession, intensified by the fact that His Majesty, by constant devotion and earnest labour in the development of his Hospital Fund—which reached the unprecedented sum this year of £100,000—has made himself the greatest and most productive philanthropist of his generation.

HISTORICAL.

This lecture, together with a similar one at the Royal College of Physicians, was founded by the late Mrs. Bradshaw, to honour and perpetuate the memory of her husband, the late Dr. William Wood Bradshaw, M.A., D.C.L. Oxford. Her action was at once affectionate and enlightened. She hoped that successive lecturers would, each in his turn, do something to advance the profession to the pursuit of which her husband had devoted his life. And it remains to each Bradshaw lecturer to give effect, to the best of his capacity and knowledge, to the objects which she had in view.

The subject I have chosen is that of Infective Arthritis; in other words, I propose to discuss those diseases of the joints which are due to infection. I have not taken this subject as one that is altogether new, for it has long been known that in such instances as septicaemia, pyaemia—and other allied conditions—the joints are liable to become infected. I have chosen it because recent advances in bacteriology and exact clinical observation have made it clear that joint affections of this origin are much more frequent than was formerly supposed. Thus while no doubt much will still be added to our knowledge, the time, I think, has come when it may be useful briefly to review what has thus far been done.

In general pathology one of the main advances recently made has consisted in the discovery of the large part which infection by various micro-organisms plays in the production of disease. Thus, for example, peritonitis, only a few years ago, was supposed to arise as an independent form of inflammation, to be, as the phrase was, idiopathic. It is now well established that no such form is ever met with, but that inflammation is, in the large proportion of cases, infective and due to micro-organisms, of which several forms have been recognized. And as it is with the peritoneum, so it is with the joints. When inflammation occurs in a joint, while it is never idiopathic, it is often infective, and is produced, moreover, by agencies which, when they were first discovered, were supposed to be limited in their action to other structures. Indeed, there exists a close parallel between the peritoneum and the joints in regard to infection; and that this should be the case appears even at first sight probable when the similarity of structure between the peritoneum and the synovial membranes is borne in mind. Both consist of an epithelial layer and a substratum rich in blood vessels along which infective agents easily pass, and through the walls of which their migration can readily take place. A notable illustration of joint infection is met with in the case of the pneumococcus, or the bacillus lanceolatus. This micro-organism, originally discovered in the saliva and shown to be a pathogenic agent by Sternberg, was subsequently proved by Fraenkel to be the active agent in the production of acute lobar pneumonia, and, in the absence of any suspicion that it invaded other organs, it was termed the pneumococcus or lung coccus. Later investigations have shown, however, that it has a far wider range, and that it not only produces pleurisy, peritonitis, pericarditis, and meningitis, but also acute primary arthritis.

In fact, the general statement may now be made that in, at all events, the great majority of specific diseases, the joints are liable to infection. In compiling the list of these diseases tuberculosis and syphilis need not now be considered, and the common forms of septicaemia and gonococcal infection are so well known that they need not be further alluded to. There will then remain the following: pneumococcal infection,

typhoid fever, influenza, scarlet fever, dysentery, erysipelas, and glanders. Lastly, there is the doubtful case of acute rheumatism.

PNEUMOCOCCIC ARTHRITIS.

The discovery in 1888, by Weichselbaum, that the pneumococcus (*diplococcus lanceolatus* of Fraenkel) has the power of producing acute inflammation of the joints, was of the highest importance, not only on its own account, but because of the impetus it gave to the study of the general subject of infective arthritis. In January, 1901, Dr. E. Cave published his paper on pneumococic arthritis, and furnished a table of the cases which had so far been recorded. The list included 31 examples. Dr. Cave's paper was so complete that little of importance has since been added to his description of this affection.

In all these cases arthritis was developed from two to fifteen days after the onset of acute pneumonia. In 2 instances, however, referred to by Cave, arthritis is said to have preceded the pneumonia, in one by three, and in the other by seven days. In 3 cases arthritis occurred independently of pneumonia. Of the 31 cases, no fewer than 25 are known to have been males. The disease in different cases attacked all the large joints in turn—the hip, however, only in 1 instance. The age varied from 5 to 71 years; but only 2 patients were children. The great majority were between 30 and 60, while 6 were over 60 and 2 between 70 and 80. In some cases only one joint was attacked, in others several were involved. The disease was somewhat more frequent in the upper than in the lower extremity.

As to the form of inflammation, suppuration occurred in as many as 27 instances. The extremely grave nature of this form of arthritis is conclusively indicated by the fact that of the 31 cases in Cave's table, no fewer than 23—that is, nearly 75 per cent.—terminated fatally.

The high mortality which attended these cases has been observed in other cases also. It is due to the fact that the pneumococcus produces a general systemic infection—in other words, an acute and profound septicaemia of which the arthritis is merely a local manifestation, others being—in addition to acute pneumonia—malignant endocarditis, pericarditis, pleurisy and empyema, meningitis, peritonitis, nephritis, and otitis media. In a case (Leroux's) mentioned by Cave, the necropsy disclosed endocarditis, a purulent meningitis, pleurisy, peritonitis, and arthritis—a group of lesions which shows that, amongst agents giving rise to general infection, the pneumococcus occupies a bad pre-eminence, while its morbid anatomy may be summarized in the statement by Cave that it is essentially the same as that of any other septic infection.

The pathology of the pneumococcus in respect to the human subject and to animals has been set forth by Foulerton in an able and exhaustive paper printed in the *Lancet*, August, 1901.

ARTHRITIS IN TYPHOID FEVER.

Since attention was first drawn to this subject by Parise in 1842 several writers have alluded to it, and recorded their observations. Among these may especially be mentioned Keen, of Philadelphia, who in 1898 published his important monograph on the complications and sequelae of typhoid fever—including a chapter on affections of the joints—and Barjon and Lesieur, whose essay on *Septicémie Eberthienne* was published in 1901.

Keen, placing himself in line with previous writers, states that there are three forms of joint disease met with in association with typhoid fever:

1. Rheumatic typhoid arthritis.
2. Typhoid arthritis proper.
3. A form of septic arthritis.

(1) Under the head of rheumatic typhoid arthritis are placed cases in which at the commencement of the illness—sometimes as the earliest symptom—the patient complains of severe pain in the knees, elbows, or other joints, which are the seat of varying degrees of swelling. Usually the affection lasts for a few days and then disappears without leaving any ill-effects. In other instances effusion increases in quantity and suppuration occurs. As a good example of this early form Dr. Percy Kidd has given me notes of the following case.

A girl, aged 17, was admitted into the London Hospital, December 9th, 1899, having complained for two or three days

of pain in her left knee, which was swollen. Her case was at first regarded as one of acute rheumatic arthritis. The knee, however, became worse, and on the 14th was explored, as suppuration was suspected. No pus was found. Two days later the temperature was 105°, and diarrhoea, faint rose spots, and Vidal's reaction disclosed typhoid fever. Death occurred sixteen days later from intestinal haemorrhage, and typhoid ulcers were found in the intestines. The joint affection had subsided. Unfortunately no bacteriological examination was made.

To speak of these cases—not that Dr. Kidd does so—as rheumatic typhoid arthritis because they in some respects resemble acute rheumatism is but another example of the confusion which results when terms, which themselves are (in the absence of precise knowledge) virtually meaningless, are so used as to suggest pathological theories. The real nature of these cases is at present undetermined. But it seems likely that the joint condition will prove, on further investigation, to be the direct result of some irritative product of the typhoid bacillus.

(2) *Typhoid Arthritis Proper.*—This is met with in the acute stage of typhoid fever, or towards its decline. Occasionally it makes its appearance during convalescence. Its features vary in different cases. Several joints may be involved (polyarticular form) or it may be limited to one (monarticular). In the former two, three, or more of the large joints may be attacked. There is pain together with more or less swelling, due in part to infiltration and thickening of the synovial membrane, and in part to the effusion of turbid fluid into the joint cavity. In some instances the attack subsides and recovery follows. In others, the joints, after prolonged inflammation, undergo fibrous ankylosis, which can be removed, if at all, only by repeated manipulation under an anaesthetic and prolonged passive movement and massage. Or, again, suppuration may ensue, necessitating free incision and irrigation.

In the monarticular variety it is the hip which in the great majority of cases is involved, and the result is usually serious, for some cases end in firm fibrous ankylosis much resembling that which ensues after gonococcal infection, while in others (and much more frequently) the capsule become distended by serous effusion, and then either spontaneously, or when the patient is lifted, dorsal dislocation occurs.

The pathology of this group of cases is as yet but imperfectly worked out. Such evidence, however, as is at present available tends to indicate that the arthritis is due to the local action of the typhoid bacillus on the tissues concerned. It is well-known that, although the typhoid bacillus is not usually present in the general circulation, it is to be found in the liver and the spleen, in bone marrow, and in the rose spots on the skin; in other words, that it is widely distributed in the tissues. It is therefore easy to see that along with other parts the joints may be invaded by it. Positive evidence, however, of the presence of the bacillus in the joints is as yet very limited. Several observers who have examined fluid withdrawn from joints in these cases have found that it was completely sterile. Too much weight must not be attached to this negative result.

Possibly the bacillus, although originally present, subsequently undergoes disintegration. It is not rare to find collections of pus in various parts which are sterile but which almost certainly were produced by micro-organisms now no longer to be found, and Cave remarks that even in some cases of suppurative arthritis occurring in association with pneumonia the pus removed from the joint has been sterile. But in such cases it is nearly certain that the pneumococcus was originally present. Or the explanation suggested by Vidal that the arthritis was produced not by the pneumococcus itself but by toxins derived from it may be correct, a view which may also be held in the case of the arthritis of typhoid.

On the other hand, Delanglade and Chibret found the typhoid bacillus in the joint fluid, while Orloff produced arthritis by injecting the bacillus into the joint of a rabbit. Further observations are required to determine the real origin of the form of arthritis under discussion, but at present such evidence as we possess points, as I have said, to the conclusion that the typhoid bacillus may produce arthritis by its specific action on the tissues, just as may the gonococcus or the pneumococcus.

(3) *Septic Arthritis.*—It will readily be understood when the

condition of exhaustion to which the patient is reduced by typhoid fever is borne in mind, that septic arthritis is a highly dangerous complication. It is, in fact, very generally fatal. It, however, presents no specific characters. It is an instance of the common form of septic infection, and is due to the presence of streptococci or staphylococci which have been absorbed from boils, or bedsores, or from the surface of intestinal ulcers. Perhaps the bacillus coli communis may also in some instances be concerned, but as to this there is, so far as I know, no positive information.

ARTHRITIS IN SCARLET FEVER.

The arthritis which is not very rarely associated with scarlet fever occurs in two forms. The first is often indistinguishable clinically from acute rheumatism. It makes its appearance either quite early—while the rash is still present—or during the desquamative period. It usually involves several joints, is not severe in character, and, in the majority of cases, soon subsides. As to its pathology, Dr. Hilton Fagge remarks, "probably most cases of synovitis following scarlet fever are true rheumatism." Osler regards it as analogous to gonococcal arthritis, while Dr. F. Taylor holds practically the same view, for he says, "although known as scarlatinal rheumatism, it is possibly an arthritis due to the direct action of the septic organism of the primary disease." This, in the present state of our knowledge, and judging by such instances as the pneumococcal, gonococcal, and probably also typhoid arthritis, is likely to be the correct view. The second form is much more severe, and not rarely ends in suppuration. It is usually, as Strümpell remarks, a part of secondary sepsis or a general pyæmia, as evinced by such other lesions as empyema, subcutaneous abscesses, splenic tumour, etc. Here, as in the septicaemic form of typhoid arthritis, the infective agent is the streptococcus pyogenes, or the staphylococcus.

ARTHRITIS AFTER INFLUENZA.

I have met with two cases which appear to suggest that, with whatever rarity, arthritis may follow influenza. On this subject Boncoumie has contributed a paper to the *Bull. Gén. de Thérapeutique*, vol. cxi, p. 5, 1900. His observations, however, refer not to any specific form of arthritis directly dependent on influenza, but to the predisposition which influenza establishes to attacks of ordinary gout. The subject is one about which little is known, but it seems possible that these cases of Boncoumie which resembled gout may have been due directly to the action of the influenza bacillus.

The first of my own cases was that of a gentleman, aged 42, whom I saw with Dr. Hetley, of Norwood. The patient had a sharp attack of influenza (which was at the time epidemic in the neighbourhood) on May 3rd, 1891. Two days later he complained of severe pain, apparently in the glands below Poupert's ligament; he had a rigor, copious sweating, and a temperature, for the next ten or twelve days, ranging between 102° and 103.5°. During this time nothing abnormal could be detected in the groin or thigh, till on May 11th a band of oedema appeared running down the outer side of the limb from the trochanter nearly to the knee, but this was neither painful nor tender. Pain was situated chiefly in Scarpa's triangle on the inner side of the femoral vessels. There was a tender spot also over the sciatic notch. On May 14th the temperature went down and the oedema subsided, but some thickening could be felt in front of the joint, and the artery seemed to be pushed forward. At this time the limb was becoming flexed and abducted. There was marked muscular wasting, and pain about the joint was so severe as to call for the hypodermic injection of morphine. Two consultations were held in the next three weeks, but the nature of the case remained obscure. Dr. Hetley, however, formed the opinion that the hip-joint was involved. I first saw the case on July 5th. The patient was very weak, temperature was 101° to 103°. The limb was flexed and adducted; there were night startings and much muscular wasting. Any attempts to move the limb produced severe pain. It was obvious that the joint was actively inflamed. Weight extension in the axis of deformity was used. The symptoms gradually subsided, and the patient gained flesh and strength and was able in about two months to be up on crutches. When I saw him again in the following December, there were $\frac{3}{4}$ in. of real shortening and

some adduction of the limb; the trochanter was considerably above Nélaton's line. The joint was stiff, but all active symptoms had ceased. At the present time the patient walks with a stick, but lameness is marked. He has only fatigue-pain. No further shortening has taken place. Any one who now examined the limb would probably regard the case as one of monarticular osteo-arthritis. A few days after the hip was attacked the patient complained of pain in the knee, which was enlarged from periarticular infiltration. There was no fluid in the synovial cavity. This condition persisted for four or five weeks but then slowly subsided. The joint ultimately completely recovered.

In the second case, a boy, aged 18, after a moderately severe attack of influenza, complained of stiffness and severe pain about the right hip. On his admission, two months later, into St. Bartholomew's Hospital, the limb was flexed and abducted, the joint was stiff, and there was much brawny oedema of the soft parts in Scarpa's triangle in the iliac fossa, and on the outer side of the joint, and below and a little internal to the iliac spine there was a suspicion of fluctuation. The inguinal glands were enlarged. The muscles were wasted. The position of the limb was corrected by weight extension and swelling slowly subsided, but stiffness remained. The brawny thickening persisted for several months and suppuration often appeared imminent. At present the joint is stiff. There is no pain, thickening around the joint remains to some extent. The patient walks with only slight lameness.

As to the sequelae of influenza I may mention in passing the case of a surgeon who suffered from acute purulent cystitis attended with high temperature, severe pain, and rapid loss of flesh and strength after a severe attack of influenza. The bladder was irrigated two or three times a day with warm boracic solution, and in about a fortnight the cystitis subsided and he ultimately made a complete recovery.

ERYSIPELAS AND ARTHRITIS.

Arthritis as a complication of erysipelas appears to have been first recorded by William Musgrave in 1702, and Boissier de Sauvage mentioned it in 1740. But the arthritis in question was believed to be acute rheumatism and was by some later writers termed erysipelatous pseudo-rheumatism.

An important contribution to this subject consists of a thesis written by Dr. René Jorrot, published in 1899. Dr. Jorrot has collected from various sources a number of cases which show that the joint affection usually occurs in the acute stage of erysipelas, though in a few instances it comes on when the disease is declining. Like the arthritis met with in septicaemia it may occur as—

(a) Transitory synovitis attended with serous effusion and involving many joints and sometimes the sheaths of tendons;
(b) As a plastic form tending towards fibrous ankylosis. This form may be persistent over many weeks; in one case it lasted for two months and a half;

(c) An acute suppurative arthritis by which the joints are rapidly disorganized.

No exhaustive bacteriological examination seems to have taken place as to the agent by which the arthritis is produced. In the early days a chain coccus was described by Fehleisen as the streptococcus erysipelatis, but this is now very generally regarded as identical with the streptococcus pyogenes. Suppurative arthritis appears to be rare as a complication of erysipelas. Jorrot calls it very rare—1 in about 200 cases.

ARTHRITIS IN GLANDERS.

Delafield and Prudden allude to arthritis as met with in the course of glanders, but they give no detailed account of the affection, nor have I found elsewhere any full description of it. It would probably be acute and destructive.

As to acute rheumatism, although it is believed by many authorities, from the evidence already obtained, that it is an infective disease, there is up to the present no agreement as to the agent by which it is produced; some believe it to be the diplococcus first described by Triloubet, and since carefully studied, among others, by Paine and Poynton; Singer, but with few adherents, holds that it is the streptococcus or the staphylococcus; others consider that the disease is caused not by any one micro-organism exclusively, but by a mixed

infection, arising from bacteria lodged in the tonsils or the adenoid tissue of the pharynx.

I have now passed in brief review the principal instances in which the joints are liable to infection in the course of the various specific diseases, and it will be apparent that the infective agent is different in different cases. In some cases it is the same micro-organism (or its toxins) as that which produces the primary disease—as, for example, when acute arthritis occurs as a complication of pneumonia, and the pneumococcus is found in the joints; or when arthritis, developed in the course, or as a sequel, of typhoid fever, depends on the local action of the typhoid bacillus.

The arthritis found in association with scarlet fever appears to be a mild form of ordinary septicaemia due to the streptococcus pyogenes or the staphylococcus, while that which is sometimes met with in dysentery is the result of a mixed infection, in which, perhaps, the bacillus coli communis plays a chief part.

In studying these infective agents, it is to be remembered that the effects which they produce may be largely influenced by the varying conditions under which they are placed, as determined, for example, by the resisting power of the individual attacked, the dose received, and the presence of other micro-organisms or their toxins. These are matters about which at present little is known. Little, for instance, I think, is known as to the influence which the typhoid bacillus or its toxins may have in modifying the action of the streptococcus or the staphylococcus, or vice versa.

INFECTIVE JOINT CHANGES.

The changes met with in joints which are the seat of infective arthritis vary considerably in different instances. In some they are slight and transient, and result from synovitis, attended with infiltration of the sub-synovial tissue and with some, but only a limited amount, of serous effusion into the cavity of the joint. This is the case, for example, in the arthritis which occurs in the early stage of scarlet and typhoid fever.

In a second group, one, two, or more of the joints are painful and swollen from effusion of fluid into the synovial cavity. These are the cases which have been so misleading in consequence of the close similarity of the appearances observed, on clinical examination, to those of acute rheumatism. Although at first, as I have said, merely turbid, the fluid in the joint in some cases soon becomes purulent.

In the third group the inflammatory process chiefly involves the periarticular tissues, and leads to considerable brawny or boggy swelling, and to reddening of the skin, which is at the same time so stretched and shiny as to suggest the presence or near approach of suppuration. This is one of the most clearly-marked types of infective arthritis, and one with which we have long been familiar in some of the cases of gonococcal infection. In this form there is frequently no effusion into the cavity of the joint, and, so far as I am aware, though it seems imminent, suppuration very rarely occurs. Clinically, these cases run a very prolonged and tedious course; they may extend over several weeks, some have lasted two or three months. Many are attended with pain, not only persistent but very severe, and they show a strong tendency to result in firm fibrous ankylosis, which apparently, in some instances, gradually passes on, by ossification of the new fibrous tissue, to complete synostosis. In these respects they furnish a close parallel to one of the forms of arthritis which is met with in the course of the course of the septicaemia which follows acute infective periostitis or osteomyelitis.

A fourth group of cases includes those in which the arthritis is, from the first, acute and destructive. Suppuration takes place early, and goes rapidly on to complete disorganization of the joint, and often to the wide burrowing of pus, which has escaped from the synovial cavity, along the intermuscular spaces of the limb. This fourth variety is met with in its most marked phase in pneumococcus arthritis.

UNCLASSIFIED INFECTIVE ARTHRITIS.

Thus far I have been laying before you such an account as time will permit of infective arthritis as it occurs in connexion with various specific diseases. But apart from this clearly defined group, and under widely varying circumstances, other examples are met with, and to these I will now ask your attention.

Every one who has many patients passing under his notice meets with cases of joint affection attended with fever as to the nature of which he is uncertain, but which are traditionally regarded as acute or subacute rheumatism. Now, just as charity covers a multitude of sins, so has the term rheumatism covered a broad expanse of loose pathology. Undefined itself, it has thrown its aegis over many conditions equally obscure. To-day, however, when it becomes more and more probable that acute rheumatism is itself an infective disease, it is seen that many cases which have been termed rheumatism may also belong to the infective class.

Certainly one of the most instructive contributions towards the elucidation of this obscure subject is the paper in the current volume of the Clinical Society's *Transactions*, in which Drs. Paine and Poynton relate the result of their investigations into the pathogenesis of two cases of arthritis under the care of Mr. Page in St. Mary's Hospital. In Mr. Page's first case, a man, aged 34, fell and sprained his right knee. About three weeks later, on his admission (January 11th, 1901) to the hospital, the joint was greatly swollen and very painful. As swelling had not diminished, aspiration was performed on January 15th. The pain, however, continued, and the swelling began immediately to return. The patient was restless and irritable, and as it were "off his head." On January 18th fluid was again removed and the joint washed out with carbolic lotion (1 in 100) till the fluid returned perfectly clear. After this steadily progressing recovery took place.

In the second case a boy of 14 was admitted March 22nd, 1901, having hurt his knee shortly before Christmas. Some swelling followed, but passed off in a few days. On March 15th he knocked his knee against the wall as he lay in bed. The blow was followed immediately by swelling and considerable pain. On admission the joint was tightly distended with effusion, and he was evidently very ill, his temperature was 100° to 101° F., and he was restless and irritable, and inclined to wander. March 26th. Fluid removed from the joint. The boy was, however, no better, but rather worse, and his temperature was higher. On April 2nd the joint was again emptied and washed through with carbolic lotion. From this time he steadily recovered and left the hospital on May 9th.

I can only briefly summarize the bacillary and experimental investigations of Drs. Paine and Poynton, than which it would be difficult to find work more admirably done or more suggestive.

On cultivation the fluid contained numerous very minute diplococci in chains. Twenty minims were injected intravenously into a rabbit. Arthritis followed in several joints. The heart's action became very excited, and a systolic murmur developed at the apex. Two months later the rabbit was killed. The necropsy showed mitral valvulitis. The joints contained exudation like raw white of egg, and showed swelling due to chronic inflammation in the periarticular tissues. Two other rabbits were treated by intravenous injection, and again a polyarthritis was produced. In none did suppuration occur. "The investigation demonstrated," the authors of the paper observe, "that the exciting cause of the arthritis in the second of the two cases described by Mr. Page was a diplococcus, and that this diplococcus resembled in its morphological and cultural characters the diplococcus of rheumatic fever rather than the streptococcus pyogenes."

On a subsequent occasion Drs. Paine and Poynton at a necropsy found in the exudation and synovial membrane of a knee-joint showing all the changes of osteo-arthritis a diplococcus which in ten weeks produced an osteo-arthritis of the right knee-joint of a rabbit. This observation is of great importance, and we may hope it will not be long before we hear more from the authors respecting this subject. The severe form of osteo-arthritis occasionally met with after parturition seems especially suitable for bacteriological investigation.

In October, 1901, at the Clinical Society, Dr. Percy Kidd showed a female patient, aged 26, who eighteen months before had become the subject of chronic bronchiectasis. Six months later the sputum was offensive and more copious, and the ankle-joints became affected. Subsequently the wrists and knees were involved. When the patient was exhibited both wrists were swollen, stiff, and tender. The fingers were generally enlarged, the joints swollen and tender, and the ends clubbed. The knees were swollen, and each contained a little

fluid. The ankle-joints were swollen but not tender. A remarkable point in this case was that the condition of the joints varied with the variation in the bronchiectasis, undergoing considerable improvement when the sputum was limited in amount and less fetid in character, but becoming worse when the sputum was more copious and more fetid. In other words, the amount of arthritis seemed clearly to depend on the amount of absorption that was taking place.

In a second case recorded by Dr. Kidd, in a man aged 30, who was suffering from a foul and copious discharge due to bronchiectasis, the knees, wrists and ankles were swollen and contained fluid. This patient stated that he had had gout on several occasions during the last six years, but the present joint affection began three years ago with pain and stiffness in the knees and ankles. He had chronic cough with expectoration for three years and for a year the sputum had had a foul taste.

The fact that this patient had suffered from gout, so that his joints were already damaged, may be noted in reference to the case recorded by Barjon and Lesieur. A girl, aged 17, having had typhoid fever when she was 8, and acute rheumatism when she was 11, was attacked on October 1st, 1900, with acute illness, accompanied by raised temperature, a cardiac *bruit* and painful swellings of several of the joints. The joint symptoms completely disappeared by October 15th. But the general illness continued and now presented the phenomena of typhoid fever: there were rose spots, profuse diarrhoea, high temperature, bronchitis, and delirium. Death occurred on October 28th. At the necropsy no intestinal lesions were found. But that the disease had been typhoid fever was proved by the discovery in the blood (removed from the patient on October 12th) of the typhoid bacillus. The view the authors held in respect to this case was that while the glandular structures of the intestine had been rendered immune by the first attack of typhoid and so escaped, the joints, already damaged by acute rheumatism, had lost their normal resistance, and thus became localities in which the irritative action of the typhoid bacillus, or its toxins could take effect.

Thus both in Dr. Kidd's case and in that just described the joints were, it appears, predisposed to attack in consequence of their diminished powers of resistance, produced by previous gout in one and acute rheumatism in the other. And bearing on this same loss of resistance, we notice that in Dr. Cave's table in no fewer than 10 out of 31 cases pneumococcal arthritis was developed in recently-injured joints; while in Vero's case pneumococcal infection declared itself a few hours after a young lady had been exposed to cold by passing through the house during a night in October in her nightdress. Within forty-eight hours the right knee-joint, which was swollen and painful, was punctured, and yellow pus was withdrawn.

The influence of injury, of whatever kind, in predisposing joints to attack by the pneumococcus and other micro-organisms should be carefully kept in view. Unless this is done, cases really depending on infection may be regarded as merely traumatic, and so the requisite treatment by evacuation and irrigation of the affected joint may be withheld until very serious mischief has taken place.

Dr. W. H. Brook recently described at the Clinical Society an epidemic which occurred in May, 1902, in Lincoln, and which was traced to infected milk. A large number of people were suddenly attacked with very severe sore throat and oedema of the fauces and uvula. The tonsils were coated in many cases with a drab-coloured fur. The cervical glands were enlarged and tender, and there was in many instances a skin rash resembling urticaria. One patient died of pyaemia. High temperature was reduced in several cases by the administration of antistreptococcus serum.

One of these patients, a man, aged 37, had, on about the tenth day of illness, intense pains in the knees, ankles, and elbows. The slightest movement produced agony. None of the joints contained fluid. There was no endocarditis. For several weeks the patient was unable to move any of his limbs. Now he can walk, but with great difficulty.

Similar joint symptoms were present in several other cases.

On the subject of infective arthritis, Mr. Clement Lucas's observations (1885) on the occurrence of infection of the joints as a complication of ophthalmia neonatorum formed a very important contribution. His views are fully recorded in his paper in the *Med. Chir. Trans.*, 1899, while other observers

have amply confirmed him. Among the most instructive papers is that by Dr. Hawthorne in the *BRITISH MEDICAL JOURNAL*, May, 1902.

In Dr. Hawthorne's case speedy and complete recovery resulted when the conjunctival discharge was removed by treatment, that is, when the supply of infective material was stopped, a result corresponding with the variation in the degree of arthritis mentioned in Dr. Kidd's case dependent on the condition of the bronchiectasis, and as to the amount and character of the discharge.

THE ACUTE ARTHRITIS OF INFANTS.

In 1874 Sir Thomas Smith described, under the title of *Acute Arthritis of Infants*, a series of cases in which young children were attacked with an acute form of arthritis attended with early suppuration and disorganization of the joints, and often terminating fatally. All the patients were under a year old, and in several the affection was developed in the first week or two of life. In some instances as many as four of the large joints were involved. In one infant swelling of the right knee was noticed very soon after birth, and in the next month the opposite knee, the left elbow, and the left ankle were attacked. In one case a female infant, 4 weeks old, along with purulent polyarthritis had empyema. In these cases on necropsy it was almost invariably found that the arthritis was secondary to an abscess which had developed in the end of one of the bones forming a joint, and had made its way through a pinhole orifice in the cartilage into the joint. When his paper was written Sir Thomas Smith was not in a position fully to explain the pathology of these abscesses, although the necropsy indicated in some of the instances that the cases were associated with pyaemia. But no one would, I think, at present hesitate to believe that they were infective in their origin. Probably the infective agent concerned was either the streptococcus pyogenes or the staphylococcus. Precisely where the primary infective process was situated is uncertain. In some instances it perhaps was connected with suppuration around the funis, in others with septic conditions in the nose or pharynx, whence, or from some other locality, the infection was carried in the blood stream into the vascular cancellous tissue in the end of one of the bones. My object in mentioning these cases is to add them, as I am sure I may do, to the list of cases of infective arthritis.

The following cases have been seen in the past few years, either in consultation or in St. Bartholomew's Hospital:

CASE I.

A man, aged 54, who had been a free liver, and had diabetes, had a large boil on the abdomen, near the groin. This was opened and 2 drachms of offensive pus were let out. Within two or three days he had very acute inflammation of the right knee-joint, which became flexed, considerably swollen, and globular from periarticular infiltration (there was no fluid in the joint) and intensely painful. The skin was red and shiny, suggesting early suppuration, but none occurred. The patient had a high temperature and was very ill. On seeing him, with Mr. John Adams, of Aldersgate Street, I suggested boric fomentations and weight extension. Under treatment, very carefully carried out by Mr. Adams's partner, Dr. Dyson, slow improvement occurred, and in six months the patient was able to be upon crutches; fibrous ankylosis, however, occurred. The patient is now well, and the joint is free from pain. The knee is slightly flexed, and quite stiff.

CASE II.

A case almost exactly similar, except that the source of infection was untraced, was that of a lady, aged 37, a patient of Dr. Haynes, of South Kensington. This lady "caught a sharp chill" at an evening entertainment and cold weather. Next day she had severe pain in the right knee, which became red on the surface and considerably swollen. Pain was so severe and prolonged that it was necessary to keep the joint on a splint for two months. For several weeks the temperature was raised. Pain and swelling subsided slowly, but the joint was left firmly ankylosed. At the present time there is no pain, but the knee is stiff and still enlarged by brawny periarticular thickening.

CASE III.

An unmarried lady, of 23, had acute arthritis of the right knee following a wrench at tennis. The joint a week after

the injury was considerably swollen, very painful, and covered with dusky red skin. The joint contained no fluid, but the periarticular tissues were thickened and brawny. The limb was kept at rest on a back splint and swung from a cradle. The patient's temperature was 99° in the morning and 101° in the evening, and for the next month pain continued to be severe, and it seemed probable that suppuration was about to occur. From this time, however, very slow improvement took place, and at the end of three months the splint was removed. The limb was in good position, but the joint had undergone close fibrous ankylosis, and the patella was firmly fixed on the condyles of the femur. The patient six months later was walking on the limb, but the joint was so completely rigid that I believe the ankylosis, which was at first fibrous, had now become bony.

CASE IV.

Mr. X., aged 24, had an illness extending over about six weeks, and attended with rise of temperature. Several of the large joints and the finger-joints were the seat of pain and swelling, but the left ankle was especially involved. It was extremely painful and considerably swollen. All the other joints recovered, but the ankle remained inflamed and painful for two months, and ultimately became firmly ankylosed. I saw Mr. X. in consultation with Sir A. Garrod, and we agreed in regarding the condition as gonococcal in its origin; but this the patient firmly denied. As the foot was in the condition of slight equinus, I manipulated it under ether and brought it up within a right angle with the leg. During this proceeding, which required some force, strong cicatricial tissue was torn through. The ankle-joint itself remained stiff, but by long-continued manipulation and exercises considerable movement was acquired in the medio-tarsal joint, so that the patient ultimately walked with only slight lameness. It is, of course, impossible to be sure that gonococcal infection did not occur in this instance; but as the patient, who was intelligent and straightforward, persistently denied it, it seems unreasonable to assume its presence, and, looking at the case in the light of recent experience, I now regard it as one of some other infection, although here, as in preceding cases, what the agent was—whether the pneumococcus, the streptococcus, or some other—I cannot say.

CASE V.

A man, aged 29 years, was admitted into St. Bartholomew's Hospital in whom all the large joints of the lower extremities had undergone ankylosis, after an illness attended with fever of thirteen weeks' duration. As to the nature of this illness no trustworthy information could be obtained, but apparently there had been no gonorrhoea. I think there need be no hesitation in pronouncing this case is one of infection; but what the infective agent was must remain uncertain.

CASE VI.

A lady, aged 35, was suddenly attacked with swelling and severe pain in the left ankle, which became uniformly and considerably swollen. Pain in the next three days increased to great severity. The temperature was $101-102^{\circ}$, and the skin was red and shiny, as if suppuration would occur. The joint was carefully supported in a poroplastic splint, and covered with warm boric fomentation. As the patient had had gout on a previous occasion, this attack was at first considered to be of this nature, and colchicum was prescribed. No improvement followed, and sodium salicylate also failed to give relief. Morphine was used hypodermically for seven or eight days to relieve the severity of the pain. The symptoms slowly subsided. Three months later the patient could walk on the foot, but the joint became firmly ankylosed. The source of the infection, which I cannot doubt was present, was quite obscure.

I have felt compelled, even at the risk of being wearisome, to state the clinical evidence which has led me to believe that, scattered about in everyday practice, are a number of cases which have hitherto been regarded as rheumatic, but which are really infective. Many such cases, it is true, when they are taken singly present no distinct resemblance to what is usually regarded as the typical form of infective arthritis. When, however, they are massed together, and their features are compared, and when they are studied in connexion with

observations which have recently been made in bacteriology, their real nature appears to admit of scarcely any reasonable doubt, and I venture to anticipate that in the future their comparatively common occurrence will be generally recognized.

PROGNOSIS AND TREATMENT.

In the first form of infective arthritis which I have mentioned, that, namely, which consists of a transient synovitis, attended with limited effusion, prognosis is favourable. The arthritis soon subsides, often in the course of four or five days, and the joints completely recover. A suitable splint and warm fomentations will be the only treatment required.

In Group II, in which the joint cavities contain fluid, the treatment imperatively called for is clear. It is the same as that which should be employed in gonococcal infection. The fluid must at once be removed and the joint freely irrigated, either with carbolic lotion 1 in 100, or with mercurial solution, of which, perhaps, the best form is biniodide 1 in 1,000. To evacuate the fluid, a full-sized hydrocele trocar and cannula may be used, and irrigation can be readily performed through the cannula: or the joint may be opened by an incision at the side of the patella. If the fluid proves to be already purulent, the joint must be freely opened and the finger inserted to break down any adhesions which may have formed, and behind which pus might be imprisoned, and then thorough and copious irrigation must be carried out. The immediate improvement and complete recovery observed in Mr. Page's cases after evacuation and irrigation was very striking and it has its parallel in what, I think, many surgeons will have observed in cases of pyaemia, namely, that, when a joint has become distended with pus, if it is freely opened and copiously irrigated, it may forthwith undergo a startling improvement and, if the patient survives, may completely recover and retain absolutely free movement.

In the third or plastic form prognosis is distinctly unfavourable. The arthritis tends, as I have said, to extend over a considerable period. A large amount of new fibrous tissue is developed both between the articular surfaces and in the periarticular tissues, and the joint is converted into a massive scar, so that firm fibrous ankylosis, which may subsequently become bony, results. This strong tendency towards ankylosis is, I think, one of the especial characteristics of infective arthritis; nor are any means at present known by which it can be prevented. As to treatment, the best that can be done is to keep the joint for the time being at complete rest; indeed, the pain is such that no alternative can be thought of. Warm boric or opiate fomentations are required during the most acute stage, but when swelling and heat have somewhat subsided a succession of small blisters will alike relieve pain and promote absorption. Later still, massage will be required to remove the brawny oedema of the soft parts. A very important question is, whether, in any of these cases, manipulation should be employed with the object of restoring movement. In the slighter cases this is advisable, but when inflammation is either severe or prolonged, the joint becomes filled up with cicatricial tissue and the restoration of movement is impossible. Forceful manipulation in such conditions is not only useless, but definitely mischievous, for it renews irritation and promotes the formation of scar tissue. Moreover, I have seen instances in which manipulation has left a joint which was previously free from pain so sensitive and painful for many weeks that it was necessary to keep it at complete rest, so that instead of being diminished, stiffness was increased.

In the fourth group prognosis is highly unfavourable, for the arthritis is but one of the manifestations of a general septicaemia, and is often associated with other lesions of the gravest kind. Often the arthritis is rendered comparatively unimportant—except for the suffering it entails—by the speedily fatal termination of the case. In those rare cases, however, in which the septicaemia is of a milder type and other local developments are absent, if the joint is at once freely opened and irrigated, repair may take place sometimes with ankylosis, sometimes with the restoration of considerable, or even completely, free movement.

In the future when our knowledge of pathology has become more exact and when each infective agent can, as we may hope, be opposed by an appropriate antitoxic serum, these cases may be treated with much more success than is at present possible.