permanently closed according to the nature of the obstruction as shown by the subsequent progress of the case.

The two cases appended to this paper, both of which occurred in my practice in the same week, will serve as illustrations of the treatment advocated. In both, the symptoms were immediately relieved by the small opening into the cæcum; in both Nature ultimately restored the natural passage, and the fecal fistulæ were closed by the simplest operation. As an illustration of how little shock is induced by the treatment suggested, I may mention that the second of my cases was a lady over 80 years of age, who completely recovered, and has since remained in excellent health.

CASE I.—In March, 1892, Dr. Jeaffreson, of Wandsworth, asked me to see a lady, aged 45, with the following history: On some four or five occasions during the last ten years she had had attacks of sharp colicky pains, lasting a few hours, and passing away without further symptoms. Seven days before she had had such an attack, but instead of the pain passing off as on former occasions it had remained, coming on in paroxysms. During the last two days she had had a certain amount of retching, but not violent. She was not seen by Dr. Jeaffreson till the sickness commenced. She had several injections of water and oil, but practically without result. On my visit the patient expressed herself as feeling better and in less pain than on the previous day, and there had been no vomiting for the last twenty-four hours. The temperature was normal, and the pulse 100. The abdomen was uniformly and greatly swollen, no tumour could be recognised, nor could anything be felt in the rectum. Owing to the improvement in the symptoms and vomiting having ceased, it was decided that no operation should be yet performed. All food was stopped by the mouth, small nutrient enemata being substituted. During the next two days there was practically no change, she was not sick, but nothing was passed by the rectum. On the third day, however, vomiting recommenced and the belly appeared a little more distended; she was restless, and the pulse 120. After consultation it was decided that operative interference should be no longer deferred.

The abdomen was opened by an incision three inches long on the left side; distended small intestine immediately crowded out. Two fingers introduced could feel nothing but distended intestine. The sigmoid flexure was then sought for and found to be contracted and empty as far as it could be examined. The wound was closed, buried sutures being used. The abdomen was then opened on the opposite side over the cæcum, and this was found enormously distended. The parietal peritoneum was carefully stitched to it with fine sutures in such a way that only a small circle of gut, the size of a sixpence, was left exposed. The cæcum was then punctured by a trocar and cannula, the puncture being made through a piece of india-rubber sheathing to protect the wound. A large quantity of wind and fluid feces escaped through the cannula. This at once caused the abdomen, which had been previously tightly distended, to collapse to a considerable extent, and immediately took off the pressure on the stitches. I then enlarged the opening so as to introduce the little finger. Previous to puncturing the bowel it would have been quite impossible to have raised it to the skin surface, but after its contents were evacuated it came to the skin level without any tension. Ten more sutures were then passed, fixing the small opening firmly to the skin.

The general symptoms were at once relieved, and the patient made an uninterrupted convalescence. Fifteen days after the operation a small quantity of motion was passed for the first time by the rectum. The quantity then increased day by day, so that by the end of a month nearly the whole of the motion passed by the natural passage. The opening into the cæcum had contracted to the size of a No. 8 catheter. Wind occasionally passed through this, and a very small quantity of fecal discharge. In May I closed the fistula. The patient walked and drove about, and remained in good health till the end of July. She then had another attack exactly similar to the first. The abdomen became greatly distended, with much pain and vomiting. Simple measures having failed, and the patient on the fourth day passing obviously into a dangerous condition, I cut through the centre of the small scar, and again opened the cæcum sufficiently to admit the little finger. A copious escape of wind and fluid feces gave rapid relief. The patient convalesced in a few days, and normal motions again passed by the rectum. The patient now has a little valvular opening over the cæcum, the size of a No. 8 catheter. A pad of wool is worn over this, and it remains practically unstained. The patient will not have this closed at present, but keeps it open by a little glass rod, passed daily. She says the opening causes her the least possible inconvenience, and she regards it as a safety-valve against the recurrence of the symptoms which she so greatly dreads.

CASE II.—In March, 1892, I was asked by Dr. Townsend to see a patient, aged 79, in consultation. She had enjoyed good health till seven days previously. She then suffered a good deal of gripping pain. Since that time she had passed no motion, and suffered severely from recurrent paroxysms of pain. During the last thirty-six hours she had had much sickness and retching. The abdomen was greatly distended and tympanitic. No tumour could be discovered either through the abdominal walls or by the rectum. We agreed that if by the following morning the symptoms having increased rather than otherwise, the abdomen was to be opened on the left side in the position already described. Distended small intestine presented itself. On further search the lower part of the sigmoid flexure was found contracted and empty. On following this up for 8 or 9 inches, a hard solid mass was felt. With some little difficulty this was drawn out of the wound, and it was found to consist of a dense hard mass of inspissated feces in the lower part of the descending colon. The mass was about 4 inches long and 2 in diameter. The colon was so stretched over it as to be semitransparent and scarcely thicker than goldbeaters' skin. Having protected the abdominal cavity by sponges, and placed a temporary ligature around the bowel above the mass, a clean incision 2 inches in length was made in the long diameter of the bowel

opposite the mesentery. The mass was then broken in two, and each portion extracted without difficulty. The bowel wound was closed with fifteen fine silk sutures, the temporary ligature removed, the intestine returned into the abdominal cavity, and the wound closed. Thirty-six hours later I was summoned to see the patient again.

Soon after the operation she had passed some wind and a small quantity of motion. By this the distension was somewhat relieved, but it increased again, and vomiting continued. I felt confident that the large intestine so long distended above the obstruction was paralysed and unable to pass its contents through the contracted intestine lying below the obstruction, thus practically the patient was dying unrelieved. Her general condition was such as forbade further delay, or even an anæsthetic, so that at once, with the assistance of Dr. Townsend, I made a small incision into the abdominal cavity, and exposed the cæcum, which was greatly distended. The parietal peritoneum was carefully stitched to this, so as to leave a circular area of the bowel no larger than a three-penny piece exposed. A trocar and cannula were then introduced into the distended gut. A considerable quantity of wind and motion escaped. An india-rubber tube the size of the little finger was then passed into the opening, in which it fitted fairly tightly, and through this the bowel was washed out with a copious supply of hot water. By carefully packing sterilised wool round the tube, the wound itself was kept comparatively free from contact of fecal material. The following morning all vomiting had ceased, and the patient's general condition had greatly improved. A large quantity of liquid feces had passed through the opening. She made an uninterrupted recovery, motions beginning to be passed by the rectum on the eighth day. At the end of a month the small opening in the cæcum was closed by operation, and the patient, now over 80 years of age, when last seen remained free from all intestinal trouble and enjoyed excellent health.

THE END OF A "PRACTICAL CURE."

[WITH CHROMO-LITHOGRAPH.]

By SIR SPENCER WELLS, BART., F.R.C.S.,
Surgeon to the Queen's Household.

In the summer of 1879 I amputated the elongated and enlarged cervix uteri of a married lady who had suffered from supposed papilloma or epithelioma, and had been treated unsuccessfully by injection of perchloride of mercury into the substance of the cervix. Examination of the part of the cervix removed proved that the enlargement was due to adenoma, or simple hypertrophy of the utricular glands. The patient was much relieved by this amputation, but about a year afterwards the remains of the cervix showed some tendency to enlarge and proliferate, and I destroyed the affected part by the actual cautery. There was some pain and fever for a few days after this, but they soon subsided, and for about seven years afterwards, until 1887, she had as a rule excellent health, surprising her friends by her improved appearance; occasionally suffering from menorrhagia but at other times walking several miles without fatigue, driving, riding, rowing, dancing, and enjoying society more than most people. Her husband died in 1887, and in February, 1888, she married again. She remained very well during that summer, and in the autumn went to Scotland, where, after a fall or blow, she suffered considerable pelvic pain. She then travelled to Edinburgh and Wales, and afterwards to Birmingham in November. She was there assured that her life was in imminent danger, that immediate operation was necessary, that she would die or become insane if it were not done, that the risk was very small, the after pain insignificant, and a cure almost certain. Her ovaries and Fallopian tubes were removed. About a month after this operation the operator exhibited these ovaries and tubes before the British Gynecological Society, and said the patient was then "practically cured." The uterine disease was entirely ignored, and I was accused of having overlooked a pyosalpinx due to a gonorrhœa from which she had suffered many years before. I took no notice of this attack for about a year, although I suspected a useless operation had been performed and the "cure" very doubtful.

Long before this I had repeatedly condemned a growing tendency to perform unnecessary mutilating operations upon women, both married and unmarried, and had protested against any teaching or example which might tend to lower the standard of surgical morality and to endanger the character of British surgery. Afterwards, when delivering the Bradshaw Lecture at the College of Surgeons, in December, 1890, I expressed my fear that "the operation of oophorectomy has been injudiciously performed without due explanation of its consequences, and with mistaken prophecies of

TO ILLUSTRATE THE PAPER BY SIR SPENCER WELLS, BART.,

Fig. 1.—*a*, Cervix from which small piece has been removed for microscopic examination; *b*, small submucous myoma; *c*, larger submucous myoma; *d*, larger submucous myoma in substance of wall; see also Fig. 1 *a*.

Fig. 2.—Mucous membrane of uterus—endometritis $\times 20$; *a*, comparatively normal gland tube near surface of mucous membrane; *b*, somewhat convoluted gland tube near surface of mucous membrane; *c*, distended tube containing altered extravasated blood, the epithelial lining is somewhat flattened, and does not stain well; *d*, cellular connective tissue (granulation tissue) in which there are numerous hæmorrhages; *e*, irregular layer of epithelium, here and there, detached.

Fig. 3.—Mucous membrane of uterus—endometritis $\times 270$; *a*, columnar epithelial lining, somewhat dilated gland tubes; *b*, small blood vessel; *c*, connective tissue cells; *d*, group of red blood corpuscles, small hæmorrhage into mucosa; *e*, granular debris of blood corpuscles in mucosa covering layer of columnar epithelial cells.

insanity or early death: that in certain circumstances the bounds generally recognised as those which should limit deliberate and consultative practice have been overstepped; that many young women who have been saved from unnecessary mutilation have afterwards borne children; that recoveries from the operation have been incorrectly counted as cures of the diseased condition, and that all failures have not been recorded." I added: "All this has excited a suspicion in the public mind—and more than a suspicion in the mind of the profession—that some of the recent expansions of abdominal surgery have not increased public respect for the profession, and require denunciation." In January, 1891, at the Obstetrical Society, I read extracts from a paper "On the Failure of the Removal of the Tubes and Ovaries to Relieve Symptoms," by Dr. Ross, of Toronto. He said "many cures I hear of as cures are not cures," and I referred to the case reported as a "practical cure" only a month after operation as, after more than a year, a "deplorable and disastrous failure." Several letters in the medical journals soon afterwards appeared, and may be seen in the first volumes of 1891. In one of these the patient herself flatly contradicted several of the statements published in the *Transactions* of the British Gynaecological Society. More than once the operator declared that the case was not a failure, but was or would be a cure; whereas I maintained that since the operation, in November, 1888, she had been a confirmed invalid, adding that "the failure to the husband has been 'disastrous' in expenditure; to two devoted daughters in health and anxiety from continuous nursing; to the patient 'deplorable,' in more than two years' almost continuous suffering and shattered health." All this may be much strengthened now. One of her daughters died of tuberculosis rapidly developed. The patient herself went from bad to worse. Menstruation, which before the operation was only occasionally excessive, became so at almost every period, and towards the end severe and dangerous hæmorrhage was almost constant. More than once she appeared to be almost dying in spite of injections into the uterine cavity and plugging the vagina, while ergot was taken very freely. In September, 1891, she went to Spa in hope that she might regain some strength there, but, in spite of constant and skilful attention from Dr. Cafferata, it was very doubtful whether she would be able to travel towards the Riviera, where she had been advised to spend the winter. She stopped at Strassburg on the way, under the care of Professor Freund. I wrote to him suggesting that extirpation of the entire uterus might be the only way to stop such repeated and alarming bleeding; but he thought it better not to do more than dilate the cervix and use the curette freely. This he did with some temporary benefit, but alarming bleeding returned while she was under the able care of Dr. Schetelig at Nervi. She became so pallid and exhausted that death was more than once expected, and it was feared that she could not survive another flooding. Vaginal extirpation of the uterus was then agreed upon after full consultation. The operation was performed in Genoa by Professor Ceci in July, 1892. The patient at first rallied well, but died on the tenth day. The uterus was sent to me by Professor Ceci; it was somewhat enlarged. The measurements were: Weight of uterus, 180 grammes; length, 12 centimetres; width, 8 centimetres. On division into two parts, the length of the uterine cavity was 9 centimetres, the distance between the two horns, 6 centimetres.

The naked-eye appearance is shown in one coloured drawing of life size, and I append a report on it, and another on the microscopical sections, both by Dr. Woodhead.

"On opening up the uterus, the mucous membrane is thickened and very irregular, and here and there are small clots, the result of hæmorrhage from the mucous membrane. On the right side near the fundus is a small submucous myoma. On the left side in the muscular wall is a similar tumour about the size of a small marble, and on incising a projection which appears on the outer surface another myoma the size of a large filbert is found (see Figs. 1 and 1a). Over the intramural and submucous tumours the mucous membrane is very much thickened, and there are well-marked extravasations of blood, not only on to the free surface, but also into the substance of the thickened membrane.

"On microscopic examination of (a) a tumour, (b) the thick-

ened mucous membrane, and (c) the cervix, the following appearances are presented:—

"(a) The tumour consists of bundles of non-striped muscular tissue, which have the usual concentric and interlacing arrangement. There is somewhat more fibrous tissue than one would expect to find in tumours of this size, from which it may be gathered, as the fibrous tissue is comparatively dense, that these tumours are of some standing.

"(b) Under a low power, in the mucous membrane, especially immediately over the small myoma in the wall of the uterus, where the evidence of hæmorrhage is most marked, the glandular layer is enormously increased in thickness; the glands, instead of being regular flattened tubes with apposed walls, and comparatively short in length, appear to have become convoluted and distended, especially at some little distance from the mucous surface. The epithelial layer on the surface of the mucous membrane is wanting at points, and appears to be readily separable. Beneath this we have what looks like ordinary granulation tissue into which there have been small hæmorrhages; the small vessels are enormously distended, and some appear to have given way. Near the mucous surface the gland tubules with their epithelium are only slightly dilated and convoluted, and the hæmorrhages into them are not specially well marked, but the deeper spaces, which are very considerably enlarged, are many of them filled with blood. In these it is often difficult to make out cylindrical epithelium; around them also the interstitial hæmorrhages are very distinctly seen, the red blood corpuscles in such cases being in various stages of degeneration.

"On examining under a high power, the matrix in which the tubules are embedded is seen to be very cellular with only delicate fibrillæ of connective tissue between. The connective tissue cells are large, the protoplasm is somewhat granular, and the nuclei are exceedingly well stained. The vascular channels are very large, and in some cases they are enormously dilated; many of them have given way, and, as a result, masses of red blood corpuscles with a few leucocytes may be seen lying in the loose cellular tissue. Near the surface the columnar epithelial cells are still distinctly seen with a layer of granular mucus on the surface, but lying in the lumen of the tube, which is often dilated, pigment granules probably derived from red blood corpuscles are found mixed with the mucus; in the deeper spaces the columnar shape of the epithelial cells is often lost, rows of flattened cells taking the place of the large columnar cells. We have here, then, endometritis of the form in which hæmorrhages are so frequently met with. There has been well-marked hæmorrhage from the vascular succulent layer, especially from the parts lying over the surface of the tumours.

"(c) In the cervix microscopical examination reveals nothing more than a somewhat atrophied mucous membrane lying on very dense fibro-muscular tissue. There is certainly nothing to indicate that recent hæmorrhages have occurred at this point, though it is probable that there has been some chronic endometritis, perhaps similar to that above, leading to the formation of fibrous tissue which, here, is even more dense than in a healthy uterus."

It can scarcely be necessary to comment on the above painful history. The case was brought forward as one of a chronic invalid whom I had treated for seven years, but "failed to relieve," although she could "have been relieved at any time." A month after operation she was said to be "practically cured." And when afterwards a repetition of so-called suppurating hæmatocœles called for treatment, the profession heard with surprise that such occurrences were not infrequent sequels to an operation which had been and still is asserted to be an almost certain cure of diseased conditions, confirming Dr. Ross's statement: "Many cases I hear of as cures are not cures," and my own that I have seen or heard of many others equally questionable.

BEQUESTS.—The late Mrs. Phoebe Springall Croker of Lavenham, Suffolk, has by her will bequeathed, among other bequests, £100 to the Suffolk General Hospital, £50 to the Zenana and Medical Mission Home and School for Ladies, and £20 to the St. Leonard's Cottage Hospital, Sudbury.

PRINCE METSKERSKY, editor of the Russian *Grashdanin*, has been sentenced to six weeks' detention in a military prison for a libel on the sanitary administration of the army.