in migraine one side of the head only may be involved in pain. The muddling of thought which occurs in headache is due to the fact that in it the special sense areas are, for reasons already given, particularly apt to be implicated, and when these are perturbed clear thought is manifestly impossible, for it is they which chiefly constitute the anatomical substratum of thought processes.

(2) The other fact is the extreme sensitiveness during headache of the special sense systems, notably of sight and hearing. The sufferer is apt to be acutely sensitive to noises and bright lights, and this alone is strong evidence of the close connection of the auditory and visual centres with the C.S. centres of the head.

It may, perhaps, be thought that all the facts just given can be explained by supposing that the connection I affirm obtains in the lowest level only, and I do not deny that some of them may be. In the photophobia of conjunctivitis, for instance, the irritation may spread from the lowest level C.S. centres of the eyeball and lids (that is, the sensory nucleus of the fifth) to the visual centres on the same level (that is, the nucleus of the optic nerve). It would be strange, indeed, were it to ascend to the highest level before passing over to the visual centres. It is quite possible, however, that the irritation spreads across at each successive level.

But I now proceed to cite a fact which not only confirms my main contention, but tends to show that the connection in question obtains in the highest nervous level—the anatomical substratum of mind. "It is very common," writes Gowers,² "for an aura first felt in the epigastrium or limbs to seem to ascend to the head, and when it reaches the head consciousness is almost invariably lost." And from this the writer concludes "that the central representation (that is, the C.S. centres) of the head, and perhaps of the brain itself" (he speaks thus guardedly, for we do not know whether the brain is centrally represented) "is connected in a special way with structures related to the mental state of consciousness." Further than this he does not go. The interpretation I put upon these phenomena is as follows. The aura, as Gowers observes, "is the result of the spread of the discharge in the brain." The discharge in these particular cases evidently spreads along the C.S. centres of the limbs and trunk until it reaches those of the head. Directly this occurs the special sense centres, from their close connection with the latter, are necessarily involved. Now all these centres (the C.S. centres and the special sense centres) are related to consciousness, but the latter are infinitely the more important in this respect, for the special sensations of sight, sound, and so forth constitute the chief elements in our concepts and percepts. Wherefore, when the special sense centres are involved in the discharge, thought necessarily becomes impossible, and very soon consciousness itself.

A CASE OF MYXEDEMA CURED BY THYROID EXTRACT.

By JOHN P. HENRY, M.D., B.Ch.DUBL.,
Lewisham.

Mrs. S. L., aged 44, suffering from myxœdema, came under my care in the spring of 1892. She dated the commencement of her illness from the birth of a stillborn child about fourteen years ago. Her face and limbs began to swell, and her mind to become torpid. Six years later she had a severe attack of illness, with great prostration. One day she was quite unconscious. Since then she has been unable to do anything, while the symptoms have gradually got worse. When I saw her the face was swollen, and the lines of expression lost. The eyelids, lips, and *alæ nasi* were enlarged and thickened. The skin was dry, branny, and anæmic, with a bright malar flush. Perspiration was absent; the hair was very thin and dry; the hands were "spade-like." There were swellings over each clavicle; the thyroid seemed to be atrophied, and there was tenderness over its site. Thought, volition, and action were very slow. She had not been able to leave the house for three or four years, having fallen several times in the street. Her memory was bad, and was getting worse. Speech was slow, and devoid of tone. She complained of an

unpleasant taste and smell. The highest temperature noted before beginning treatment was 96.6° F., and several times the mercury of my thermometer would not rise out of the bulb,

Before Treatment.

when it must have been at least as low as 94° F. The bowels were obstinately costive. She had violent epistaxis at

After Treatment.

the ages of 21, 31, and 41, and again about a year ago on several occasions. Menstruation did not commence till she was 19.

² *A Manual of Diseases of the Nervous System*, vol. II, p. 796.

It stopped in 1888, and occurred once two years ago, and again once again in June, 1891. She passed about forty ounces of urine daily. Sp. gr. 1020, acid, turbid, trace of albumen, no sugar. She has two children living. Her daughter and sister always feel cold. Her father died of liver disease (probably cirrhosis); her mother and one sister of phthisis.

Having first photographed her, I commenced on May 14th to treat her by hypodermic injections of thyroid extract, prepared, according to Dr. Murray's method, by Messrs. Brady and Martin, of Newcastle. She had biweekly injections of 30 minims from then until the beginning of August, with the exception of a fortnight in July. From August 3rd till October she had two injections in every fortnight. Unpleasant symptoms were only observed on three occasions. After the first injection she had an epileptiform fit, after which she was unconscious for an hour. Next day she felt better and warmer, the temperature having risen from 96.6° F. to 97° F., and on the second day to 98° F. After the fourteenth injection a small abscess formed. The nineteenth injection had to be stopped after 6 minims were given, owing to the onset of faintness, with great flushing of upper part of body, and pains in back. Two days after the first injection she sweated profusely, and the menses reappeared. Menstruation has been regular almost every month since then. She felt warmer than she had done for years. Her temperature usually ranged from 97° F. to normal. On each occasion a rise of about 1° F. was noticed within five minutes after the injection. The swelling of the face and limbs gradually disappeared.

On June 19th she got off her wedding ring for the first time for twelve or thirteen years. Her colour became more natural, the anæmia disappearing, and the malar flush dying away. On June 27th a dense undergrowth of young hairs was observed on the head, and on July 21st hairs were seen growing on the arms and the backs of the hands. On July 21st she had a slight attack of hæmoptysis. Shortly after she developed a cough, and some physical signs of incipient phthisis at the apex of the left lung. These have persisted since, but do not show much sign of increase, being, if anything, slightly better.

Her condition now is vastly improved. The œdema has quite disappeared, and her face looks perfectly natural, and at least ten years younger. A son who had not seen her since March says he would not know her. Her speech is rapid and natural. Her spirits are bright and memory good. She can move quite actively. The skin is moist, and there is a thick growth of hair on her scalp, and the hairs on the body have returned. The bowels are regular. The quantity of urine has increased, though I did not notice the diuretic effect after each injection noted by Mr. Hurry Fenwick.

Since the publication of Drs. Mackenzie's and Fox's papers I have been feeding her with raw thyroids. Though this method is in many ways an improvement on the hypodermic, being cheaper, less troublesome, and less risky, I cannot say that in this case it seems equally efficacious. It has prevented any return of the œdematous symptoms, but its effect on the temperature seems less marked and more evanescent. To keep her fairly warm she has to take one lobe of a sheep's thyroid every second day. She felt some nausea at first, but now takes it well. The bowels and menses, too, have not been so regular since the cessation of the injections.

In conclusion, I must bear testimony to the excellence of the extract supplied by Messrs. Brady and Martin.

THE TREATMENT OF MYXŒDEMA BY FEEDING WITH THE THYROID GLAND OF THE SHEEP.

B. J. DEE SHAPLAND, M.R.C.S.ENG.,
Exmouth.

E. P., aged 52, has been ill about ten years, her distressing symptoms gradually increasing in intensity until November 12th, 1892, when she presented the appearance shown in photograph No. 1. The head was scurfy and the hair gradually falling off, the skin cold and non-perspiring, and the nose was so enlarged that she could not breathe through it. She spoke with great difficulty and pain, and drawled her

words. She was so deaf that she could not hear the clock strike. She had a good deal of pain over the shoulders,

Fig. 1. - Before Treatment.

clavicles, and round the neck. She could only raise her head with difficulty and pain. Her heart and chest sounds were

Fig. 2. - After Treatment.

normal; the kidneys and liver were also healthy; both wrists were so enlarged that the pulse could not be felt.

I commenced treatment on November 12th, and immedi-