It stopped in 1888, and occurred once two years ago, and again once again in June, 1891. She passed about forty ounces of urine daily. Sp. gr. 1020, acid, turbid, trace of albumen, no sugar. She has two children living. Her daughter and sister always feel cold. Her father died of liver disease (probably cirrhosis); her mother and one sister of phthisis.

Having first photographed her, I commenced on May 14th to treat her by hypodermic injections of thyroid extract, prepared, according to Dr. Murray's method, by Messrs. Brady and Martin, of Newcastle. She had biweekly injections of 30 minims from then until the beginning of August, with the exception of a fortnight in July. From August 3rd till October she had two injections in every fortnight. Unpleasant symptoms were only observed on three occasions. After the first injection she had an epileptiform fit, after which she was unconscious for an hour. Next day she felt better and warmer, the temperature having risen from 96.6° F. to 97° F., and on the second day to 98° F. After the fourteenth injection a small abscess formed. The nineteenth injection had to be stopped after 6 minims were given, owing to the onset of faintness, with great flushing of upper part of body, and pains in back. Two days after the first injection she sweated profusely, and the menses reappeared. Menstruation has been regular almost every month since then. She felt warmer than she had done for years. Her temperature usually ranged from 97° F. to normal. On each occasion a rise of about 1° F. was noticed within five minutes after the injection. The swelling of the face and limbs gradually disappeared.

On June 19th she got off her wedding ring for the first time for twelve or thirteen years. Her colour became more natural, the anæmia disappearing, and the malar flush dying away. On June 27th a dense undergrowth of young hairs was observed on the head, and on July 21st hairs were seen growing on the arms and the backs of the hands. On July 21st she had a slight attack of hæmoptysis. Shortly after she developed a cough, and some physical signs of incipient phthisis at the apex of the left lung. These have persisted since, but do not show much sign of increase, being, if anything, slightly better.

Her condition now is vastly improved. The œdema has quite disappeared, and her face looks perfectly natural, and at least ten years younger. A son who had not seen her since March says he would not know her. Her speech is rapid and natural. Her spirits are bright and memory good. She can move quite actively. The skin is moist, and there is a thick growth of hair on her scalp, and the hairs on the body have returned. The bowels are regular. The quantity of urine has increased, though I did not notice the diuretic effect after each injection noted by Mr. Hurry Fenwick.

Since the publication of Drs. Mackenzie's and Fox's papers I have been feeding her with raw thyroids. Though this method is in many ways an improvement on the hypodermic, being cheaper, less troublesome, and less risky, I cannot say that in this case it seems equally efficacious. It has prevented any return of the œdematous symptoms, but its effect on the temperature seems less marked and more evanescent. To keep her fairly warm she has to take one lobe of a sheep's thyroid every second day. She felt some nausea at first, but now takes it well. The bowels and menses, too, have not been so regular since the cessation of the injections.

In conclusion, I must bear testimony to the excellence of the extract supplied by Messrs. Brady and Martin.

THE TREATMENT OF MYXŒDEMA BY FEEDING WITH THE THYROID GLAND OF THE SHEEP.

B. J. DEE SHAPLAND, M.R.C.S.ENG.,
Exmouth.

E. P., aged 52, has been ill about ten years, her distressing symptoms gradually increasing in intensity until November 12th, 1892, when she presented the appearance shown in photograph No. 1. The head was scurfy and the hair gradually falling off, the skin cold and non-perspiring, and the nose was so enlarged that she could not breathe through it. She spoke with great difficulty and pain, and drawled her

words. She was so deaf that she could not hear the clock strike. She had a good deal of pain over the shoulders,

Fig. 1. - Before Treatment.

clavicles, and round the neck. She could only raise her head with difficulty and pain. Her heart and chest sounds were

Fig. 2. - After Treatment.

normal; the kidneys and liver were also healthy; both wrists were so enlarged that the pulse could not be felt.

I commenced treatment on November 12th, and immedi-

ately the beneficial action of the gland was visible by inducing refreshing sleep and gentle perspiration. On November 19th she expressed herself as feeling better than she had done for years. On November 22nd the swelling of the face was much diminished, and she could now hear the clock strike.

On December 3rd the wrist pulse could be felt. On December 17th she said that she felt quite well and could easily breathe through her nose. On December 28th the swelling of the legs had all but disappeared. On January 20th photograph No. 2 was taken.

I gave her half an underdone thyroid gland the first thing every morning taken with ordinary food, from November 12th to January 12th; no bad symptoms arose during the treatment, except the last week, when there was some feeling of discomfort (tired) about the neck and shoulders.

February 5th. She is now free from pain and feels quite well.

CANCERUM ORIS, FOLLOWED BY EXTENSIVE ULCERATION OF THE CHEEK AND ANKYLOSIS OF THE JAW: RECOVERY.

By JOHN WARD COUSINS, M.D.LOND., F.R.C.S.,
Senior Surgeon to the Royal Portsmouth Hospital and the Portsmouth
and South Hants Eye and Ear Infirmary.

N. R., aged 10 years., was sent to me by Dr. B. H. Mumby, the medical officer of the Infectious Diseases Hospital, Portsmouth. The child had been under treatment for a severe and prolonged attack of typhoid fever, which had been followed by destructive ulceration of the right cheek. On admission, on March 15th, 1892, the little patient was much exhausted, and bedsores had formed over the hips and sacrum. The perforation of the cheek was as large as a crown piece, its edges were extremely thin, and several teeth were visible through the opening. (Fig. 1.) The dense cicatricial tissue

Fig. 1.

bound the maxillæ firmly together, and the buccal cavity was completely obliterated. The jaws were so tightly closed that the child was fed with difficulty through a gap between the

teeth. After a few weeks' careful treatment the bedsores began to heal, and the general condition of the patient considerably improved. A false joint was then made in front of

Fig. 2.

the line of adhesion by subcutaneously dividing the lower jaw, and excising a small wedge of bone from the lower border. Three weeks after the operation solid food was taken readily,

Fig. 3.

and mastication was performed with comfort. The dense cicatricial structures around the opening in the cheek were