from the bases of his lungs; he was able to move his neck more freely, and altogether felt stronger.

On February 19th Dr. Carter again saw him with me, and expressed surprise at the manifest improvement. We noted that he was passing more than a quart of urine in twenty-four hours; it was concentrated, deposited urates on cooling, specific gravity 1028, acid, and contained a faint trace of albumen. The oedema had disappeared from the subcutaneous structures, the breath sounds and percussion notes were normal, and there was no evidence of fluid in the peritoneal cavity. He now lay down comfortably, the voice was clear and strong, and he had no stridor. The mobility of his neck had greatly increased, and there was decided softening of many of the glands which had previously been so hard and large. The neck now measured $2\frac{1}{2}$ inches in circumference.

More than a month has passed since this change in the patient's condition began, and he has never relapsed into his former state of danger and misery. Considering the long period during which, week by week, he became worse and worse, and at length appeared to be near death, and the rapid improvement which followed the administration of the marrow, one is tempted to think and to hope that something more than coincidence has accounted for the altered condition in which we now find him. The glands are certainly much softer, and if there has been no actual decrease in their size, they certainly have not increased, either in their number or dimensions, so that something like an arrest in the progress of the disease seems to have taken place.

The extraordinary success which has resulted from the treatment of myxœdema by the thyroid secretion during the past year is, I think, explainable on the same principle which I have laid down as the basis of these remarks. Here it has been proved, both clinically and experimentally, that the absence of the thyroid secretion entails the development of a condition in which there is an excess of mucin not only in the subcutaneous tissues, but also in many other connective tissues. It is well known that mucous tissue is developmentally the normal predecessor of ordinary connective tissue, the white fibres being produced in the jelly-like matrix (according to Rutherford) by the production of a periplast by the cell of the mucous tissue, which elongates, and eventually becomes a white fibre. In myxœdema, therefore, there is an over-production of the rudimentary progenitor of connective tissue, a certain consequence of the loss of a secretion which normally prevents its formation, and it seems a reasonable explanation that the thyroid adds something to the blood which controls the growth of the predecessor of connective tissue, and, therefore, of the fully formed tissue itself. This appears to me to be more probable than the theories which ascribe to it powers which prevent auto-intoxication, and it does not seem to me impossible that the application of this principle may ultimately lead us to treat some local overgrowths of tissues either embryonic or more fully formed, such as sarcoma and carcinoma, by the introduction into the system of materials which may inhibit their production.

REMARKABLE CASES OF GASTRIC ULCERATION AND THEIR RESULTS.¹

By SAMUEL WEST, M.D., F.R.C.P.,

Assistant Physician to St. Bartholomew's Hospital, Senior Physician to the Royal Free Hospital, etc.

GASTRIC ulcer is associated in many, perhaps in most, cases with such striking and definite symptoms as to make the diagnosis both easy and positive. Yet of the two characteristic symptoms—pain and hæmatemesis—the pain may be due to dyspepsia only, and hæmatemesis occur without any actual ulcer. Hæmatemesis, again, is not a necessary, but only an accidental phenomenon even when gastric ulceration exists. As long as pain is the only symptom and hæmatemesis has not yet occurred, the diagnosis between dyspepsia and gastric ulceration must, in most cases, remain uncertain. To make the difficulties greater still, gastric ulceration may

exist and last for a long time, not only without hæmatemesis, but even without pain or gastric symptoms of any kind. Such cases as these may be described as latent gastric ulceration.

Post-mortem observation proves that gastric ulcer is a far commoner lesion than clinical records indicate, for a scar, or even many scars, may yield conclusive evidence of past ulceration, for which there may be little or nothing definite in the clinical history to account.

The object of this paper, however, is not to prove that gastric ulceration may be latent, in other words, may exist for long without definite symptoms, for this requires no further proof, but to show by a series of cases which have been under my own observation, in what strange form these latent gastric ulcers sometimes present themselves in practice.

CASE I. Chronic Gastric Ulcer for Eighteen Months; Acute Peritonitis; Gangrenous Abscess in Left Loin; Incision; Suppurative Peritonitis the Result of Perforation; Death.—T. L., 34, was active and well, and had never had any illness until eighteen months ago, when he was suddenly seized with hæmatemesis. Before that time he had never had any pain or discomfort in the stomach, and believed himself to be perfectly well. This attack kept him in bed for a week, at the end of which time he was able to return to work. Three months later he had another attack of pain, but without hæmatemesis, and since then has been subject to attacks of pain, often of great severity, and has hardly ever been free for more than a few days at a time, so that he has lost much flesh and strength. A fortnight before admission he was attacked with cold shivering, diarrhoea, and vomiting. Although the shivering and vomiting were not constant, the diarrhoea continued without intermission, and was very frequent.

The man looked very ill and anxious. The temperature was 100° only, the pulse 108, and the respirations 24. The tongue was dry, red, and glazed. He complained greatly of thirst, and was constantly sick, being hardly able to retain anything at all on the stomach. The abdomen was distended, and in the left lumbar region was a local swelling occupying the whole side. This felt very tense and elastic, but resonant everywhere except at the lower part as the patient lay on the back. Opium relieved somewhat the vomiting, but the next day, the patient's condition not having in any way improved, I invited the assistance of Mr. Battle, who agreed with me that the swelling in the loin was due to a localised peritonitis, and that it should be opened. Accordingly this was done by him, and a large cavity was opened from which a great quantity of fetid pus and foul gas escaped, showing the swelling to have been a gangrenous abscess. The cavity extended apparently behind the intestines, and was thought to be shut off from the peritoneal cavity. Upwards it passed toward the middle line, and extended further than the finger could reach. It was well washed out, and a drainage tube inserted. The operation, though it relieved the patient, did not cause any material improvement in his general condition, for the prostration continued to increase, and the diarrhoea and vomiting persisted, though perhaps hardly to the same degree as before the operation. Four days later the urine was retained, and, as the patient was found to have a bad stricture, the bladder was aspirated, after which he passed water naturally.

The strength continued to fail, and the patient died of exhaustion on February 26th, three weeks from the time that he had been taken ill. There was nothing further worthy of note except that two days before death he developed a parotid abscess. At the *post-mortem* examination, the abscess cavity in the left loin was found to extend into the pelvis, and from thence upwards to the diaphragm and under the left lobe of the liver to the stomach. Here the stomach and the liver were adherent, and a perforation was discovered the size of a threepenny-piece. On opening the stomach, this perforation was found to correspond with the centre of an old gastric ulcer of no great size. An abscess existed on the right side in a similar position between the right lobe of the liver and the right kidney, and a third, though smaller one, lay in the middle of the matted coils of intestines. Besides this suppurative peritonitis, there were no lesions in the body. There can be little doubt that in this case it was the old gastric ulcer that was the cause of the peritonitis, which had probably commenced as a small abscess near the pylorus and spread to the left loin, the other abscesses being of later date.

CASE II. Chronic Gastric Ulcer; General Suppurative Peritonitis; Suppurative Pylephlebitis; Death.—A. G., a domestic servant, aged 22, was admitted into the hospital looking very pale, ill, and with drawn and sunken features; respirations, 36; pulse, 136; temperature, 101° . She lay with the knees drawn up; the abdomen was rigid and tender, and did not move with respiration. The diagnosis was made of acute peritonitis, possibly suppurative. She gave the following history that ever since the age of 15 she had suffered on and off from pain and vomiting after food, the vomit having sometimes been streaked with blood, but no more. During the last six months she had been much better, until on April 9th she slipped on the stairs, and was at once seized with such severe pain that she had to lie up. A day or two later she became hot and thirsty. She did not shiver at any time, but sweated profusely. The diagnosis when she was admitted into the hospital was acute peritonitis, probably due to the perforation of a gastric ulcer, or to the rupture of some adhesions surrounding it. From this time, however, all abdominal symptoms vanished, and pain was complained of in the left pleura. Cough developed, and a patch of dulness was found in the left base. This was thought to be an empyema, and a needle inserted for the purpose of diagnosis drew off a little pus. The diagnosis of empyema seemed to be confirmed, when a few days later she began to expectorate pus. It was decided then to explore the chest with a view to incision, but though a small needle was inserted several times no pus could be found, and the operation was abandoned. The temperature chart was very irregular, but not characteristic of anything except hectic of an irregular type. There was a remarkable absence of rigors, the patient had two shivering fits, and those within a week of one another, and the first not until she had been many weeks ill. It was evident that the patient had pus somewhere; at first it was thought in the abdomen and then in the pleura; but exploration having failed in the pleura, and the abdominal symptoms having subsided, and the abdomen being not

¹ Read before the East Surrey District of the South Eastern Branch of the British Medical Association.

tumid but rather retracted, and not tender to the touch, it was thought that the pus must be deepseated, and out of reach. The patient died of asæthia.

On *post-mortem* examination the peritoneum was found full of pus, most in the pelvis, into which it had gravitated. The intestines were adherent to each other and to the diaphragm, as well as to the liver and spleen. The adhesions were most dense near the stomach, and on opening that organ a large ulcer was found on the lesser curvature, nearly circular, and measuring three-quarters of an inch across. Its edges were rounded, and almost cartilaginous, the base was thickened and irregular, and formed of thickened omentum. The ulcer was adherent to the under surface of the liver, and lay in about the centre of the chief mass of fibrous adhesions described, but there was no communication with the peritoneum.² The liver was a little enlarged, and on section seemed to contain many abscesses, but these abscesses proved on examination to be branches of the portal vein distended with pus, suppurative pyelophlebitis. The main trunk of the portal vein contained an old suppurating clot, and the splenic vein, like the branches of the portal vein, were filled with pus. Neither the portal or the splenic vein contained any red blood or any recent clot. In this again, as in the last case, the cause of death was suppurative peritonitis, but the cause of the peritonitis lay in the old gastric ulceration.

CASE III. Chronic Ulcer of Stomach and Duodenum; Perforation of Duodenal Ulcer; Local Abscess opening into Transverse Colon.—The patient, a woman aged 43, a servant, had been in good health and active work until seized one day a month before admission with violent pain in the pit of the stomach. The pain at first was very severe, like cramp, and seemed to take away all her strength, but she did not vomit. She was taken to bed, where she remained for a fortnight, when she tried to get up, but had to go back again, and the doctor told her she had inflammation of the stomach. The pain increased, and a week later diarrhoea set in, and has continued more or less ever since. Throughout she has never vomited, but has been able to take food, both solid and liquid, though without appetite. She has lost flesh and strength rapidly, and during the last few days has been troubled with cough and perspirations, especially at night. The patient looked pale and somewhat dusky, and was bathed in perspiration. The pulse was 140, the respirations 60, and the temperature only 99°. There was considerable general bronchitis, and the heart sounds were feeble. The abdomen was distended and contained a little fluid. The right hypochondriac region was tender and slightly prominent, and the liver appeared to extend from the fifth rib to the umbilicus, but the lower part of this area was resonant. The feet were slightly swollen. Beyond that the patient was in a septic state, and probably had some clots in the lungs, no exact diagnosis appeared to me possible. The subsequent history threw no light upon the case, nor did any fresh symptoms develop, except that two days before death some dark blood was passed by the bowel. The patient rapidly sank and died of exhaustion four days after admission.

At the necropsy, on making the incision into the abdomen, a superficial abscess was opened, which lay immediately below the abdominal walls. The lower edge of this corresponded with what was thought to be the liver, and, as it contained air, accounted for this part being resonant. Otherwise the area ascribed to the liver was correct, for that organ extended from the upper border of the fifth rib to nearly the level of the umbilicus. The abscess described had two openings in its walls; the one of small size led into the transverse colon. It was obviously recent, and probably accounted for the hæmorrhage during the last few days of life. The second opening was of larger size and older date, and lay more posteriorly. It proved ultimately to be connected with the duodenum. On laying the stomach and duodenum open, two chronic ulcers were found, one in each, with thickened fibrous walls, and evidently of long standing. That in the stomach was seated on the lesser curvature, and was about the size of a five-shilling piece. It had made its way through the walls of the stomach, and its floor was formed by an excavation in the liver, but no communication existed between this ulcer and the abscess cavity. The duodenal ulcer was of smaller size, not quite so large as a shilling, and in its centre was the larger of the two openings described in the abscess walls. Besides that, both femorals were occluded by clot, which accounted for the œdema; there was nothing further of note in the rest of the body.

The *post-mortem* examination gives a full explanation of the clinical history of the case. The only diagnosis ventured upon during life was confirmed by the discovery of the air-containing abscess, and the cause of this was found in the perforation from the duodenal abscess. The hæmorrhage from the bowel was no doubt due to the bursting of the abscess into the colon. The gastric ulcer took no part in the production of symptoms, but the thickness of its walls, the excavation it had formed into the liver, and the induration round it were conclusive proofs of its long duration. The duodenal ulcer, though subsequent in formation to that in the stomach, must also, from the thickness of its walls, have been of some standing. Now it is remarkable that not only had these ulcers existed without symptoms until that in the duodenum perforated, but that even after the perforation there was not the slightest other symptom to suggest their existence.

CASE IV. Malignant Ulceration of the Stomach; Perforation of the Colon.—The patient, a man, aged 40, was admitted into St. Bartholomew's Hospital with the diagnosis of chronic intestinal obstruction. He stated that he had been in good health and at work until three weeks before admission, and that then he was suddenly seized, while carrying a heavy weight, with severe pain in the epigastrium. He had taken his dinner about half an hour before, and had suffered no discomfort from it. The abdomen swelled soon after, and severe vomiting set in, which relieved the pain. Since that time he was laid up, and was more or less in bed. The pain returned after every meal, and continued until relieved by vomiting. He had not at any time brought up blood. The bowels were irregular, but never obstinately confined. A week after his first attack the vomit became foul smelling, and continued to be of the same character on and off, though at times a day or two would pass without its being offensive or consisting of anything but partly digested food.

On admission the patient did not look specially ill. The tongue was

slightly coated. The breathing was not free, because of pain in the epigastric region, caused by taking a deep inspiration. The abdominal pain was more or less constant, being usually of a gnawing character, but occasionally gripping. There was great hyperæsthesia, as well as tenderness on firm pressure, and considerable muscular resistance. The abdomen was not tumid, nor was there any distension of either small or large intestine. The temperature was normal, and the pulse and respirations only slightly accelerated.

The symptoms not appearing urgent, the patient was placed upon milk and lime water as diet, and bismuth and opium as medicine, and an enema administered. The diagnosis presented many difficulties. The intermittent symptoms, long duration, and the absence of abdominal distension, seemed to negative acute obstruction of either the large or the small intestine. The more chronic form could not be so definitely excluded, but the absence of abdominal distension rendered it unlikely. Impaction of feces was negated by rectal examination, and no growth or stricture could be felt within reach. Chronic peritonitis might explain the irregular vomiting and action of the bowels, but, if so, it would have to be localised. The most probable diagnosis was thought to be that of a communication between the stomach and colon, but against this was the absence of any previous history of mischief either in the stomach or colon. The age, especially, was opposed to malignant ulceration, which such a communication appeared to require. The diagnosis, it was concluded, must, however, be between gastro-colic fistula and a localised peritonitis.

On August 18th, three days after admission, the patient brought up suddenly a large amount of vomit, in colour and odour similar to a copious motion which was passed almost at the same time. On the 21st, 24th, 30th, and 31st similar attacks of fecal vomiting occurred. Although no further special symptoms developed, the patient all the time was rapidly losing ground, and by September 2nd it was obvious that if no relief could be given he would shortly die.

Under these circumstances the advice of my colleague, Mr. Cripps, was taken, and he advocated an exploratory incision into the abdomen. An incision was accordingly made into the left loin, but nothing abnormal could be found, and the wound was stitched up again. Two days later the patient died. At the *post-mortem* examination the transverse colon was found to be adherent to the greater curvature of the stomach and communicating with the stomach through a ragged hole three quarters of an inch in diameter. On opening the stomach this hole was found to be in the centre of a large ulcer, evidently malignant. The ulcer extended round the stomach transversely, except for 2 inches at the lower curvature, and was in its widest part 3 inches broad and nowhere less than 1 inch. With the exception of two small nodular masses in the adhesions between the stomach and colon, there were no secondary deposits. Nor was there elsewhere in the body any disease. Microscopical examination showed the tumour to be a medullary carcinoma.

During life the symptoms were most puzzling, for although on their merits they seemed to point to a gastro-colic fistula, still the difficulties of explaining the origin of such a lesion seemed insurmountable. In the first place, there had been no evidence whatever of stomach trouble previous to the supposed perforation, and, secondly, such perforations were generally due to malignant disease of either colon or stomach, and against the assumption of malignant disease was the age of the patient as well as the absence of symptoms. The *post-mortem* examination, however, made it clear, not only that ulceration existed, but that it was malignant. Moreover, the extent of the ulcer showed that it must have existed for some considerable time; while the history makes it equally certain that the ulceration continued without symptoms of any kind until perforation occurred.

ON CENTRAL BIRTH PALSY.¹

By THOMAS OLIVER, M.A., M.D., F.R.C.P.,
Physician to the Royal Infirmary, Newcastle-upon-Tyne.

At a previous meeting of this Branch I read a paper on paralysis occurring in children, accompanied by loss of speech, and due to meningitis. The boy whose case I now describe is suffering from the effects of a lesion developed at the time of his birth, to which the term "birth palsy" is applied.

CASE I. Difficult Parturition; Child Resuscitated with Difficulty; Convulsions; Atrophic Hemiplegia.—R. H. is 14 years of age. When he first came under my notice three years ago the following was the note taken: There is nothing of importance in the family history. His mother states that she had good health during the pregnancy which concerns our patient; that the labour was a severe one, the forceps requiring to be used. When the patient was born there were no signs of vitality in him; breathing was completely suspended, and it was some time before the infant could be sufficiently roused to utter a "cry." Very shortly after his birth convulsions developed and continued more or less for the next three days. Since then there has been complete freedom from convulsions; the patient's health has been remarkably good.

He is somewhat deformed, and when standing or walking he leans rather to the right side, with his left foot slightly drawn up. His left arm is flexed at the elbow and is held forwards, so that when walking the left arm is dangled in front of him and is observed to be the seat of tremor. His mouth is always half open, and, though this gives him rather a "weak look," there is no mental defect; far from it, the lad is above the average in ability; he is prompt and exact in his replies to questions addressed to him at school and elsewhere. This is a point of great interest, for his speech is slow and drawing, and it is in these cases where articulation is defective that there is generally noticed some intel-

¹ Read at a meeting of the North of England Branch of the British Medical Association.

² Specimen and condition described in *Path. Soc. Trans.*, xli., 146.