the main proteid constituent of all protoplasm is nuclealbumin.

PRACTICAL BEARINGS.

During these lectures I have from time to time stepped aside from my subject proper, in order to point out certain practical bearings which I considered to be of importance. I wish that I had had time, and, still more, the necessary experience to have done so more fully. If, however, any grains of usefulness are scattered through what I have said, there is no doubt that others will pick them up, and work out their practical side. In connection with blood coagulation and thrombosis I need not stop to point out the immense importance of a correct knowledge of this subject to the pathologist, and I will content myself with two remarks only on its practical aspects.

The first of these is to point out the great resemblance between the phenomena observed in the animals in the experiments described, and those cases of pulmonary thrombosis which are occasionally seen in the human subject. One can hardly doubt that here we have to deal with poisoning by the same material. Professor Gerald Yeo was witnessing some of our experiments a few days since, and he described to me a case he had had in King's College Hospital. It was one of amputation of the breast for cancer, and the wound was perfectly aseptic. But ten days after the operation, without any warning, the patient died suddenly with great dyspnoea. Examination of the brain revealed no lesion, but a firm fibrinous clot was drawn like a tree with its branches from the pulmonary artery at the necropsy.¹¹ The dyspnoea in this and similar cases points to the cause of death in the lungs themselves rather than in the respiratory centre; this difference is easily explicable on the grounds that the dose of poison enters the circulation more slowly than in our experiments, where it is injected.

The second practical point which I wish to emphasise is a word of warning. When I have seen animals die in a few seconds, as though by a lightning flash, after a dose of 5 cubic centimetres of this poison, which could not have contained more than a few centigrammes of the material, I feel very loth to urge that any practical use of such substances should be made in the human subject, either for the stoppage of hæmorrhage, or for the production of a clot in aneurysm. The cure would in such instances be worse than the disease.

THE ANTI-CHOLERA VACCINATION :

AN EXPERIMENTAL CRITIQUE.

Read before the Pathological Society of London.

By E. KLEIN, M.D., F.R.S.

HAVING for some time been engaged for the medical department of the Local Government Board in investigating the antagonistic action of various pathogenic bacteria, I have during the last year extended these investigations to the anti-cholera vaccination, as described by M. Haffkine of the Pasteur Institute. Haffkine, by an extensive series of experiments, has shown that if a large dose of choleraic comma bacilli, grown on and then removed from the surface of solidified nutrient agar, be injected (as a suspension in sterile beef bouillon) into the peritoneal cavity of a guinea-pig, a fatal result is produced in, or within, twenty-four hours, the peritoneal cavity containing fluid exudation full of the living comma bacilli. R. Pfeiffer¹ had already shown that this result is produced by injecting intraperitoneally large masses of the cholera bacilli, either living or after having been previously killed by heat or drying, by chloroform, ammonium sulphate, alcohol, or other means, and he justly concludes from these experiments that the comma bacilli themselves contain in their protoplasm a poisonous principle. The fatal result in the guinea-pig thus obtained with large doses of the cholera bacilli was shown to be due to acute severe peritonitis, and it was further shown that the same result is obtained by the same methods with the vibrio of Finkler and the vibrio of Metschnikoff, so that also these two species include in

their substance a similar poisonous principle. Notwithstanding these facts, Pfeiffer regards the intracellular poison of the cholera bacilli as "the primary cholera poison."

Haffkine starts from these experiments, and finds that by transmitting, by intraperitoneal injection, the (peritoneal) exudation of a guinea-pig dead after the above injection through a succession of guinea-pigs (twenty to thirty generations), an exudation fluid is ultimately obtained of which the cultivation on agar possesses a high degree of virulence, "virus fort," so much so that that a small dose of this cultivation injected intraperitoneally into the guinea-pig produces a fatal result in eight to twelve hours, the fatal dose of this "virus fort" being in amount only a-fifth, a-sixth, or less of that required of the original cholera culture. Repeated subcutaneous injections of small doses of this "virus fort" produce in the guinea-pig a transitory constitutional illness; at the seat of inoculation there appears a soft oedematous swelling, which, after a few days, diminishes in size, becoming at the same time firmer, and is either totally absorbed or more generally produces a local necrosis and ulceration of the skin, which ultimately completely heals up. Guinea-pigs which have thus been twice subcutaneously inoculated with small doses of the "virus fort," generally attenuated by the addition of phenol, prove themselves refractory—that is, "vaccinated" against the intraperitoneal injection of a fatal dose of the "virus fort," a dose that invariably kills control guinea-pigs in eight to twelve hours. I have made a considerable number of experiments on guinea-pigs in a similar direction; the results of these I now propose, with the sanction of the Medical Department of the Local Government Board, to communicate to the Society. These results may be arranged in the following order:

1. Experiments of intraperitoneal injection were made with beef bouillon suspension of various microbes, grown on the surface of solidified nutrient agar; the amount used for each animal being in all cases one-fifth to one-half of a culture tube; each culture tube presenting an agar surface of 6 centimetres by 2 centimetres. The microbes used were: (a) the cholera bacilli; (b) the vibrio of Finkler; (c) the bacillus coli; (d) the proteus vulgaris; (e) the bacillus prodigiosus; and (f) the bacillus of typhoid fever.

2. In all cases the intraperitoneal injection of one-fifth to one-half of a culture was followed after a few hours (4 to 8) by distinct illness; the temperature is lowered, the animals are quiet, do not move and do not feed; after about 8 to 12 hours, sometimes earlier, this condition is more pronounced; next day, that is within 20 to 24 hours, the animals as a rule are found dead. On *post-mortem* examination there is found intense peritonitis with pseudo-membranes on the omentum and liver; the intestine and particularly the serous covering of the stomach are much congested. The small intestine is either relaxed and filled with sanguineous mucus, or is more or less contracted. The peritoneum contains more or less turbid, slightly sanguineous, fluid exudation filled with the microbes used for the injection. The heart's blood also contains the microbes, as cultivations prove; the colonies obtained by such cultivations are always fairly numerous, but of course not so numerous as those of the peritoneal exudation. It must be insisted on, that in all this there is absolutely nothing to warrant the assertion that the results produced by the cholera bacillus are comparable to cholera. The symptoms of the disease and the pathological changes produced in the guinea-pigs by the intraperitoneal injection are for all the above species exactly the same, but it ought to be mentioned that the cultivations of bacillus coli, bacillus of typhoid, and bacillus prodigiosus act more virulently than those of the cholera vibrio or the vibrio of Finkler, since the dose of the latter, in order to be fatal, has to be larger than that of the former, and the appearances produced by the same amount of culture, though identical in character, are more intense in the case of the bacillus coli and typhoid, and bacillus prodigiosus, than in that of the vibrio of cholera and of Finkler respectively.

3. The same result is also produced by the intraperitoneal injection of sterilised culture (exposure to 62° to 65° C. for 10 to 12 minutes), the amount required to produce a fatal result being, however, a little larger than that required of the living bacilli.

4. As only the bacteria themselves (scraped off from the

¹¹ For numerous similar cases see Sir J. Fayer's *Clinical and Pathological Observations in India*, pp. 94 et seq.

¹ *Archiv. Hygiene und Infektionskrankheiten*, vol. x1, part 3, p. 393.

surface of the agar culture) are used for these injections, it follows that all these six different species of bacteria contain in their protoplasm a poisonous principle, which appears to be the same for all species, since the disease produced and the anatomical changes found on *post-mortem* examination are in all cases identical.²

5. Non-fatal doses (one-fifth to one-tenth of a culture) injected intraperitoneally produce a transitory illness; after recovery, the guinea-pigs are found refractory against the action of otherwise fatal doses of the same species, injected intraperitoneally—that is to say, doses which are fatal to control guinea-pigs. This refractory condition is obtained by non-fatal doses both of the living and of the sterilised cultures.

6. The refractory condition produced by intraperitoneal injection of a non-fatal dose of one of the above six species holds good against all the other five; thus guinea-pigs after surviving the intraperitoneal injection of non-fatal doses of vibrio of Finkler, of proteus, of bacillus coli, of bacillus prodigiosus, or of typhoid bacillus respectively, are found refractory against a fatal peritoneal dose of cholera bacilli. I have experimented with a considerable number of guinea-pigs which had survived the peritoneal injection of non-fatal doses of the vibrio of Finkler, of the bacillus coli, and bacillus prodigiosus respectively; on a second intraperitoneal injection with fatal doses of the cholera bacilli—nay, even of Haffkine's "virus fort," they proved themselves completely protected. These experiments furnish a further confirmation of the above conclusions—namely, that the intracellular poison of these different bacterial species is identical, since they mutually protect the animals.

7. The subcutaneous injection of cultures of all these different species into guinea-pigs produces the identical disease; a transitory general illness, a local soft cedematous swelling, which after a few days diminishes in size and becomes firm, and either altogether becomes absorbed or leads to local necrosis and ulceration of the skin; the latter only takes place after injection of large doses of the culture. The effect produced by bacillus coli, bacillus prodigiosus, and bacillus of typhoid is a little more intense than that produced by the others; living culture is more virulent than sterilised culture, but in its general characters is the same for both.

8. Just as is the case in Haffkine's experiments with the cholera vibrio, so also with the vibrio of Finkler, the bacillus coli, the bacillus of typhoid, and the bacillus prodigiosus, the intraperitoneal transmission through a series of guinea-pigs produces a marked increase in the virulence of the peritoneal exudation. In the case of the bacillus coli, bacillus prodigiosus, and bacillus of typhoid, already after two transmissions the subcutaneous injection of a few drops of the peritoneal exudation or of a culture grown from this produces a fatal effect, whereas a considerably larger dose of the original culture fails to do more than produce a local tumour. In the case of the vibrio of Finkler also, the transmission of the peritoneal exudation through a few successive guinea-pigs produces an exudation, of which a few drops subcutaneously injected produce a pronounced local tumour, leading to necrosis and ulceration of the skin, such as is only produced by large doses of the original culture.

9. Animals which have been repeatedly injected subcutaneously, either with living or with sterilised cultures (all from the surface of solid agar), prove refractory against the intraperitoneal injection of fatal doses of the living cholera cultures. I have a number of guinea-pigs which had been injected subcutaneously on several occasions with sterilised cultures of vibrio of Finkler, of bacillus coli, and of bacillus prodigiosus. They were, after the lapse of ten to twelve days, subjected to intraperitoneal injection with fatal doses of Haffkine's "virus fort," but they proved themselves refractory. I may mention one such series of experiments: (a) Six guinea-pigs were subcutaneously injected with sterilised (62° C. for ten minutes) vibrio of Finkler on four occasions (one-fifth of a culture tube being used for each animal on the first, third, fourth, and sixth day). Twelve days after the

last injection they were injected intraperitoneally with Haffkine culture of "virus fort," one-twelfth of a culture of this latter being used for each animal. Two control guinea-pigs injected at the same time with one-twelfth culture of this same "virus fort" were found dead the next morning, but only one of the above six Finkler guinea-pigs died, after about thirty hours, the other five remained alive. (b) Four guinea-pigs were subcutaneously injected with sterilised culture of bacillus prodigiosus on four occasions (one-fifth of a culture tube being used for each on the first, third, fourth, and sixth day). Twelve days after the last injection they were each injected peritoneally with one-twelfth of Haffkine culture of "virus fort." All remained alive. (c) Four guinea-pigs were subcutaneously injected on three occasions (one-fifth of a culture tube being used for each animal on the first, second, and fourth day) with sterilised culture of bacillus coli. Twelve days after the last injection they each received into the peritoneum one-twelfth of the same Haffkine's "virus fort." All four guinea-pigs remained alive.

10. The different species of microbes thus experimented with, including the cholera vibrio, contain in their substance the same poisonous principle, but this must be carefully distinguished from the poisonous substances (ptomaines, specific toxins, and others) which are elaborated by them in nutritive media, including the animal body. While the intracellular poison, as is shown by the above experiments, appears to be identical for all the above species (judging from their identical action on the guinea-pig after subcutaneous and intraperitoneal injection, and from their mutual protection), the latter, that is the specific toxins produced in the nutritive media, differ (a) from one another, and (b) from the intracellular poisons. As regards (a) this is well known; as regards (b) several cases in point may be mentioned. Tuberculinum of Koch contains the intracellular poison of the tubercle bacilli; as is well known, it produces on subcutaneous injection into tuberculous individuals a definite phlogo- and pyo-genic reaction; but when the tubercle bacilli grow in the body of a suitable animal, they produce additional pathological processes, namely, the characteristic giant cell formation, the spreading necrosis and caseation. Or take the proteus vulgaris: the effect of large doses of the bacilli themselves after intraperitoneal and subcutaneous injection, particularly the latter, have been described above; these effects are different from those resulting from the subcutaneous injection of small doses of the ptomaines produced by the proteus growing in albuminous matter.

Again, in the case of the diphtheria and tetanus bacilli, the well-known effects produced on inoculation of these bacilli are those of their specific toxins elaborated by them at the seat of inoculation and absorbed by the system, not of the bacilli themselves—that is, the intracellular poison—since these microbes do not live in the blood or tissues of the infected bodies.

A further consideration to be added here is this: When the action of an agar culture of the above-named six species is compared with that of the peritoneal exudation even of the first guinea-pig in the series, it will be found that the latter fluid is much more virulent, and considerably smaller doses produce a fatal result than in the case of the former. The explanation, I presume, is this: In the case of the agar culture the action is chiefly that of the intracellular poison, while in the case of the peritoneal exudation there is in addition the action of the ptomaines or toxins that had been elaborated by the bacilli in the peritoneal fluid. For the same reason also the fatal dose of the living agar culture injected into the peritoneal cavity is smaller than that of the sterilised culture as has been stated above, for in the latter the effect produced is only that of the intracellular poison injected, whereas in the former must be added also the action of the poisons elaborated in the peritoneal fluid by the living bacilli.

That there exists a notable difference between the intracellular poison and the toxins elaborated by the bacilli in the culture medium is strikingly shown by the fact that protected guinea-pigs which have survived the intraperitoneal injection of fatal doses of Haffkine's "virus fort," mentioned on former pages, when subsequently inoculated peritoneally with gelatine culture (liquefied by the growth) of the cholera bacilli promptly succumb, thus proving that, though pro-

² The similarity of the results obtained by Roux and Rodet after intraperitoneal injection into guinea-pigs of cultures of bacillus coli and bacillus of typhoid fever do not therefore prove anything as to the relationship of these two species, since the identical results are obtained with the proteus, bacillus prodigiosus, cholera vibrio, and vibrio of Finkler.

tected against the intracellular poison, they are by no means protected against the toxins elaborated in the culture medium (gelatine) by the bacilli.

11. It follows from all this that if in any case a refractory condition against the intracellular poison is established it does not follow that this refractory condition necessarily holds good also against the specific toxins or against the growth and multiplication of the bacilli and the production by them of their specific toxins in that refractory body. As we have shown, the previous injection of the intracellular poison of the vibrio of Finkler, of the bacillus prodigiosus, or of the bacillus coli, is capable of rendering the body refractory against the intracellular poison of the cholera bacillus or of the typhoid bacillus, but it would not be justifiable to assume that such a refractory condition is an index of insusceptibility against natural infection with cholera or typhoid fever; as a matter of fact, guinea-pigs protected against the intracellular poison of the cholera bacilli succumb to the toxins produced by these in the culture medium.

FUNCTIONAL OPHTHALMOPLÉGIA WITH GENERAL PARALYSIS AND IMPLICATION OF CRANIAL NERVES IN YOUNG WOMEN.

By C. W. SUCKLING, M.D. LOND., M.R.C.P.,
Professor of Medicine, Mason College, Birmingham.

I HAVE recently met with two cases in young women of ophthalmoplegia with general motor weakness and other paralytic symptoms, which are very similar to two cases recorded by Dr. Bristowe, and which I believe illustrate a distinct type of disease allied closely to exophthalmic goitre. The affection is probably functional, and in my opinion is decidedly not hysterical. In my two cases exhaustion seems to have been the exciting cause, and in each case there is a neurotic family history.

One patient is 18 years of age and very tall, the other 21 years and tall also. Both came to me complaining of general weakness. One patient is now quite unable to stand, her knees giving way under her, and for some months past she has fallen down at times; the other patient can only walk about a hundred yards and is then exhausted. Both patients are unable to hold their arms out or to place their hands on their heads, complaining that their upper limbs are very heavy. In both the movements of the eyeballs are much restricted and attended with nystagmus, there being ophthalmoplegia externa but no implications of the internal muscles. In both there is diplopia and squinting at times.

In both cases there is difficulty of articulation after the patient has been talking a little time, articulation then becoming an effort and the words slurred. There is no aphasia, motor or sensory. In both cases there is some difficulty of swallowing, and in one case after mastication the lower jaw drops for a time. In both cases mastication is difficult. In neither case has there been headache, vomiting, epileptic attacks, febrile attacks, nor exophthalmic goitre. In one there is slight chorea present. Both patients have been ailing about three years, the illness in each commencing gradually and slowly increasing, the prominent symptom being general weakness. In neither case is there any loss or perversion of sensation or affection of the special senses. The reflexes are the same in each case, the knee reflex being a little increased, but there is no ankle clonus. In one patient the hands and feet are always cold and blue, and the plantar reflex, though at times present, is generally absent. The fundus oculi is normal in each case. In both cases there is no disease of any thoracic or abdominal origin, and no urinary affection. In both there is complete command over the bladder and rectum. In neither case has there been any symptom of hysteria nor any mental peculiarity.

In one girl there have been suffocative attacks, the recurrence of which she greatly dreads. When I attempted to examine her larynx an attack was brought on by my slightly pushing her head backwards. She became cyanosed, and inspiratory stridor was well marked. While she is resting breath-

ing is quiet and easy, but any effort causes slight inspiratory stridor. She is always propped up in bed, and cannot lie down, dreading the difficulty of breathing which is apt to ensue if she does so. Although I have not seen the vocal cords in this case, I feel sure that there is paralysis of the abductor muscles, and I consider the patient to be in great danger. I am impressed with the relationship of exophthalmic goitre to this affection. At the present time I have two cases of exophthalmic goitre under my care, in one of which there is ophthalmoplegia externa and in the other paraplegia. I met with one of these cases of ophthalmoplegia some months before the other, and did not at all understand the case till I met with the second one. I showed both patients before meetings of the Pathological and Clinical section of the local branch of the British Medical Association a few weeks ago.

These cases have not exactly the symptoms which were observed by Dr. Bristowe in his, but they are evidently of the same nature, and, in my opinion, prove that there is a functional disease which chiefly affects young women, and which is characterised by ophthalmoplegia, more or less complete, with other motor symptoms, such as hemiplegia, general paralysis, dysphagia, dysarthria, etc., and that the affection is chronic and closely allied to exophthalmic goitre. Recovery may take place, or the patient may die from paralysis of the abductor muscles of the vocal cords or from some intercurrent disease, for example, bronchitis or pneumonia. The affection is probably one of the motor nuclei in the iter and floor of the fourth ventricle, and the general motor paresis is probably due to extension of the disease into the motor tracts. In both my cases the nuclei of the third, motor nuclei of the fifth, nuclei of the facial (slightly) spinal accessory, and hypoglossal nerves were probably affected. In both cases also the shoulder muscles were most affected, the nuclei for the movements of the shoulders being the highest. A lesion spreading downwards along the motor nuclei would involve these first. There were no electrical alterations in either case.

Dr. Bristowe, in his clinical lectures on diseases of the nervous system, describes cases of what he calls functional ophthalmoplegia. Three cases are described, one of which proved fatal from bronchitis, in which the necropsy revealed no lesion of the medulla or pons, but the condition of the nuclei of the third nerves was not made out. One case recovered, and the other improved. Two of the cases occurred in young women, the third in a man. All the cases were chronic in duration, lasting two years or more. In two of the cases right hemiplegia occurred with epileptic fits at intervals, the fits being preceded by a rise of temperature; in one of these cases there was hemianæsthesia. In the case of the man there was ophthalmoplegia externa and interna, partial anæsthesia of the head, neck, and chest; epileptic fits and attacks of intense dyspnoea dependent upon paralysis of the abductors of the vocal cords. The case of the man was probably dependent upon degenerative changes induced by syphilis.

The cases of the two women much resemble the two cases which I record. In one of Dr. Bristowe's cases exophthalmic goitre had preceded the ophthalmoplegia, in the other chorea developed during the progress of the illness.

Dr. Bristowe did not consider his cases to be hysterical for the following reasons:

(1) The gradual and uniform progress of the symptoms from bad to worse, there was never any variability, never any shifting of paralysis or anæsthesia; whatever fresh symptoms accrued were permanent.

(2) The character of the fits, which were clearly epileptic.

(3) The remarkable prevalence of febrile temperatures without any obvious cause.

Dr. Bristowe is of opinion that the disease was functional, affecting the floor of the fourth ventricle and walls of the iter, with extension into the neighbouring sensory and possibly even neighbouring motor tracts.

CASE 1.—A young woman, aged 21, came to me complaining of extreme general weakness. She could scarcely stand, and could not raise her arm. Her head dropped forwards and she could not keep it erect. A brother suffers from epilepsy, and a sister had paralysis. Her illness began about three years ago, with weakness of her arms, so that she could not arrange her hair, and had to prop her arms up to do so. She attributed her illness to overwork, having served behind a bar in an hotel for several years for many hours a day, constantly using her arms and stand-