RUPTURED TUBAL PREGNANCY: ABDOMINAL SECTION: REMOVAL OF FŒTUS AND GESTATION CYST: RECOVERY.¹

By JOHN W. BYERS, M.A., M.D., M.Ch.,

Physician for Diseases of Women to the Royal Hospital, Belfast, and Physician to the Belfast Hospital for Sick Children; Examiner in Obstetric Medicine, Gynaecology, and in Diseases of Children to the Royal University of Ireland.

ON September 30th, 1892, I was aroused early in the morning to see a patient, with her medical attendant, Dr. Austin, of Belfast. On arriving, I found her in a state of profound collapse, pallid, with cold face and extremities, and with a rapid "fainting" pulse. She was quite clear, and complained of pain in the "pit of the stomach," and at this spot (epigastrium) there was tenderness on pressure. It was evident that there had been a severe internal hæmorrhage, and, although physical examination of the abdomen and pelvis was negative—except for the pain and tenderness on pressure in the epigastrium—I came to the conclusion that the bleeding was due to a ruptured extrauterine gestation, mainly from the previous history, which had been most carefully noted by Dr. Austin.

It seemed that the patient, who was 33 years of age, had been married eleven years, and had one child 10 years old. About four years ago she had a miscarriage; and in July, 1891, I had seen her, with Dr. Wilson, of Whiteabbey, when she was residing a short distance from Belfast. At this time she was in a state of collapse, and I thought the illness was due to a ruptured tubal pregnancy. On August 1st, 1892, she went to a seaside resort a week after her period, and two weeks later—that is, three weeks from the original period—she had a watery discharge which lasted three days, her usual illness being five days in duration. This ceased for two days, and then reappeared and continued for two weeks. No clots nor pieces of "flesh" came away, but her breasts were sore and somewhat enlarged and she thought she might be pregnant. She came home to Belfast on August 26th—sooner than she had intended owing to the loss still continuing—and placed herself under Dr. Austin's care. The discharge then ceased, and she became "unwell" at her usual time, a week after this; and on September 14th she got up, although still "ill," wishing to be about when her husband would come home, and while lying on a couch in a room on the ground floor in the evening she was suddenly seized with pain and fainted. At 7.30 p.m. on that evening (September 14th) Dr. Austin was at once summoned to see her, and on his entering the room she said: "I feel the same as I did last year, and I am going to have a miscarriage." She was put to bed, and although nervous, pale, and weak, she was conscious and not much collapsed. The pulse was good, and on physical examination there was nothing pointing to pregnancy or to abortion. She soon rallied, and on September 15th the "loss" had ceased. Next day she vomited green bile, and during the following week she had attacks of vomiting and of cramp in the bowels and was tender over the epigastric region. On September 19th Dr. Wilson, her original medical attendant, saw her with Dr. Austin. On September 22nd she became unwell, and for eight days a dark red discharge continued, but no decidua came away, and by September 28th she had considerably improved; the abdomen was small and flat, and she was so well that it was arranged she was to have been up the next day or the day following.

On September 30th, at 5 a.m., Dr. Austin was urgently summoned, and found her intensely collapsed, with all the signs of a severe internal hæmorrhage. It was on this morning that I saw her, and came to the conclusion that the profound collapse was due to the rupture of a tubal pregnancy. The diagnosis was based mainly on the previous history (as given by herself and by Dr. Austin), and on the fact that no other source could be discovered for the hæmorrhage, the physical signs being of no value as localising the seat of the lesion; although there was the circumstance, to which the patient herself frequently drew attention, that she had been ill much in the same way in July, 1891, when I had seen her with Dr. Wilson. She gradually recovered from the collapse,

and at 10 p.m. on September 30th, when Dr. Austin saw her, she had quite rallied, and was fairly strong; but a short time after his visit he was sent for to see her, and on arriving he found her again collapsed. She now for the first time complained of pain in the left iliac region, so acute that she could not bear being touched in that part. Dr. Austin stayed all night, and by 10 a.m. the next morning reaction had set in, although she was pale and weak. At 1 p.m. on that day (October 1st, 1892) I saw her for the second time, in consultation with Dr. Austin, and we both urged an operation; and it was arranged that at 5.30 p.m. this should be done, Professor Sinclair being asked to meet us at that time. However, at 4 p.m. a letter was received from the patient's husband to the effect that he had "decided not to go on with the operation." It seemed that the patient's relatives could not reconcile the necessity or advisability of an operation with the fact that in the previous year (July, 1891) she had recovered without surgical interference from a condition which she and they considered quite as alarming as that in which she now was.

I learned from Dr. Austin that on October 2nd, at 6.30 in the evening, she was seized with faintness and abdominal pains, and when he saw her at 9.30 p.m. he believed that bleeding had again been taking place. At this time she was very pale, decidedly worse, and was thought to be dying. The abdomen was distended and tympanic in front, but there was dullness in the flanks, with an obscure feeling of fluctuation. With the aid of hot bottles, champagne, and morphine she rallied, when at 2 a.m. Dr. Austin (who once more urged the absolute necessity of an operation) went home. On that morning the friends at last saw that her condition was hopeless unless something operative could be done, and I was again sum-

Fig. 1.—Fœtus, in amniotic cavity, removed from abdomen.

moned, but, owing to being engaged in another case, I did not see her until 7 p.m. in the evening of October 2nd, when Dr. Austin, Dr. Wilson, Professor Sinclair, and myself met in consultation. At this time she had rallied, and the pulse, although very quick and weak, was quite perceptible. She was extremely exsanguine, her skin of a pale lemon hue, and the abdomen, which was distended, evidently contained a quantity of fluid. There was tenderness in both iliac regions, but on pelvic examination the os uteri was closed, the uterus was movable and not enlarged, and no tumour could be felt.

On passing a fine trocar into the right flank the presence of blood was demonstrated, and her friends then consented to allow an operation to be done, and she was removed to the Royal Hospital, where, the next morning, assisted by Professor Sinclair, and Dr. Austin and Dr. Wilson being present, I opened the abdomen at 9 a.m. The parietal incision showed the abdominal walls to be very exsanguine, and on reaching the peritoneum that part was noticed to be of a dark purple hue. On opening the cavity an immense quantity of blood clot and serum welled out. The left tube—the seat of the gestation—was found ruptured, and it was removed after tying the broad ligament with the Staffordshire knot. The peritoneum was then flushed out with warm water, and after some clots had come away, the fetus surrounded with the amnion was washed out from amongst the intestines. After thoroughly flushing the abdomen, and clearing out in

¹ Read at a meeting of the North of Ireland Branch of the British Medical Association.

this way large quantities of blood and clots, a glass drainage tube was inserted, and the cavity closed. The patient made an uninterrupted recovery, with the exception of a small stitch abscess, and the drainage tube was removed on the fourth day. The accompanying plates, from photographs taken immediately after the operation, show the fœtus and the ruptured gestation tube, which were removed from the abdomen.

REMARKS.—The points worthy of remark in this case are in reference to the diagnosis and treatment.

1. *Diagnosis.*—When I first saw the case with Dr. Austin, although the symptoms were very obscure, I believed that it was one of ruptured tubal pregnancy. The problem to be solved then was: Given a married woman suddenly seized with profound collapse and symptoms of internal hæmorrhage, but with an absence of any special abdominal or pelvic

Fig. 2.—Left Fallopian tube, ruptured at seat of gestation cyst.

signs on physical examination, what was the source of the bleeding? Her previous history: Irregularity of menstruation, uterine hæmorrhage, and a period of sterility afforded the clue to the diagnosis of a tubal gestation, which had ruptured, and the subsequent attacks of pain and collapse with the abdominal signs confirmed this view of the case. The condition of matters present in July, 1891, when her relatives as well as myself thought her profoundly ill, is difficult to explain. At that time her symptoms may have depended on a ruptured tubal pregnancy, from which she recovered, or on an ordinary or a tubal abortion.

2. *Treatment.*—The line of treatment indicates the enormous advance made in recent years in abdominal surgery; and the case is of special interest as being the first one operated on in Belfast. The patient is now quite well.

DONATIONS.—The Fishmongers' Company have made a grant of £150 towards the funds of the Metropolitan Hospital, Kingsland Road.

THE PREVENTION AND MITIGATION OF SMALL-POX.

By SIR PETER EADE, M.D., F.R.C.P.,
Consulting Physician to the Norfolk and Norwich Hospital.

THE increasing prevalence and diffusion of small-pox in England at this time necessarily attract the attention of the medical profession to the subject of this disease; the more especially as it is well known that there is a large and increasing number of unvaccinated, and therefore unprotected, children and persons in our midst.

As in the case of the other acute zymotic diseases, we know of no internal remedy which can cut short the disease. Nor can we do more by way of treatment than relieve some of the more urgent symptoms and miseries which attend its course.

We all watch with interest the vital experiment which Leicester, as well as some other places, is making in regard to variola, and we mark the repetition of the known facts of the disease which are there being illustrated. We can scarcely doubt that the result will be—as is, indeed, already becoming the case—to show once more that a serious proportion of the unvaccinated die, and that there is but one real known remedy against an attack of small-pox, and that is the preventive protection of the population by universal vaccination.

But even if there were any reliable method of treating the disease when once acquired, "prevention is better than cure;" and the prevention of it, by efficient vaccination and revaccination at proper periods, is not only better, but a means of saving a large amount of human life. And the professional mind, as a whole, remains quite unshaken in the conviction of the general efficacy of these processes. This, then, is the first and the greatest of antivariolous preventive measures. But in the face of a present widespread epidemic, and with the possibility of very valuable lives at stake, it seems worth while to recall other measures more or less effective to prevent or mitigate which have from time to time been recommended, after exposure to the contagion or during the progress of the disease itself.

The view has been entertained, and there are many strong arguments in favour of it, that small-pox contagion is really and actually received upon the cutaneous and mucous surfaces, and that it grows and develops thereon, or rather in their follicles or involutions, prior to the infection of the blood and general system—just as in diphtheria we believe that the specific poison first alights upon the tonsils, and, vegetating there, produces secondarily the further local and general symptoms which characterise the disease.

Now, after exposure to variolous contagion there are twelve days before the seed sown begins to show its local reproductions, before the distinctive papular foci appear and develop into the typical papules and pustules. We cannot reach or treat any seed spots sown upon the internal mucous membranes, but has any sufficient trial ever been made during this incubatory period, and after exposure, of the method of inuncting the outer surface, and especially the more exposed parts, such as the face, neck, chest, and wrists, with some sufficiently potent antiseptic or bactericidal application? This might be either sulphur or carbolic ointment well rubbed into the pores of the skin of these parts, or they might be saturated with some cleaner application, such as strong salicylic lotion. We know that the danger of an attack of small-pox may be largely measured by the number of pustules which form. Therefore even a considerable diminution of these, or their early abortion, would be almost certain to benefit the patient and increase his chances of recovery.

Again, after the eruption has appeared and the papules have begun to soften, it has often been proposed to apply to these some bactericidal application which may prevent the normal and thorough development therein of the variolous germs. External spongings at this stage with lotions, or the outside smearing of the surfaces with antiseptic applications, however strong, although apparently of some value, are insufficient for abortive action, because the agents used do not come in contact with the ripening germs. But a large amount of benefit has been found to be derived from their insertion into the individual pimples, just as has been found from a similar internal application of bactericides to the