means of Loeffler's alkaline methylene blue (aqu. sol. dilute before use with an equal vol. water) the nerve cells and the connective tissue throughout the section can be stained, but the nerve tubes are poorly shown. Excess of dye is removed by the piridin which follows. Picrocarmine (half an hour) stains all the structures present; but I have hitherto got the most satisfactory results by staining first in anilin blue-black, of the strength given, for 20 minutes, and subsequently in picrocarmine for 15 to 30 minutes. The staining of the grey matter will, I hope, be presently improved upon. My immediate object is more especially to bring to notice the fact that by means of piridin sections of the fresh cord can be fixed and preserved unwrinkled, each nerve tube being well defined. Sections prepared and mounted in the manner described show scarcely any diminution in size when compared with the original cord. The rapidity of the method is noteworthy, for by it the cord can be removed, cut, stained, and mounted in the course of a day; indeed, it is probable that from 2 to 4 hours would suffice.

In conclusion, I may state that I have hitherto practised with the cord of the cow, cat, and rabbit respectively, in each case successfully, with the reservation regarding the grey matter alluded to. The longest period which has elapsed between the death of the animal and the time of cutting sections (after which the elements were thoroughly fixed) is an hour and a-half (cow's cord). I have not yet cut any later than this.

Judging from certain specimens in which nerve roots appear outside the cord in transverse section, I am disposed to think that the same method may be advantageously employed for peripheral nerves.

HYPERTROPHIC ELONGATION OF THE CERVIX AS A CAUSE OF OBSTRUCTED LABOUR.¹

By SAMUEL GRIFFITH, M.D.,
Portmadoc.

HYPERTROPHY and elongation of the cervix and os uteri, or hypertrophic elongation of the cervix, is a very uncommon cause of obstructed labour. In an experience of forty years I have met with only the two following cases.

CASE I.—On November 17th, 1888, I was called to visit Mrs. W., stated to be in labour of her second child. I found her complaining of very severe but irregular pains in the back. The os uteri was between the vulvæ, looking excessively swollen and livid, with a posterior and anterior slit at least an inch deep in the cervix; the cervix was also very hypertrophic, and was elongated to the extent of three inches. Both the external and internal os, as well as the cervical canal, were so patulous that I had no difficulty in inserting my finger and feeling the presenting part above the brim. There was no bulging of the membranes with the pains, so I left the house, prescribing rest, opiates, and purgatives. I was not summoned again until November 24th, and as I was not at home when the messenger came, my assistant went there and remained with the patient all night. On November 25th, about midday, he sent for me, stating that the patient was making no progress, that she had been having pains all night, but they were not at all strong; that shortly before he sent they had ceased altogether, and that the patient looked very weak. When I arrived it was quite evident that rupture of the uterus had taken place.

Tait's modification of Porro's operation having just been made known, I was very anxious to perform it, and for that purpose sent for my friend and former pupil, Dr. Jones-Morris, to assist me. The husband, however, refused his consent unless we could guarantee that the operation would be successful; this, of course, we were not able to do, but we assured him it was the only chance for the poor woman. She died in about fifteen hours after the rupture had taken place.

I was allowed to make a *post-mortem* examination. The rupture was a transverse one, and exactly on a level with the peritoneal surface of the roof of the vagina—that is, imme-

diately above the internal os; there had been very little hæmorrhage, and evidently death was caused by shock.

CASE II.—In August, 1892, Dr. Jones-Morris requested me to see a multipara with him, as the case resembled very much the one he had seen with me. There was, however, good dilatation, and he had applied the long forceps, but could make no headway. The pulse was fairly good, but the pains weak. We applied the forceps again, but as soon as we did so the pains stopped altogether; we had, therefore, to remove them, and wait for pains to return. This was on Sunday evening, the pains did not return with any force until Monday evening; we then applied the forceps again, and delivered her of a stillborn child after a very tedious and difficult operation; metritis set in, and in five or six days the patient succumbed.

This complication is so rare that in most of our obstetric literature it is not even hinted at, but in Spiegelberg's *Textbook of Midwifery* "prolongation of the cervix, especially of its intravaginal portion," is just mentioned. It is stated that it must not be confounded with prolapse, that the condition is unimportant, and causes no trouble during labour, except, perhaps, by a delayed softening of the lips of the os, interfering with its dilatation. This, however, is wrong, as it is clear that hypertrophic elongation of the cervix may lead to very serious difficulties in delivery. Professor Simpson² describes in the Edinburgh Obstetrical Society's *Transactions* a case in which "it became necessary for him to break up the head of the child in a woman in whom the cervix was between four and five inches long, and correspondingly thickened; and he mentions other cases where lateral incisions nearly two inches deep had to be made through the cervix before delivery could be effected."

MEMORANDA: MEDICAL, SURGICAL, OBSTETRICAL, THERAPEUTICAL, PATHOLOGICAL, Etc.

PSOAS ABSCESS TREATED BY REPEATED ASPIRATION AND INJECTION OF IODOFORM EMULSION.

H. B., aged 20, in August, 1891, complained of weakness and lassitude, which had come on after two severe attacks of influenza. In October, 1891, he complained of aching in the lumbar region, but careful examination showed nothing to indicate spinal mischief. In a few weeks the pain disappeared; but in January, 1892, he came again with a swelling in the right groin, which proved to be a large psoas abscess; there was no deformity or tenderness of the spine. In fact, the patient had been daily following a somewhat laborious occupation. The apices of both lungs showed impaired resonance and marked crepitations.

Treatment.—It was decided to try a method of aspiration recommended by Mr. A. W. Mayo Robson in the *BRITISH MEDICAL JOURNAL* of April 25th, 1891. On January 26th the first aspiration was performed, with antiseptic precautions, and 24 ounces of pus drawn off. Aspiration was stopped immediately the pus became tinged with blood. Pads were applied above and below Poupart's ligament, then cotton wool and a bandage, and over all elastic webbing, so as to get considerable elastic pressure. The patient was kept perfectly at rest on his back in bed, and disturbed as little as possible, whilst cod-liver oil and as much nourishing food as possible were given. The temperature, previously hectic, became at once almost normal, with a variation of 1° between morning and evening. In two weeks the temperature became hectic again (a variation of 3° between morning and evening), and the abscess cavity was found to be refilling. On February 15th it was again aspirated, 17 ounces of pus drawn off, and 1 ounce of iodoform emulsion injected. The after-treatment was the same as before.

On March 7th the cavity was again aspirated; 11 ounces of pus drawn off, and 1½ ounce of iodoform emulsion injected. The temperature again improved for a time, and then resumed the hectic type.

On April 5th aspiration was performed for the fourth time,

¹ Abstract of a paper read at a meeting of the North Wales Branch of the British Medical Association.

² Quoted in Lawson Tait's *Diseases of Women and Abdominal Surgery*.