(d) Excitation of the same convolution higher up, at 4, evoked movements of the wrist and some twitching of the fingers.

It will be seen that excitation at 1 evoked a movement of the head towards the stimulated side; this movement was similar in character to that which was observed in the epileptiform convulsion, and it is therefore probable that this kind of movement was represented in this area of the cortex. In addition to this result, which, being the opposite to that ascertained by experiment on the monkey, is itself noteworthy, there are two other circumstances to which I wish to draw attention.

The first is the great inexcitability of the area marked 2. A similar inexcitable area has been previously noted by Beevor and Horsley in the orang, and has been observed also in the few recorded instances in which this particular region has been stimulated in man. Its significance, lying as it does between the lower limit of the excitable area of the upper limb and the upper limit of that of the head and face, has been remarked on by the authors just referred to. The present result confirms the previous ones.

The second circumstance is one of much importance physiologically, and is connected with the enormous intensity of the stimulus which it is necessary to employ in the case of the more excitable areas. Here, too, the orang experiments pointed to the existence of a similar necessity. It was found that a series of far more intense induction currents had to be used to evoke movements from the cortex of this animal than was necessary in the case of such smaller and lower types of monkey as the common macaque. Further, the cases in which the human cortex has been excited seem to exhibit the same remarkable feature. In comparison with the strength of stimulus which is found adequate to excite most of the excitable tissues, that now employed was astonishingly great.

Two factors seem to especially determine the relation between strength of exciting induction currents and the production of nerve impulses down motor nerve paths of sufficient power to evoke muscular contractions. There are first the physical alterations in the distribution of the currents due to the mass, moisture, temperature, etc., of the tissue which these traverse, and secondly, the physiological resistance of tissue to the arousing agent. The first factor is undoubtedly largely dependent upon the mass of the tissue, especially when this is a moist conductor, and it is possible that the increased spread of current deviations through such a large bulk as that of the human brain may, to some extent, explain the high intensity which has to be used. It is, however, far more probable that the physiological factor is the more potent. The more highly differentiated cortex seems to possess greater inertia than the less highly focalised one; there is a greater disinclination in such a cortex to pass from the static to the dynamic condition. The contrast between the ease with which epileptiform attacks are evoked by artificial stimulation of the cat's brain, and the difficulty of similarly evoking such convulsions from the orang, is an indication of this. But it must not be forgotten that there is a third possibility. In the case of the spinal cord there is abundant experimental evidence to show that a discharge of impulses from its motor side is far more easily evoked by electrical or mechanical irritation of the posterior roots or posterior columns than by similar irritation of the grey matter.

A similar condition may rule in the sensory motor region of the cerebral cortex. Indeed, it is possible that this may be only capable of responding reflexly, that is, to the stimulation of its entering afferent nerves. The comparative thickness of the human cortex would thus necessitate the employment of a current of very considerable intensity, since only under these circumstances would a derivation of adequate strength traverse and excite a sufficient number of the deeply-situated afferent bundles of nerve fibres.

THE AFTER-EFFECTS OF ABLATION.

There is little to be said on this side of the question. The area ablated is indicated in the diagram, and the results to which attention is here directed may be briefly summarised as follows:

1. The temporary loss of voluntary power in the grasp of the left hand was associated with complete inability to com-

pletely extend the thumb, and with a very marked diminution in the number of fits.

2. A restitution of power in hand and thumb occurred in a very short time (a week to ten days), and, coincident with this restitution, was a renewal of the fits at more frequent intervals.

3. There was no impairment of sensibility, but on this point the method of testing employed soon after the operation was not a satisfactory one. There was, however, a prolonged subsidence of the sensory aura (tingling, etc.), which does not appear to have been permanently lost, since the boy has informed Mr. Parker and myself that quite recently, in some of the few fits which he still has, the aura has been present again for the first time since the ablation.

A CLINICAL STUDY OF THE CAUSATION AND TREATMENT OF CERVICAL DYSMENORRHOEA.

By M. HANDFIELD-JONES, M.D. LOND.,

Senior Obstetric Physician to St. Mary's Hospital, and Lecturer on Midwifery and Diseases of Women to St. Mary's Medical School.

MANY theories, little knowledge. This saying undoubtedly applies to the subject of dysmenorrhœa, for numerous though the theories are which have been put forward to explain and account for severe suffering in connection with menstruation, yet it cannot be said that much exact and scientific knowledge of the subject exists at present. Doubtless in such a complex process as menstruation, possessing as it does numerous local and constitutional phenomena, and depending as it does on a healthy anatomical tract, on a normal blood condition, and on perfect nervous function, there are abundant opportunities for disordered working of some of the component parts. In the search for truth it would appear that some investigators, close students of the anatomical tract, have unduly magnified certain mechanical obstructions, and made overmuch of one possible factor; while, on the other hand, some who longed for a severe simplicity of explanation, have found in spasm the "fons et origo" of all the menstrual suffering which woman has endured since our primeval mother had her first monthly sickness. In the present communication I have no wish to be dogmatic, nor do I desire to lay stress on any one theory to the exclusion of all others; but having gone over the clinical records of all the cases of dysmenorrhœa which have occurred in my private and hospital practice during the last five years, I shall endeavour to set down the deductions I have drawn from a study of these cases. To avoid possible misunderstanding, the term dysmenorrhœa is used throughout this paper to describe pelvic pain, which occurs during the menstrual bleeding or during the two or three days which immediately precede the onset of the flow.

NEURALGIC DYSMENORRHOEA.

The first group of cases comes under this heading, and a single one will serve as an example:

Miss E. B., aged 19, is a well-developed girl, healthy, but not very robust. She menstruated first at 13, and has her periods with fair regularity. The periods last as a rule six days; no clots are passed; for the first three years there was no pain at the times. During the last three years she has had pain with the periods; usually at its worst on the first day. Now the interesting feature about this pain is this: The girl is engaged in domestic service, and spends part of the year in London and part by the seaside. When she has been in London some weeks, has been up late, and is somewhat overworked, then she has menstrual pain, but she loses the latter entirely on returning to the country house, where work is light and exercise in the fresh air constant. Naturally no vaginal examination has been made in this case, nor indeed in any which occur in this group, but it is impossible to believe that pain which can thus come and go is of other than neuralgic origin.

INFLAMMATORY DYSMENORRHOEA.

The majority of the cases which come under this heading occurred in women who had passed, more or less recently, through an attack of pelvic peritonitis or cellulitis. The disease in some cases had been partial and localised to one region of the pelvis; in others it had been general, involving the whole of the pelvic peritoneum or cellular tissue. In the majority of the patients menstruation had always occurred before the inflammatory attack, with little or no suffering; in all after the disease dysmenorrhœa had been a prominent

symptom. It is easier to account for the return of the pain each month in these cases than in any other. Imagine a woman with cellulitis of the hand and forearm who is lying in bed with the arm raised on a pillow; now let such a patient hang the arm over the side of the bed for a considerable time. Soon the weakened blood vessels are gorged with blood, the recently-inflamed nerve endings are again irritated, the arm begins to throb and burn again, and the disease is, to an extent, restarted. This is exactly what occurs in the pelvis every month for some time after an attack of pelvic peritonitis or cellulitis, for with the rise of vascular tension and engorgement of the pelvic blood vessels absence of pain must be an impossibility. The liability to return of the inflammatory condition at each monthly period, unless care is taken, and the existence of the term "recurrent menstrual peritonitis" are evidences of the truth of this pathology. In inflammatory dysmenorrhœa the pain begins constantly two or three days before the commencement of the blood loss, and increases in degree as the onset of menstruation approaches. It is commonly remarked by the patient that the more she loses the greater is the relief experienced.

MEMBRANOUS DYSMENORRHOEA.

Regarding this form I do not intend to say anything, for my cases are too few to draw any useful deductions from, and the subject is too complex and intricate to be treated shortly. In his recent Harveian Lectures Dr. Champneys has compiled the work done by various authors on this question so fully and so ably that everyone interested in the phenomena of this form of dysmenorrhœa should carefully study those writings.

Having selected and classified the cases thus far, there remains now a large group which cannot be referred to either of the foregoing divisions, and in which the causation of the pain is not so easily capable of explanation. This is the group to which the terms "mechanical," "spasmodic," "obstructive" have been applied by some writers, the group on which most difference of opinion rests, and to the study of which this paper is especially directed. The terms "mechanical" and "obstructive" are correct in a certain sense, but not, I imagine, in the sense in which they have been applied by their authors. It is probably true that the uterine contractions required to expel clots or membranes may cause suffering; but then the same suffering is felt during the expulsion of the fetus in normal labour, and yet we do not talk of mechanical or obstructive labour pains. The term "spasmodic" is more correct if it be used to imply uterine contractions painful because excessive; but if it means that the uterine contractions of menstruation are abnormal, partial, and unconsentaneous, then it implies a condition the existence of which is purely hypothetical, and the proof of which must rest merely on inference. The term "cervical dysmenorrhœa" is used in this communication because it seems capable of proof that the causation of menstrual pain in the cases now under consideration is allied with certain abnormal conditions which have their seat principally at and near the internal os. Before proceeding to quote cases in proof of this assertion, it will be well to inquire whether we can show or bring forward any clinical evidence to prove that uterine contractions do normally occur during the menstrual epoch; what is the effect of these contractions, if they do occur, on the uterine canal; and the relation of these contractions to pain.

DOES THE UTERUS CONTRACT IN CONNECTION WITH MENSTRUATION?

Menstruation has often been described as a miniature labour, and the analogy between the phenomena of menstruation and those of labour has been carefully worked out. If this comparison is worthy of belief, it follows that at every menstrual epoch intermittent contractions become established in the uterus, and dilatation of the cervical canal either wholly or in part occurs. Following out the comparison with labour still farther, it is clear that the internal os will be the part first affected by this dilatation. Long ago Dr. Braxton Hicks and other authorities expressed their belief that rhythmical uterine contractions did occur during normal menstruation, but I am not aware that anyone has demonstrated the effects of these contractions on the cervical canal. From the very nature of the case it is clear that any

such demonstration must rest solely on clinical evidence, for there is no reasonable probability that pathological experiments or *post-mortem* examinations will throw much light on the subject. The two following clinical records will supply evidence regarding the condition of the "internal os" during menstruation.

CASE I.—In December, 1889, I was asked to see Miss X., aged 33, who had been suffering for some months from very profuse menstruation. Her medical man had examined *per vaginam* on several occasions, and had given an opinion that some intrauterine growth was present. The patient had been menstruating three days when I saw her, and was still losing heavily; but as I had to return to town and an immediate opinion was requested, I examined by the vagina and found the following condition: The external os was dilated to the size of a threepenny piece, the cervical canal was almost obliterated, and a tumour the size of a very small hen's egg was presenting with its lower end almost resting on the external os. Gradually insinuating my finger through the yielding external os, I could explore the uterine cavity and make out that the tumour was attached high up to the fundus. I could not make out any trace of the internal os. About a fortnight later I had the opportunity of examining again, menstruation then having ceased about ten days, and found now that the condition of the parts had greatly altered. The external os was indeed but little changed, but the cervical canal was of normal length, though patulous in its lower part, and the internal os was so tightly closed that it was impossible to introduce the finger through it in order to explore the uterine cavity. On several occasions afterwards I had the opportunity of repeating these observations, and I found that while the internal os was always more or less firmly closed during the intermenstrual epochs, yet at the time of menstruation the inner os was freely patulous and admitted the examining finger. The notes of a case in which precisely similar phenomena were observed have been supplied to me by Dr. Travers.

CASE II.—Miss L., aged about 28, was brought to me complaining of great suffering in connection with menstruation. She had grumbling pain in both ovarian regions for two days before the onset of the monthly discharge, but the first day was her time of greatest suffering; lately the loss had become more profuse, and was accompanied by the passage of clots. Various medicines, douches, and vaginal supports had been tried, but had failed to relieve the pain, and each month the suffering was becoming greater. Vaginal examination showed that the uterus was slightly anteverted, and that its tissue was soft and flabby; in other respects everything seemed normal. As other means had been tried and failed, the patient was advised to submit to dilatation of the cervical canal without further delay. In dilating, my usual plan is to give ether and dilate up to 11 or 12 Hegar at one sitting, and then keep the patient in bed for five or six days; in the present case, however, this plan was not feasible, as the patient declined to desist from work and lie up. It was therefore resolved to dilate gently with the graduated steel dilators on two occasions in each month, and this plan proved entirely successful. The following point was noticed in the course of this treatment: in the few days which followed the close of a period it was easy to pass the dilators up to No. 9, but in the week which preceded a period it was impossible sometimes to introduce No. 4, and never was it done without much pain and difficulty. At all times it was easy to pass the dilators up to the level of the internal os, but it was at this point that the sounds were arrested. The increase of difficulty preceding menstruation was too pronounced to be dependent on any swollen condition of mucous membrane or turgescence of the tissues from hyperæmia, even if it could be shown that such conditions were actually present.

One more piece of evidence I may be permitted to quote in support of this view, that the internal os is drawn open by the uterine contractions during the onset of menstruation. In his work on *Diseases of Women*, the late Dr. Matthews Duncan says:

In the various cases that have been in "Martha" within this last year, we have found no stricture, no contraction of the passage through the womb, except in one case. Especially did we find no contraction when the woman was suffering from the pain; for in order to satisfy ourselves as to the nature of the disease in some of the cases, we passed a bougie into the womb while the woman was in the agonies of dysmenorrhœa, and we found that the passage was clear.

Though the proof is not absolute, yet the inference to be deduced from the above quotation is plainly this: that the result of acute pains (due, of course, to uterine contractions) is to render the cervical canal especially clear, or, in other words, to dilate the internal os. Though the evidence may not be as yet very strong, still, reasoning from the allied phenomena of labour, and having regard to the clinical facts already brought forward, it seems justifiable to take the three following propositions as proven: (1) that rhythmical uterine contractions do occur during the early part of the menstrual process; (2) that inhibition of the cervical or lower uterine sphincter fibres does exist; and (3) that, as a result of the foregoing, dilatation of the cervical canal does take place.

WHEN DOES THIS DILATATION COMMENCE, AND WHEN IS IT COMPLETED?

Our present knowledge of the phenomena of menstruation is so limited and still so theoretical that it is impossible to answer these questions with accuracy. If, however, we fall back once more on the analogy to labour, and regard the first day of menstruation as representing the period of an ordinary labour, as we understand the term commonly, then we should

expect the dilating process to commence some hours before the onset of the flow, and to be completed before the close of the first day. Of course, it is assumed that in such a case the conditions are all quite healthy; for it is clear that in cases where the tissues have undergone various pathological changes, or in which thickened membrane has to be thrown off, the dilatation might be much delayed, might require to be carried to a greater degree, and the various stages would be prolonged and accompanied with pain of varying intensity and duration.

IS THIS DILATATION NORMALLY ATTENDED WITH PAIN?

Having regard to the previous arguments, one might jump at once to the conclusion that the menstrual act must at all times, and even in perfectly healthy women, be attended with suffering. It is undoubtedly true that few women pass through the monthly sickness without more or less discomfort, but a little consideration will show that, under normal conditions, dilatation of the cervical canal to the amount attendant on menstruation need not be, and is not, attended with pain. Anyone who has examined healthy women two or three days before the onset of labour pains must have noticed how often the internal os has been taken up into the uterine cavity, how the cervical canal has disappeared, and the external os has been opened up to the size of a sixpence, and yet all this has been effected without the woman experiencing any discomfort. Again, numerous instances are on record of children born during sleep, or of no discomfort being felt until the head had entered the pelvis; Brunon describes such a case in the *Archives de Tocologie* for November, 1890. Few men have been long in practice without having seen cases of primiparae in whom the whole of the first stage has been completed with little more than slight discomfort, and in whom real pains only commenced with the onset of expulsion. One such case I may quote as an example of painless first stage:

Mrs. G., primipara, aged 24, noticed, while driving about one afternoon in her carriage, she being at that time a fortnight from her full term, that some discharge was coming away from her. On arriving at home she found her linen stained with blood, so sent at once for assistance. Before medical help was obtained pains had commenced sharply, but within ten minutes of their commencement the os was found to be fully dilated, and the head in the cavity of the pelvis.

There is evidence also supplied by those who have had the opportunity of observing the act of parturition among South African tribes, which leads to the belief that among these savage women the first stage is commonly a painless one, and that suffering attends the expulsive stage alone. If this be true in labour, when the cervical canal has to undergo complete dilatation, it is clearly probable, in the highest degree, that the comparatively small amount of dilatation required during the menstrual process should be obtained without making itself felt by the presence of pain, provided always that the woman is healthy, and that the condition of the various uterine tissues is normal. We are brought now face to face with this question: If granted that in normal menstruation dilatation of the internal os and cervical canal does occur, and usually does occur without the setting up of pain, what conditions may and do exist which will interfere with the process of dilatation, and so lead to the establishment of pain or, in other words, to dysmenorrhœa?

In attempting to answer this question, reference may be made again to the process of labour at term, and a comparison may be instituted between the causes which delay the completion of the first stage of labour and the causes of dysmenorrhœa.

A. MALPOSITION OF UTERUS.

It is a fact well known to accoucheurs that when the full term uterus is allowed by a flaccid condition of the abdominal walls to drop forward into a position of acute ante flexion, then the first stage is indefinitely prolonged, the os does not open up, and the suffering is acute. As soon, however, as the uterus is pushed back and maintained *in situ* by a binder, then the pains improve, the os quickly opens up, and the first stage may be completed rapidly. In the vicious position of the uterus the longitudinal fibres of that organ cannot act at advantage, and dilatation is delayed; moreover, suffering is greatly aggravated. What is true of the pregnant uterus is equally true of the non-pregnant. The two following clinical records well illustrate this.

Miss E., aged 26, was sent to me by Dr. Arthur Rowe, of Margate, for the treatment of acute suffering, which usually commenced some hours before the onset of her period, and lasted into the second day. The dysmenorrhœa had been present for some years. Ether was administered, and, an acute retroflexion having been recognised, a Hodge pessary was inserted. The next period came on with very little pain, and now, six months later, all acute pain has disappeared, and the monthly time is no longer dreaded.

Miss C., aged 28, has for a long time suffered severe pain during the first day, or sometimes two days of her menstruation. Medicines have failed to relieve. Vaginal examination reveals the presence of an acute retroflexion. A Hodge was inserted, and the uterus maintained in good position. The next period was nearly painless, and the following one entirely so.

Of course minor degrees of uterine displacement exist without the presence of menstrual suffering, as may be noted in cases in which the abnormal position is not sufficiently pronounced to affect the action of the muscular fibres of the womb injuriously; and, again, menstrual suffering may exist with slight abnormalities of position, and yet be really dependent on other causes. In the non-gravid condition retroflexions and retroversions will necessarily be more commonly a cause of dysmenorrhœa than anterior displacements, for owing to anatomical conditions the former will be usually more pronounced than the latter. The readiness with which relief from pain was obtained in the two cases quoted above, as soon as the displaced organ was raised and a support adjusted, is a strong argument in favour of the explanation put forward. Of course prolapse is sometimes associated with retroflexion or retroversion, and then reposition relieves the dragging on the broad ligaments and lessens venous congestion of the uterus, but in the instances already given this complication was certainly absent, and may be safely discarded.

B. MUSCULAR SPASM OF THE SPHINCTER FIBRES OF THE INTERNAL OS.

There are two drugs which relieve pain principally by relaxing spasm, namely, chloral and belladonna; the action of the former in cervical spasm of the first stage is well known to obstetricians. Both these drugs act admirably in some cases of dysmenorrhœa, as the following case will show:

Mrs. G., aged 24, began to menstruate at 16, in regular monthly, loss free, and continues four days. She has always had on the first day severe pain accompanied by vomiting and faintness; the pain lasts about ten hours. She has been married nine months, and her menstrual suffering has increased since marriage. By vaginal examination the womb appears quite normal. Full doses of belladonna taken immediately the pain of the next period commenced brought almost complete relief, and the same happy effect was brought about in the second period. After this she became pregnant. There can be little doubt that the belladonna brought about relief by relaxing a muscular spasm of the cervical sphincter fibres.

Probably this cause of dysmenorrhœa is not very common, but that it does occasionally occur there can be little doubt. Dilatation is not satisfactory in these cases as a means of cure, for the relief gained is very temporary, and the effect of the dilatation is soon lost. In passing Hegar's dilators on these subjects it is interesting to note how gradually the muscular resistance yields before the pressure of each dilator, and how quickly the internal os closes down again as each dilator is withdrawn and the next size larger is sought to be introduced. The sensation imparted to the operator's hand is very different to that felt when the resistance is due to a fibroid condition of the os internum.

C. FIBROID THICKENING AT THE LEVEL OF THE OS INTERNUM.

Histologists have pointed out that the connective tissue element is relatively more abundant in the cervix than in the body of the uterus, but under normal conditions there is no reason to suppose that this in any way interferes with the expansion of the os during labour. If, however, in any case it could be shown that the tissues of the upper part of the cervix were unduly dense and resistant, one might reasonably expect that dilatation would be delayed, and painful uterine contractions might be set up. Certainly, in cases of thick fibrosed cervixes depending on past chronic areolar hyperplasia, the first stage of labour is much prolonged, and the pains of this stage are much exaggerated. Still carrying out the comparison between menstruation and labour, one might reasonably expect that fibroid deposits or degeneration in the region of the internal os would lead to painful menstruation. An examination of the following cases will afford evidence in support of this view of the causation of dysmenorrhœa.

Miss G., aged 22, was admitted to St. Mary's Hospital, complaining of

great pain in connection with the monthly periods. Menstruation began at 13, is regular monthly, lasts seven days. Pain has always accompanied the process, but is becoming each year more intense. The suffering commences some hours before the onset of the flow, and lasts into the second day. Drugs have failed to relieve her. The patient was anaesthetised about a week before the onset of the period, and Hegar's dilators passed up to No. 12. The period which followed this dilatation was almost painless.

The most striking point in the course of the dilatation was the intense resistance offered by the tissues at the level of the internal os; it seemed as if one was endeavouring to stretch a ring of cartilage. To the hand accustomed to these dilating operations, it is easy to distinguish between the way in which a muscular spasm gives, and the dull tough resistance offered by a dense ring of fibroid tissue. Moreover the former quickly closes down when the dilator is withdrawn, while the latter, though yielding less readily, remains longer patent. In another case of a similar nature there were two distinct zones of fibroid stenosis at the level of the internal os about a quarter of an inch apart, and so dense and hard were these rings that it was necessary to nick them with a bistoury before dilatation could be satisfactorily completed. In both these cases the uterus was well developed, and the uterine sound passed 2½ inches. Space does not permit of the narration of more of these interesting cases, but a brief consideration of the two examples already given will enable us to see what a hindrance to dilatation of the internal os at the menstrual epoch is offered by the anatomical condition, and what a cause of menstrual suffering is necessarily present. To anyone who has watched the slow and painful dilatation of a fibroid cervix at full term, the comparison cannot fail to be instructive. The following case is also of such interest in this connection that I cannot refrain from quoting it:—

Mrs. O.C., aged 28, came to me with the following history: In her last confinement some five years ago, her cervix was lacerated bilaterally right up to the vaginal roof. Neither before this labour nor for twelve months afterwards did she have any dysmenorrhœa, but about a year after the birth of this last child she had trachelorrhaphy done, and ever since this operation she has suffered exquisite pain before and with the monthly period. Drugs do her no good, and relief is only gained by occasionally dilating the cervix between two menstrual epochs. Vaginal examination shows that the plastic operation was completely successful, but on each side of the cervix, and reaching up to the level of the supra-vaginal portion is a dense fibroid cicatrix. It is clearly in the resistance to dilatation offered by these dense deposits that the causation of the dysmenorrhœa is to be found.

D. HYPERÆSTHESIA OF THE NERVE ENDINGS AT THE LEVEL OF THE OS INTERNUM.

It is seldom that the uterine sound can be passed without giving rise to some pain, when the point of the instrument passes through the internal os, but under certain morbid conditions the ordinary pain becomes so intensified that it amounts to real agony, and patients will sometimes volunteer the remark that the pain produced resembles exactly that experienced during the onset of a menstrual time. If we note such cases carefully, it will be observed that one of two morbid states is usually present, either the patient is suffering from some existing inflammatory affection, such as endocervicitis or endometritis, with the discharges characteristic of these diseases, or else there is evidence that inflammatory troubles have existed, though their active stage is over and discharge has ceased; and here an exquisitely tender condition of the recently irritated nerve endings is all that is left to tell us of what has come and gone. In either of these cases pressure or dragging on the hyperæsthetic nervous tissue, such as may be caused by the passage of a sound or bougie, is sufficient to cause horrible suffering. A moment's consideration will make clear to us that the dragging open of the os internum by the uterine retraction connected with the early stage of menstruation will act precisely in the same way as pressure from within, and thus in these cases the exquisite pain felt in the first part of the monthly sickness is readily accounted for. In this connection it will be interesting to study a quotation from Dr. Matthews Duncan's paper on Spasmodic Dysmenorrhœa;¹ comparing the after-pains of parturition to spasmodic dysmenorrhœa, the great obstetrician remarks:

You will often read in books that when a woman has after pains there is a clot or a retained bit of placenta, or something which the uterus is attempting to expel, and this may be true, but such after-pains are not severe. That is not a disease, that is a healthy condition of the womb; the womb is doing its duty, as it were, and such after-pains are not very

painful. The real disease of after-pains is a disease in which the recently emptied uterus goes into the most violent and painful contractions without any discoverable object in view; and a severe case of this kind is a most painful disease, far more painful than the after-pains which come to expel a clot or a bit of retained placenta. Now these violent after-pains are, I believe, connected not only with a morbid condition of the muscular tissue, but chiefly or primarily with a catarrhal condition of the mucous membrane covering the inside of the body of the uterus, a condition not without several analogies with the healthy menstruating uterus.

In other words, contractions, which are only moderately painful when nerve endings are healthy, become the cause of the most acute suffering when catarrhal inflammation has left these endings in a condition of over-sensitiveness. It is scarcely necessary to quote cases illustrative of this variety of dysmenorrhœa, or to dwell longer on the causation of the pain, for instances will come readily to the memory of anyone who has much to do with obstetric medicine.

A more practical point is the question of suitable treatment. Briefly, this may be summed up as follows: When endocervicitis or endometritis is present, these conditions should be removed by the use of leeching, saline aperients, the application of carbolic acid or iodised phenol, and hot douching; but when no evidence of active disease can be found and a tender condition of the nerve endings is alone recognisable, then rapid dilatation will bring about a cure most speedily. How this latter remedy acts is not quite clear, but it is an undoubted fact that a very moderate amount of dilatation will diminish to a most marked extent the sensitiveness of the tissues at the level of the os internum.

In reviewing the subject of dysmenorrhœa, then, it is not for a moment claimed that a complete explanation has been afforded of the causation of menstrual suffering in every case which may come under observation, nor that in any individual case the pain must be attributed to one, and only one factor. Doubtless in some instances several causes of pain may exist, and it is only by a careful study of each case, and by endeavouring to recognise and remove in turn each factor, that a complete cure can be effected. A single case will illustrate this point:

Mrs. R., aged 35, has borne five children; since the last confinement four years ago, she has had increasing pain with her monthly times. The pain commences some twenty-four hours before the onset of the flow, and lasts into the second day. There is some clotting at the times, and moderate yellow discharge between the periods. The passage of the sound gives great pain; the canal measures 3 inches. Uterus is normal in position; the cervix is hypertrophied, and its tissues unduly hard. Leeching and the application of iodised phenol on a Playfair's probe to the cervical canal greatly relieved the dysmenorrhœa, and checked the yellow discharge. It was not, however, until ether had been given and the canal had been dilated up to Hegar No. 12 that perfect freedom from pain was obtained. Here, doubtless, hyperæsthesia of the nerve endings, resulting from cervical catarrh, was one cause of the pain, and rigidity of the cervical tissues another; both had to be corrected before a perfect cure was obtained.

Simplicity of explanation always carries a charm with it, and far-fetched theories seldom convey the germs of truth; it is possible, however, to err on either side, and, in avoiding stretches of mere imagination, to reduce simplicity to absurdity. The only safe and scientific method in any investigation is to study clinical facts, to compare one natural process with analogous workings of Nature, and, when facts do not perfectly fit with theory, to wait till more extended investigation gives a clearer and more perfect knowledge. Much of the difficulty which attends the study of dysmenorrhœa arises from the efforts of well-intentioned workers to make one theory explain all the phenomena of menstrual pain, and from neglect of the wise adage of the great French physician.—that before fact theories must always bend.

METROSTAXIS AND MENSTRUATION AFTER OPERATION ON THE BROAD LIGAMENT.

By W. JAPP SINCLAIR, M.D., M.R.C.P.,
Professor of Obstetrics and Gynaecology Owens College, Manchester.

FREQUENTLY after operation of any kind on the broad ligament, and invariably after operations involving the ligation of any considerable number of veins in the broad ligament, there occurs an oozing of blood from the uterus. It begins twenty-four or forty-eight hours after the operation, and continues, as a rule, for several days. This discharge of blood

¹ *Diseases of Women*, p. 132.