AN INQUIRY

REGARDING THE

IMPORTANCE OF ILL-EFFECTS

FOLLOWING THE USE OF

ANTIPYRIN, ANTIFEBRIN, & PHENACETIN

CONDUCTED BY

THE THERAPEUTIC COMMITTEE OF THE BRITISH MEDICAL ASSOCIATION.

D. J. LEECH, M.D., F.R.C.P., *Chairman.*WILLIAM HUNTER, M.D., *Honorary Secretary.*

SCOPE AND CHARACTER OF THE INQUIRY.

THE Committee have recently conducted an inquiry into the importance of the ill-effects which occasionally attend the use of the three chief antipyretic and analgesic agents phenazone (antipyrin), acetanilide (antifebrin), and phenacetin.

Since their introduction these drugs have had a very extended use at the hands of the profession; and so well have the merits of at least one of them (antipyrin) as an analgesic agent become known, that a large demand has arisen for it amongst the general public, independent altogether of medical recommendation or supervision.

An impression, however, has undoubtedly at the same time arisen that their use is frequently attended by ill-effects, and that these are of such a character and frequency as to limit very materially the general usefulness of these drugs. This impression is based on the numerous reports of isolated cases that have from time to time appeared in the medical press; and so general is the impression that it has sufficed to deter a certain number of practitioners from using the drugs at all. Considering the very widespread use which these drugs still continue to have, the Committee felt it to be important to obtain more trustworthy data regarding their alleged ill-effects than have hitherto been forthcoming.

In determining what form their inquiry should take, the Committee were guided by the following considerations. To obtain the necessary data it was important to have a certain variety as well as volume of experience represented. The experience of a few individuals, however large it might be, or that gained in hospital practice, even if available, was insufficient for their purpose. In hospital practice the drugs are only used for some very definite and immediate purpose—to control temperature or allay pain—under circumstances where their action can be carefully watched, and any ill-effects can be immediately observed and effectively combated.

In private practice, on the other hand, the sphere of employment of these drugs is a much more varied and extended one, corresponding to the wider and more varied demands that are there met with. A wider licence has necessarily to be given to the patient in using them; and ill-effects, if they arise, cannot be immediately recognised and dealt with, and become therefore more developed. It is under such conditions, then, that ill-effects are most likely to be met with; and it was therefore from the experience of private practice rather than from the narrower and less varied experience of hospital practice that the information sought for could best be obtained.

And here one of the chief difficulties of such an inquiry presented itself. It was obviously undesirable for many reasons to make the inquiry a general one. The drawbacks connected with the collective investigation method of acquiring information are too numerous and striking to require any setting forth here. As a method for the investigation of problems requiring accurate observation

and sound judgment, the method is not only useless—it is positively misleading. It can at its best be only a record of impressions, and impressions can never prove a substitute for carefully observed facts.

In the present instance the inquiry was framed in such a way as to reduce this source of error to a minimum. It was directed not so much to elicit the nature of the ill-effects—a subject regarding which we already possess fairly definite information; still less to ascertain under what conditions and in which diseases such ill-effects are most frequently met with, though this is a subject of very great importance, regarding which we know as yet but little. Information on these and other collateral points would be interesting, but to be of value it would require to be based on careful comparative observations carried out by skilled observers over a large number of cases and diseases. The scope of the inquiry was, on the contrary, definitely limited. Its very object was to ascertain the "impression" which any ill-effects met with had made on the mind of the observer; in particular whether that impression was of such a kind as to prevent him using the drugs as frequently as he would otherwise have done.

To enable the Committee to gauge the relative value of these individual impressions each observer was asked to state (1) the amount of his experience in the use of the drugs, (2) the dosage he employed, and (3) the nature of any ill-effects he had observed, giving at the same time such details as might appear to him necessary having regard to the primary object of the inquiry.

And here, in passing, a word may be said regarding the field left open for such investigations. It is a very limited one, both as regards the drugs deserving the application of such a method of investigation to them, and still more as regards the kind of information to be sought for. So long, however, as new therapeutic agents of the character and potency of those now under consideration are introduced, so long will questions of the kind above indicated from time to time arise.

A drug cannot be potent for good in any specially marked degree, especially if its chief action be on the blood or the nervous system, without possessing necessarily great potentialities for evil if used too frequently or in excess. Notwithstanding this obvious consideration there is a great tendency on the part of a certain class of observers to push every new drug to the extent of producing ill-effects. Cases of this kind are at once recorded; and thus it may easily happen that an extensive literature may arise relating to the ill-effects connected with the action of a drug, while its beneficial properties pass without record. An exaggerated and possibly an utterly false impression may thus be created, which it may require some investigation of the kind here indicated to clear away.

In the case of the drugs now under consideration, the result has fully demonstrated the necessity which existed for such an inquiry; for the inquiry has supplied clear evidence that the frequency and importance of the ill-effects attending their action have been greatly exaggerated, to the extent, in the case of certain of them, of producing an entirely false impression. The results demonstrate further that in the very large majority of cases the ill-effects have been the direct result of injudicious and excessive dosage, so that the real fault lies not so much with the drug as with its mode of administration.

METHOD OF INQUIRY.

The following was the method of inquiry adopted:—A letter was sent to the Secretaries of thirty-four Branches, describing the object of the proposed inquiry, and inviting them to furnish the Committee with a limited list of names of the members of their Branch to whom, in their opinion, such an inquiry might usefully be addressed.

In response to this, names were furnished by the secretaries of the following twenty-seven Branches, namely:

1. Aberdeen, Banff, and Kincardine.	10. Lancashire and Cheshire.
2. Bath and Bristol.	11. Midland.
3. Birmingham and Midland Counties.	12. North of England.
4. Border Counties.	13. Northern Counties of Scotland.
5. Cambridge and Huntingdon.	14. North of Ireland.
6. Dorset and West Hants.	15. Oxford.
7. East Anglian.	16. Perthshire.
8. East York and North Lincoln.	17. Reading and Upper Thames.
9. Edinburgh.	18. Shropshire and Mid-Wales.
	19. Southern.
	20. South-Eastern.

21. South-Western.
22. Staffordshire.
23. Stirling.
24. Thames Valley.

25. West Somerset.
26. Yorkshire.
27. South Wales and Monmouthshire.

The following letter of inquiry was then issued :

The Therapeutic Committee is desirous of gaining information regarding the nature, frequency of occurrence, and importance of ill-effects following the use of the above drugs (antipyrin, antifebrin, and phenacetin).

There seems to be an impression that these effects are of such a character and frequency as materially to limit the usefulness of these drugs as antipyretic and analgesic agents.

The Committee wish to ascertain how far such an impression is well founded.

Would you kindly state for the information of the Committee, giving briefly such details as may appear to you necessary, having due regard to the object of the inquiry as above stated :

1. The amount of experience you have had in the use of these drugs, whether as antipyretic or analgesic agents.
2. The doses you habitually give.
3. The nature of any ill-effects you have observed.
4. Their comparative frequency.
5. Your opinion as to their comparative importance. Do they materially limit the usefulness of the drugs ?

In the case of one Branch, the South-Wales and Monmouthshire, the inquiry was conducted independently by a local Therapeutic Committee, with Dr. D. R. Paterson as honorary secretary, acting in co-operation with the Central Committee. The terms of the inquiry were the same as those formulated by the Central Committee.

The Committee desire to acknowledge the help thus afforded them in the conduct of the inquiry by the secretaries of the various local Branches.

In reply to their circular, reports to the number of 220 have been received. These have been tabulated and carefully analysed, and the following is a summary of their chief contents.

It is particularly to be noted that throughout this report the sphere of usefulness of the drug is not touched upon. The value of the drug was a question purposely omitted from the inquiry. This was strictly limited to the question to what extent, assuming that the drug was a valuable one—as it undoubtedly is—this value was lessened by the occurrence of ill-effects.

PHENAZONE (ANTIPYRIN).

The amount of experience with this drug represented in these reports is a very large one—very much larger, as might be expected, than with the other two.

Dividing the reports into two groups, according as observers state their experience to be (1) extensive, or (2) limited, it is found that 167 out of the 220 reports (67 per cent.) are to be referred to the first group.

As regards the purpose for which the drug has been employed, the great majority have used it both as an antipyretic and as an analgesic agent. In the early years after its first introduction it was used very largely as an antipyretic; but there is practically a consensus of opinion on the part of those who report, that its chief use, and in many hands its sole use now, is as an analgesic. As an antipyretic it has thus fallen somewhat into disfavour, while as an analgesic it has in recent years gained steadily in repute.

NATURE OF THE ILL-EFFECTS.

There is great unanimity amongst observers as to the nature of the ill-effects occasionally met with. The only difference of opinion is as to their degree—"slight," or "marked," or "alarming," or "dangerous," as the case may be. They range in severity from the mildest and most evanescent of rashes to the most alarming and even fatal collapse.

The rashes may be dismissed with a word. They are variously described as erythematous, measly, or urticarial in character, with or without considerable accompanying œdema. They have been met with by every observer who has had any extended experience of the drugs. In the great majority of cases they have been the result of idiosyncrasy on the part of the patient, independent altogether of the size of the dose or the nature of the disease. Thus, in one instance a 10-grain dose caused urticaria with salivation twice in the same patient; in another a dose of 3 grains caused an urticarial rash, with dizziness and loss of power in legs; in another, swelling of face with lividity and dyspnoea was always caused even by the smallest doses.

Only a few observers, however, take any special notice of the rashes; the large majority content themselves with merely noting their occasional occurrence. As an ill-effect they cannot properly be regarded. At most they occasion a little temporary alarm to the patient till their true nature is explained; or, if met with at the outset of a febrile illness, may lead to a little difficulty in deciding whether we are not dealing with the specific eruption of measles or scarlet fever.

The character of the ill-effects proper may be judged of from the subjoined table, in which is comprised the experience of twenty-two observers. Where stated, the dose which occasioned the ill-effect is also given; where it is not expressly stated, the usual dose employed by that particular observer is appended within brackets.

It will be noted that the ill-effects group themselves clinically into 2 divisions: (1) Those referable to an action on the nervous system, including the varying degrees of vasomotor disturbance, profuse perspirations, enfeeblement, cardiac depression, and irregularity, nervous excitement and collapse, in exceptional cases such marked effects even as loss of power of speech, and complete mania; and (2) those referable to an action on the blood and circulation—namely, the breathlessness, varying degrees of cyanosis, and lividity.

The ill-effects described are, indeed, precisely of the character which one would be led to expect from a consideration of the physiological and therapeutic action of the drug. A drug capable of reducing febrile temperature with the rapidity and to the degree which antipyrin in many cases admittedly can—still more, one capable of relieving the intense neuralgic pains of functional disturbance with the rapidity and completeness with which antipyrin so frequently acts—must possess a very powerful action on the nervous system, and through it on the metabolic processes going on within the body.

TABLE I.

No.	Nature of Ill-Effect, as Specified by Observer.	Dose in Grains.
1	Weakness and shakiness ...	(20 to 40).
2	Serious collapse in a case of typhoid fever ...	30 twice.
3	Syncopal attack on one occasion ...	15.
4	Symptoms of collapse ...	20.
5	Enfeebling in its action ...	(5 to 20).
6	Depressant ...	(12 to 20).
7	" ...	(20).
8	Cardiac weakness and irregularity ...	15 every 4 hours.
9	Great depression ...	30 at hourly intervals.
10	" and exhaustion ...	15 every 4 hours.
11	Alarming depression ...	45 in 8 hours.
12	Cyanosis and dangerous cardiac depression ...	20.
13	" " " " ...	20.
14	Depressant on heart ...	(15 to 20).
15	Excessive sweating, cyanosis, and partial collapse ...	(20).
16	Vasomotor pains, lividity, profuse perspiration, tendency to collapse, and to pneumonic congestion ...	(30).
17	Collapse and death in a case of rheumatic fever ...	80 in 5 hours
18	Alarming faintness in anæmic individuals ...	(15 to 20).
19	Languor and depression ...	(15).
20	Weakening effect in 10 per cent. of cases ...	20 repeated in an hour.
21	Serious collapse in a case of typhoid ...	2 doses of 30.
22	Loss of speech, lasting 24 hours, in a case of commencing meningitis which subsequently proved fatal ...	7½ repeated in an hour.
23	A condition of mania from long-continued use of the drug, recovered from when drug withheld ...	
24	Dyspnoea and much nervous excitement once out of many hundreds of cases ...	5.
25	Salivation with urticarial rash twice in same patient ...	10.
26	Dizziness and loss of power in legs ...	3.

Ill-effects of the above character may as naturally be expected to follow the administration of large doses of this drug as excessive stupor and twitchings that of morphine and strychnine respectively. Their occurrence in exceptional cases would not, therefore, *per se* prove the drug to be a dangerous one. For this it would be necessary to show that their occurrence was independent of the size of the dose administered.

The relation of ill-effects to dosage employed becomes thus an important consideration in estimating the harmfulness of a drug. This is a point which will presently be considered. In the case of antipyrin it will be found to be an interesting and instructive one.

THEIR COMPARATIVE FREQUENCY.

The mere list of ill-effects above noted is a somewhat formidable one, but it loses much of this character when their frequency of occurrence is considered.

Thus, out of 189 observers who report on antipyrin, no fewer than 138, or 73 per cent., have never observed any ill-effects at all worth mentioning.

Considering how large is the experience of the action of this drug represented in these reports, this large proportion of observers who have no ill-effects to record is very remarkable. Almost more than any other fact could do, it seems to testify to the comparative harmlessness of this drug when properly administered.

Further testimony, no less strong in the same direction, is borne by the experience of the remaining 51 observers (27 per cent.) who describe ill-effects; for of this number 17 expressly state that any ill-effects they had observed were of so slight and trivial a character, and only met with in isolated instances, or after injudiciously long-continued use of the drug, that they could not attach the slightest importance to them.

The number who describe ill-effects is thus reduced to 34, the character of these being indicated in the foregoing table. Of these only two venture to give their experience as to their frequency of occurrence in statistical form, one stating that he has met with them in about 1 per cent., the other in about 10 per cent. of cases. Two others state that ill-effects are comparatively frequent in their occurrence, one of these, who gives it in doses of 10 grains, stating that it causes "distinct lowering of the vital powers"; the other, who formerly gave it in 30-grain and who now employs it in 10-grain doses, describing the ill-effects as including "vasomotor pains, lividity, profuse perspiration, tendency to collapse, measly rash, and tendency to pneumonic congestion." The remaining 30 either state that the occurrence of ill-effects is extremely infrequent, or expressly state that any ill-effects they have observed have occurred as isolated instances in the course of many years' experience of the drug, and extending over many hundreds of cases. So rarely, indeed, have they met with any ill-effects, that they attach little or no importance to them, and have in no way been deterred from use of the drug by fear of them.

It is clear, then, if one may be allowed to judge from these reports, that ill-effects following the use of antipyrin are not only relatively but absolutely infrequent in their occurrence. One observer expresses his opinion that they are "rarer than the idiosyncrasies of the iodides or quinine;" and certainly so far as the above reports go, this would appear to be the case.

The impression to the contrary, which is widely prevalent, and which is testified to by one observer who states he has never observed any ill-effects himself, but has been deterred from its freer use by the fear of its depressing effect, receives therefore no support whatever from these reports.

THEIR IMPORTANCE.

The main object of the inquiry, as has been stated, was to ascertain how far the ill-effects occasionally met with were of such a kind as to limit the general usefulness of the drug. The answer given by these reports to this question has been already in great part anticipated by the consideration just given to their frequency. It has been indicated that ill-effects may naturally be expected to follow the injudicious use of so powerful a drug as antipyrin undoubtedly is. To what extent, then, are the ill-effects above recorded to be ascribed to this cause; or, on the other hand, are they directly traceable to the dangerous and uncertain action of the drug itself?

In answering this question, it becomes important (1) to ascertain what is the relation of ill-effects to the dosage employed; and (2) to consider the circumstances of each particular case in which ill-effects have been observed.

1.—*Relation of Ill-effects to Size of Dose.*

In estimating dosage, regard has to be had to the frequency of administration as well as to the size of the individual dose. Ill-effects may arise from small doses too frequently repeated as much as from larger doses. In estimating the influence of dosage, a considerable difficulty arises. Hardly any two observers administer the drug in the same way. It is difficult, then, for the purposes of such an inquiry as the present to tabulate the dosage. One observer employs a dosage of 5 to 10, another of 10 to 20, and a third of 20 to 30 grains. It would be possible to strike an average in each case, and to compare these with one another—to state that the average dose was $7\frac{1}{2}$, 15, and 25 grains respectively. Certain results might thus be got, but how far they would be trustworthy it is permissible to question. A more reliable method, as it seems to the Committee, is to have regard to the minimal dosage alone, to the exclusion altogether of the larger doses occasionally employed—and for the following reason.

The minimal dose employed by an observer is to a certain extent an important index as to the degree of care he is accustomed to exercise in beginning the administration of a drug. If one observer states that he uses a drug in doses of 5 to 10 grains, while another states that he habitually employs doses of 15 to 30 grains, it is clear that in the case of a powerful drug ill-effects connected with the size of the dose are more likely to be met with in the practice of the one who usually commences with 15, than of the one who habitually begins with 5-grain doses.

It is of interest, then, from this point of view, to compare the minimal dosage employed by the various observers. Out of 167 observers who record their dosage, 134, or 80 per cent., employ doses of 10 grains or under as their usual minimal doses of antipyrin—30 per cent. commencing with not more than 5 grains; while the remaining 33, or 20 per cent., habitually employ as their minimum doses ranging from 15 to as high as 30 grains. The interesting point, then, comes out that it is almost exclusively in the practice of the latter that ill-effects have been noted.

For among the 138 who have not met with any ill-effects, no fewer than 119—86 per cent.—habitually employ doses of 10 grains or under as their minimal dose; while on the other hand, out of the 51 who describe ill-effects, 37—73 per cent.—are in the habit of giving as their minimum doses as much as 15, 20, and 30 grains.

So far, then, as these reports are concerned, there would appear to be a close and direct relation between the size of the dose and the occurrence of ill-effects. The larger the initial dose the more frequent are ill-effects met with.

2.—*The Circumstances of each Particular Case.*

The accuracy of this conclusion is further borne out when we consider the circumstances of each particular case in which ill-effects have been noted. These have already been tabulated (Table I). It will be noted that, with only a few exceptions, the dose habitually employed by the observer who describes ill-effects, or the dose which produced the ill-effects in the particular case, was a large one—in certain cases, indeed, surprisingly so. For example, it need occasion little surprise that 30 grains of antipyrin administered hourly caused great cardiac depression, that weakness and shakiness were caused in certain cases by doses of 20 to 40 grains, or that 80 grains in five hours caused collapse and death in a case of rheumatic fever.

Of the 51 observers who record ill-effects, 17 expressly state that they were of such a slight character, or so rarely met with, as to be of no importance whatever. The remaining report cases such as the foregoing (Table I). It will be noted that in most of the cases the dose has been 15, 20, or 30 grains, repeated after an injudiciously short interval. A few describe ill-effects produced by doses of 10 grains or under; but, with a few rare exceptions, these have only been after frequently repeated administration of the drug. The exceptions referred to are obviously cases of idiosyncrasy such as are common with almost every drug. For example: (1) dyspnoea and much nervous excitement, reported in one instance out of many hundreds by a dose of 5 grains; (2) urticaria and salivation following a 10-grain dose twice in the same patient; (3) dizziness and loss of power in legs

should be borne in mind as the proper initial dose of this drug. It will presently be considered to what extent the dosage employed has, in the case of antifebrin, as in that of antipyrin, been accountable for the ill-effects noted.

THEIR COMPARATIVE FREQUENCY.

The list of ill-effects noted with antifebrin is not only a very formidable one in itself, but, unlike the case of antipyrin, it loses none of this character when we consider the frequency of their occurrence. In the case of antipyrin, notwithstanding a very large experience of the drug, only 27 per cent. of observers (37 out of 189) had any ill-effects to describe. With antifebrin, on the contrary, from a much more limited experience (not exactly estimable, but probably not exceeding one-hundredth), as many as 44 per cent. of observers (44 out of 100) have met with ill-effects. This fact must be regarded as pointing to the much more frequent occurrence of ill-effects with this drug than with antipyrin. It is probable that the recognition of this has to some considerable extent contributed to the preference which is given by most observers to antipyrin over antifebrin.

A further difference in the case of the two drugs is to be noted—namely, that while most of the cases in which ill-effects have followed the use of antipyrin have been isolated instances in the experience of the individual observers; with antifebrin, on the other hand, out of a much smaller experience, dangerous effects have been noted several times by the same observers.

So far, then, as these reports go, we must conclude that ill-effects following the administration of antifebrin are both absolutely and relatively more frequent in their occurrence than is the case with antipyrin.

THEIR IMPORTANCE.

The importance to be attached to them is also greater than we have seen to be the case with antipyrin. This is evidenced in several ways, both directly and indirectly, in these reports. In the first place, the ill-effects are of a more severe character and more frequent occurrence than with antipyrin; secondly, with but few exceptions observers are agreed that the action of this drug requires more careful watching; and, thirdly, there is the indirect evidence afforded by the preference which the great majority of observers give to antipyrin or phenacetin over antifebrin. This preference may, of course, be due to the greater range of usefulness of antipyrin as compared with antifebrin, and be independent altogether of their relative powers for evil. Be that as it may, it is clear from the list of ill-effects noted with antifebrin (Table II) as compared with that found with antipyrin (Table I) that so far as ill-effects are concerned the preference must be given to antipyrin. Notwithstanding these ill-effects, a few observers still give the preference to antifebrin. Thus one who had not met with any ill-effects from antipyrin, but had more than once observed alarming collapse with antifebrin, expresses his opinion that antifebrin is the most perfect antipyretic we have. Another, who observed great cyanosis after two doses of 10 grains, nevertheless considers that the ill-effects are of no importance. And a third even goes so far as to say that ill-effects are more common with antipyrin than with antifebrin. Most, however, consider that the liability to the occurrence of ill-effects does to a certain extent interfere with the usefulness of the drug, the chief drawback being its markedly depressant action.

We have now to consider the two points we have seen to be of importance in gauging the relative importance of ill-effects following the use of a drug, namely, the relation of these to the dosage employed, and the circumstances of each particular case.

1. *Relation of Ill-effects to Dosage.*—No such direct relation is here to be made out as was seen to be the case with antipyrin. The dosage employed appears to have been much the same with those who have observed ill-effects as with those who have not. In both cases the proportions are almost the same, 72 and 75 per cent. of observers respectively having used doses of 4 to 10 grains as their minimum. This difference in the case of these two drugs would be an interesting one if it could be clearly established. If it could be shown that the occurrence of ill-effects was to a certain extent in-

dependent of the largeness of dose, this fact more than any other would tend to prove that antifebrin was more inconstant in its action, and correspondingly more dangerous than antipyrin.

A powerful drug can be used with safety so long as its action is a fairly constant one, varying chiefly with the dose employed. Danger at once arises, however, if, along with the power of doing harm in over doses, the drug is found to be uncertain in its action. In the case of antifebrin, the evidence we are now considering would appear to point to its being less safe and less constant in its action than antipyrin. It is probable, however, that with more data to go on than are to be found in these reports it may appear that with it as with antipyrin ill-effects were generally brought about by injudicious dosage. A large number of observers do not state any dosage at all.

And, indeed, when we consider (2) *the particular circumstances under which ill-effects are described* (Table II), the conclusion is, perhaps, admissible that the dosage employed by observers in the case of this drug has been singularly injudicious. Similar large doses have apparently, however, been given by others without any ill-effects.

SUMMARY.

Notwithstanding the experience of many apparently to the contrary, the conclusion must be permitted that to give antifebrin in doses of 5, 6, 8, and even 10 grains, still more to repeat these after a short interval, is a highly injudicious procedure. Such doses are altogether excessive. They are equivalent to about 25, 30, 40, and 50 grains of antipyrin. The repute of the drug has probably suffered in the past from the circumstance that this fact of its greater strength has been overlooked. Indeed, the dosage employed has been so large as to lead one to surmise that in the minds of many observers antifebrin is regarded as a drug of the same strength as antipyrin.

PHENACETIN.

The amount of experience of this drug represented in the reports is about the same as that with antifebrin, if anything probably larger. It is apparently not so widely employed as the latter, only 80 observers reporting regarding it as compared with 100 who report regarding antifebrin. A somewhat larger proportion of this number, however, already possess a considerable experience of its action—namely, 55 per cent., as compared with 44 per cent. in the case of antifebrin. It has been used largely both as an antipyretic and as an analgesic, and apparently with equally good results. This drug appears to be in exceedingly good repute with those who have any extensive experience of it, more especially as an analgesic.

NATURE OF ILL-EFFECTS.

Out of a total of 80 observers, only 7 have any ill-effects to record

TABLE III.

No.	Nature of Ill-Effect. as specified by Observer.	Dose in Grains.
1	Collapse on one occasion	5 every 3 hours for 3 days.
2	Extreme weakness, cyanosis, and feebleness of pulse	(10 to 15 every 4 hrs.)
3	Cyanosis once	7 every 4 hours for 3 days.
4	Slight giddiness once	(5 to 10).
5	Depression, although not often	(5 to 10).
6	Lividity and diaphoresis	(10 to 20).
7	Subnormal temperature, coldness, shivering	(5 every 4 hours).

If we eliminate the last 4, in which the ill-effects described are obviously of slight degree and importance, there remain only 3 where the ill-effect was at all specially marked. And it was of the same character as that already noted with regard to antipyrin and antifebrin, namely, cyanosis and collapse. They were only met with on one occasion by each observer, and their presence is sufficiently accounted for by the dosage used, namely, 5 grains every three hours for three days, 10 to 15 grains every four hours, and 7 grains every four hours for three days.

FREQUENCY AND IMPORTANCE.

Ill-effects with this drug have thus, so far as the experience in these reports goes, been strikingly infrequent. As regards the importance attached to them when they have been met with, only one (No. 5) of seven observers who have noted them has been led on their account to prescribe the drug less frequently. The others expressly state that they are of no importance whatever, and affect in no way the great value of the drug. There is, indeed, striking unanimity amongst observers as to the great value of this drug, especially as an analgesic.

As regards the dosage employed, observers are about equally divided, one-half using 5 grains or less to begin with, the other half using doses of 8 to 10 grains.

The drug appears thus to have a notable freedom from injurious action. Doubtless, if used injudiciously, it would, like antipyrin or antifebrin, although possibly to a smaller degree, produce similar untoward effects.

GENERAL CONCLUSION.

As regards their freedom from ill-effects, so far as these reports show, the drugs may be placed: (1) Phenacetin, (2) Antipyrin, and (3) Antifebrin.

THE HEALTH OF THE PRINCESS OF WALES.

VARIOUS disquieting rumours have been current during the week regarding the health of the Princess of Wales. We have, however, excellent authority for the reassuring and satisfactory statement that Her Royal Highness is now progressing satisfactorily, and that there is no reason for alarm. We understand that, after passing through an attack of influenza, Her Royal Highness suffered a relapse which so seriously affected the throat that fears of diphtheria were entertained. Under these circumstances we believe that Sir William Broadbent was hastily summoned from London. These symptoms, however, quickly passed away, and since then there has been no reason for anxiety. Unfortunately, influenza is a malady which leaves considerable debility and demands care during convalescence—a necessity which has involved arrangements suggesting to some a present danger which we are happy to be able to state no longer exists.

HOSPITAL AND DISPENSARY MANAGEMENT.

THE CITY OF LONDON LUNATIC ASYLUM.

ADMISSION OF PAYING PATIENTS.—From the last annual report we learn that the percentage of recoveries calculated on the admissions was 28.73. This low percentage is accounted for by the admission of a large proportion of incurable cases. In fact, at the close of the year 1892 there were only 13 patients in the asylum who were deemed curable. Post-mortem examinations were made in 32 out of the 35 patients who died, and the proportion of deaths to the total number under treatment was 7.02, or 1.17 per cent. below the average of similar institutions. We are glad to note that the Committee have availed themselves of the power conferred by recent legislation and have commenced to admit paying or private patients. This will prove a great boon to many a little above the pauper class who can afford to pay the reasonable charge of 21s. per week, and we are not surprised to find that in the first year of the experiment there were 26 private patients admitted. Dr. Ernest White, the medical superintendent, reports favourably of the change of dietary which has been effected during the year, with the result that the patients are healthier in appearance, and in most cases have gained weight. Three cases of typhoid fever occurred, due probably to defective and out-of-date sanitary fittings and drainage, and this is receiving the attention it urgently needs.

WESTERN OPHTHALMIC HOSPITAL, MARYLEBONE ROAD.

MR. HASTINGS NEALE forwards us the following extract from the minutes of a special general meeting of governors of this hospital, held on December 20th, 1893, with the request for our opinion thereon:

Agenda.—To consider the freeing of the hospital and to frame rules. After considerable discussion the following resolution was carried by a large majority, there being no votes against, but two governors did not vote:

"That in future the hospital be absolutely free. That no single person whose income exceeds 30s. shall be eligible as a patient. No married persons whose joint income exceeds 50s. excepting where there are more than four children, when the limit is to be £2 15s.

Unless an investigation should show any reason to the contrary, this rule to be rigidly observed."

It was further unanimously resolved that the framing of rules to carry above resolution into force be referred to the committee.

We have much pleasure in giving publicity to the step which the governors of the Western Ophthalmic Hospital have taken towards freeing that institution from the incubus of the "hospital-letter" system. No doubt that system is one which is oppressive to the poor and unsatisfactory to the subscribers. The former, if deserving and really ill, are put to serious annoyance and hardship by the necessary search for letters, while the latter have really no means, and often no leisure, to discriminate between the deserving and undeserving applicants, the appropriate and inappropriate cases.

We have received several letters bearing on this matter published in a Birmingham paper, which testify to the fact that public attention is now becoming awake to the evils of this system. So far, therefore, the governors of the Western Ophthalmic Hospital have our hearty good wishes for the success of their efforts to convert their institution into a "free" hospital. But in this, as in most things, the method is of as much consequence as the aim. Hitherto the governors have merely resolved on the establishment of a wage limit. We feel rather doubtful whether this, though it is no doubt an important, ought to be the sole, consideration in fixing the eligibility of a candidate for hospital treatment.

It seems obvious that a person who requires long or expensive or difficult treatment is less able to pay for it than one of the same pecuniary position whose case is less grave; so that it seems to us to follow inconceivably that the medical nature of the case ought to be taken into consideration as well as—and even, we should say, before—the patient's pecuniary circumstances. This can be done by requiring a medical certificate with all cases which are not urgent. We must again repeat that the vast majority of out-patients, ophthalmic and others, are suffering from diseases which are in no respect urgent, and in which a short delay for ascertaining their suitability for hospital treatment is of no consequence whatever; and further, that they are as able to pay for their medical treatment in ordinary exigencies as for any other necessary of life, and ought to do so, whether on the provident principle or otherwise. At the same time, if the disease is obstinate or complicated, the medical attendant may well wish to refer the case to a hospital either for an opinion or for treatment.

An inquiry, therefore, into the mere pecuniary condition of the patient seems to us unsatisfactory. Yet even this inquiry is by no means a simple matter. If insisted on, it must be made a reality, and can only be made so by means of some such machinery as is adopted by the Charity Organisation Society—and everyone knows how unpopular such methods are with the working classes. So that it seems to us that the governors of the Western Ophthalmic Hospital have undertaken a task the difficulties of which will become daily more and more apparent. Still, any honest effort to render medical charity more discriminating and more effectual deserves support, and we wish the present attempt all success.

FIRES AT LONDON HOSPITALS.

A FIRE broke out, early on the morning of January 5th, in the upper storey of the City of London Hospital for Diseases of the Chest, Victoria Park, E., which was not subdued until some considerable damage was done. One of the night nurses noticed a smell of burning in the attics, and opening the door discovered the corridor to be in flames. An alarm was raised, and the patients were removed under the supervision of the resident house-surgeon to a place of safety without any injurious effect. The firemen, who arrived soon after the alarm, extinguished the flames within an hour. The conflagration was confined to the upper floor and the roof. The origin of the outbreak is unknown, but it is thought to be due to a defective flue.

On January 6th a fire broke out at the Royal Free Hospital, Gray's Inn Road, but, owing to the prompt and energetic action of the officials of the institution, the outbreak was prevented from causing extensive damage. The room in which the fire occurred was one of the top isolated wards. It was much damaged, as was also an apartment below used as an isolated ward, and the board room.

EDINBURGH ROYAL INFIRMARY.

THE annual meeting of the contributors to the Edinburgh Royal Infirmary was held on New Year's Day, the Lord Provost presiding. From the report submitted, it appeared that the daily average of children in the hospital was 72, the great majority of whom were in the surgical wards. The daily average of adult patients was 690, the maximum at any one date 734, and the minimum 561. The average time under treatment was 28.4 days, as against 27.4 in the previous year; 4,133 patients were admitted from the country districts. The percentage of deaths to the whole of the medical and surgical cases under treatment was 7.00; deducting the deaths (142) that occurred within forty-eight hours after admission, the percentage falls to 5.40; 25,000 out-patients were treated. Till the projected larger extension of the hospital has been effected increased accommodation of a temporary kind had been provided in the property belonging to the Corporation in Lauriston Lane. The extension of the Convalescent House at Corstorphine had increased the accommodation by 40 beds, large day rooms, etc., and the home now contains a daily average of 77 patients.

METROPOLITAN HOSPITAL SATURDAY FUND.

FROM the report of the Council of the Hospital Saturday Fund just issued, it will be seen that the receipts from the workshops to November 25th were £10,847 as against £11,215 in the corresponding period of last year; and the street collection this year amounted to £3,095 as against £3,684 in 1892. Even with a balance of £1,114 remaining from 1892 as against £245 reserved from 1891, the total income of the fund on November 25th was £17,056 as against £17,176 at the corresponding date last year. The diminution of receipts is mainly attributed to trade depression.