A POST-GRADUATE LECTURE

ON

THE INFLUENCE OF THE ARTERIOLES IN
RELATION TO VARIOUS PATHOLOGICAL
CONDITIONS.

By SIR GEORGE JOHNSON, M.D., F.R.S.,

Emeritus Professor of Clinical Medicine, Consulting Physician to
King's College Hospital, and Physician Extraordinary to
Her Majesty the Queen.*(Concluded from page 843).*

THE PHYSIOLOGY OF NITROUS OXIDE ANÆSTHESIA.

IN the phenomena of apnœa with the resulting rapid impediment to, and even the complete arrest of, the circulation which occur when nitrous oxide gas is inhaled as an anæsthetic, we have an interesting confirmation of the results obtained by excluding air from the lungs; and on the other hand the knowledge obtained by experiments on animals killed by apnœa enables us more completely to interpret the facts of nitrous oxide anæsthesia.

Through the courtesy of the staff of the Dental Hospital in Leicester Square I have on many occasions during the last few years had the opportunity to observe the phenomena which result from the inhalation of the gas, and, in particular, I have carefully noted the changes in the character of the pulse during the successive stages of the inhalation.

In the early stage the respiration is deepened and quickened, while the pulse, which is usually quickened, always becomes fuller and firmer. A few seconds later the pulse is less full and more compressible, and if the inhalation be continued, the pulse becomes so small and feeble as to be scarcely, or even not at all, perceptible; at the same time the breathing is stertorous, the features are livid, and the pupils dilated, while there is more or less jactitation of the extremities, and often general tonic contraction of the muscles. Micturition rarely occurs, defecation still more rarely. That these symptoms are of an epileptiform character is, I think, unquestionable. The convulsions are sometimes so violent, especially in children, as to throw the patient from the chair.

After the readmission of air the pulse again becomes for a time full and strong, and less frequent, the lividity passes away, and consciousness quickly returns. There is obviously a very close resemblance between these symptoms and those which result from apnœa. When an animal is killed by inhaling nitrous oxide gas convulsions always precede death, and when the chest is opened immediately after death, the right side of the heart is found to be greatly distended while the left is nearly empty, the blood on the two sides of the heart being equally dark. The lungs are pale, their minute vessels are bloodless, and the lungs consequently collapse into a very small space when the chest is opened. These appearances are identical with those which are found after death from apnœa. What, then, is the explanation of the phenomena? It has been ascertained that at the temperature of the body nitrous oxide gives up no oxygen to the blood or the tissues. The gas rapidly replaces the oxygen in the blood and in the tissues; the full and strong pulse, the result of contraction of the arterioles excited by the passage of venous blood into the systemic arteries, is the counterpart of the high pressure in the arteries of animals deprived of air, while the small and weak pulse which follows, like the rapid fall of pressure in the advanced stage of apnœa, is the result of the impeded flow of blood through the lungs caused by contraction of the pulmonary arterioles, the heart at the same time being weakened by the small amount of blood, and that unoxxygenised, which passes into its nutrient vessels. Again the convulsive movements which are common to the two classes of cases admit of the same physiological explanation. When in an animal the inhalation of the gas is pushed to a fatal termination, convulsions are as constant and as violent as when death is caused by a ligature on the trachea. The spectroscopic examination of the blood im-

mediately after death shows that the hæmoglobin has been entirely deoxidised.

I find that some anæsthetists who are daily occupied in the beneficent work of administering nitrous oxide, with perhaps a natural unwillingness to admit that the freedom from pain which they ensure is the result of replacing vital air by an azotic gas, maintain that apart from or in addition to its acknowledged deoxidising effect nitrous oxide has a special anæsthetic influence. Of this specific action, however, so far as I can learn, they adduce no proof. True Dr. F. Hewitt states¹⁹ "that the anæsthesia is not due to asphyxia is probable from the fact that it is possible to secure absolute unconsciousness to pain, muscular relaxation, etc., from nitrous oxide mixed with such a proportion of oxygen that the development of obvious asphyxial phenomena is rendered physiologically impossible." Now in making this statement Dr. Hewitt has not taken into consideration the influence of the great solubility of nitrous oxide in the blood as compared with that of oxygen or nitrogen. If a mixture of nitrous oxide with 10 or 15 per cent. of oxygen were inhaled for a minute or so and if the expired gases were analysed it would be found that the relative amount of nitrous oxide and of oxygen has undergone a change, the nitrous oxide being less and the oxygen more in proportion than before the mixed gases were inhaled. The explanation is that a larger proportion of nitrous oxide than of oxygen is for a time retained in the blood, partly in solution and partly in a state of loose combination, and this excess deoxidises the hæmoglobin sufficiently to effect anæsthesia without the development of so-called asphyxial symptoms.

Then it is to be observed that the mixture of oxygen with nitrous oxide considerably retards the production of deep anæsthesia. Thus Dr. Hewitt states that while the average period of inhalation required to produce deep anæsthesia with nitrous oxide alone is 51 seconds²⁰ the mixture with oxygen requires that the inhalation should be continued for an average period of 110.5 seconds.²¹ The presence of oxygen, when not in excess, delays but does not entirely prevent the process of deoxidation.

Nitrogen is much less soluble in the blood than either nitrous oxide or oxygen, yet Dr. Hewitt has obtained, in my presence, distinct anæsthesia in two cases by administering a mixture of nitrogen with as much as 7 or 7½ per cent. of oxygen.²² Since that proportion of oxygen does not prevent, though it delays, the anæsthetic action of nitrogen, it is quite intelligible that with a larger amount of oxygen the much more soluble nitrous oxide should be an efficient anæsthetic. This, however, requires further explanation.

But before entering upon this there is one important practical point relating to the action of nitrous oxide as an anæsthetic to which I am anxious to direct especial attention, and that is the manner in which the circulatory failure occurs. This failure is not the result of ordinary syncope, but, as numerous observations and experiments have shown, the immediate cause is the arrest of the pulmonary circulation and the consequent over-distension of the right side of the heart. The remedy for this is the free access of air to the lungs, and in the almost infinitely small number of cases in which this fails the cautious inhalation of the vapour of nitrite of amyl, by relaxing the pulmonary arterioles, may succeed in overcoming the impediment. An experiment described in the earlier part of this paper affords a striking illustration of the influence of the vapour in removing the paralyzing over-distension of the right side of the heart, even while nitrous oxide gas was still being thrown into the lungs. Another method of relieving the over-distension of the right side of the heart and so increasing its contractile power is to open the external jugular vein. Dr. John Reid in an interesting paper "On the Effects of Venesection in Removing and Increasing the Heart's Action under Certain Circumstances,"²³ has proved by numerous experiments on animals that when, as a result of apnœa, the heart has nearly or quite ceased to beat, its contraction may be renewed or invigorated by "disgorging the right side of the heart."

THE ANÆSTHETIC ACTION OF NITROGEN.

It occurred to me, about four years ago, that some light might be thrown upon the anæsthetic action of nitrous oxide by observing the effect of inhaling nitrogen as nearly pure

as it can be obtained. Accordingly I obtained from the Scotch and Irish Oxygen Company of Glasgow a metal cylinder containing 100 cubic feet of compressed nitrogen, in which the proportion of oxygen present was only 0.5 per cent., with 0.3 per cent. of carbonic acid. As a preliminary trial Mr. Woodhouse Braine did me the favour to administer this gas in five instances to members of the staff of King's College who volunteered to submit to the experiment. The result in each case was rapid and complete anæsthesia, with general phenomena precisely similar to those resulting from the inhalation of nitrous oxide; consciousness quickly returned when the facepiece was removed, and there were no unpleasant after-effects.

After this Mr. Braine felt justified in administering the gas as an anæsthetic to patients at the Dental Hospital. On the first occasion nine patients took the gas. In every case there was complete anæsthesia, with general phenomena precisely similar to those resulting from nitrous oxide inhalation. The pulse was first full and throbbing, then small and feeble. In the advanced stage there was cyanosis, the pupils were dilated, and there was more or less jactitation of the limbs. The only difference, in the opinion of some of those present, was that the anæsthesia was rather less rapidly produced and somewhat less durable than that from nitrous oxide, though in each case the tooth was extracted without pain.

On a subsequent occasion the gas from the same cylinder was administered by Dr. F. Hewitt at the Dental Hospital.²⁴ Nine patients took the gas, with results identical with those observed on the former occasion. Dr. Hewitt says "an onlooker could not have detected any difference between the phenomena produced and those usually met with under nitrous oxide." The maximum period of inhalation required to produce anæsthesia was 70 seconds, the minimum 50 seconds, and the mean 58.3 seconds.²⁵ In one case two teeth were extracted without pain. In only one case was pain experienced, and in that case, the tooth having been broken and not extracted, the patient said she felt "a smashing up."

Having on several occasions witnessed the administration by Dr. Hewitt of nitrous oxide mixed with about 10 per cent. of oxygen, I determined to try a mixture of nitrogen with a small proportion of oxygen. I therefore obtained from the company before mentioned a cylinder containing 40 cubic feet of nitrogen with 3 per cent. of oxygen, and a second cylinder of the same capacity containing nitrogen with 5 per cent. of oxygen. The 3 per cent. gas was given by Dr. Hewitt to five patients, and the 5 per cent. to four patients. With the 3 per cent. gas the time required to produce anæsthesia varied from 60 to 75 seconds, the mean time being 67.5 seconds. In each case the tooth was extracted without pain, the duration of the anæsthesia being somewhat longer than with nitrogen alone. In each case there was lividity, dilatation of the pupils, and more or less jactitation.

With the 5 per cent. gas the time required to produce anæsthesia ranged from 75 to 95 seconds, the average being 87.5 seconds. In each case the anæsthesia was complete, during which one patient had three molars extracted; and although she said she "felt the last two," the sensation appears to have been that of a mechanical pull and not acute pain. In all four of these cases there was slight lividity before the facepiece was removed, but in only one was there slight jactitation of the limbs. Dr. Hewitt, in describing the results of these experiments with mixed nitrogen and oxygen, mentions one fact which I had not noted, namely, that "there was some excitement after the inhalation in a few of the cases." Such excitement is not uncommon after nitrous oxide inhalation.

In undertaking these experiments with nitrogen, I had little expectation that it would ever come into use as a substitute for nitrous oxide; the practical impossibility of keeping the gas in a liquid form would alone forbid this, but it occurred to me that if the inhalation of a gas which could not be supposed to possess a specific anæsthetic property were found to induce insensibility of which an experienced anæsthetist might say "an onlooker could not have detected any difference between the phenomena produced and those usually met with under nitrous oxide," this would have gone far to prove that nitrous oxide, like nitrogen, produces anæsthesia by depriving the blood and the tissues of oxygen.

It is a confirmation of this view that a longer inhalation

is required to produce anæsthesia when a small proportion of oxygen is mixed with either nitrous oxide or nitrogen, while as a result of this the available period of anæsthesia for operation is somewhat prolonged. The obvious explanation is that more of the gas is required to deoxidise the blood and the tissues when vitalising oxygen is mixed with an azotic gas, while a prolonged inhalation carries the process of deoxidation deeper into the fluids and tissues, and so lengthens the period of anæsthesia.

I learn from Dr. Hewitt that in giving nitrous oxide with oxygen he begins with two per cent. of oxygen and gradually increases it to about ten per cent. If nitrogen were given in the same manner and with the same proportion of oxygen the result would probably be identical.

If the late Dr. Snow had lived to see the general use of nitrous oxide as an anæsthetic, he might have referred to it as affording additional evidence in support of his inference from numerous observations and experiments that "the action of the volatile narcotics is that of arresting or limiting those combinations between the oxygen of the arterial blood and the tissues of the body which are essential to sensation, volition, and all the animal functions."²⁶ He demonstrated that these substances modify, and in large quantities arrest, the animal functions in the same way and by the same power as that by which they modify and arrest combustion, the slow oxidation of phosphorus and other kinds of oxidation unconnected with the living body when these narcotics are mixed with the atmospheric air.

PREVIOUS EXPERIMENTS WITH NITROGEN.

Until I read Dr. F. Hewitt's interesting book on *Anæsthetics*, I was not aware of the fact that about a quarter of a century ago Dr. Burdon Sanderson, Dr. John Murray, and Mr. Turner had administered nitrogen to six patients for tooth extraction at the Middlesex Hospital, the results of which are published.²⁷ The object of the experiment was to solve the question: "Does nitrous oxide act as an anæsthetic by excluding oxygen?" In the first two cases the inhalation was discontinued, as "the inhaler did not effectually exclude air." In the other four cases, "with a perfectly fitting inhaler," the results were as different from those which we obtained with nitrogen at the Dental Hospital as the authors rightly declare them to be from the action of nitrous oxide. Thus none of the patients showed any marked effects until the inhalation had been continued for two minutes, and for the production of insensibility the inhalation had to be continued for 3 min., 3 min. 10 sec., 4 min., and 4 min. respectively. There was no lividity, "a marked contrast" (as they say) "to the deep blue produced by nitrous oxide." In two of the cases extraction was perfectly painless, the other two said that although they had no pain they "felt it."

The report concludes by the statement that "the facts above related indicate that there is an essential difference in the mode of action of the two gases" namely, nitrogen and nitrous oxide.

What, then, is the explanation of the contrasted results of these experiments and of those above recorded as having been obtained at the Dental Hospital? The nitrogen may have contained a large admixture of oxygen, which would account for the length of time required to produce anæsthesia. If the gas was pure nitrogen it must have become largely mixed with air during the process of inhalation. Every anæsthetist knows how difficult it is, without a well-shaped and padded facepiece, to exclude air during the inhalation of any gas. It is certain, from our experience at the Dental Hospital, that no one could breathe pure nitrogen for even the shortest time mentioned in the Middlesex Hospital report. Even with a mixture of 5 per cent. of oxygen the longest period of inhalation required to produce unconsciousness at the Dental Hospital was 95 seconds.

It is not without interest to compare the results of the experiments at the Middlesex Hospital with those obtained by Sir Humphry Davy with nitrous oxide. The great chemist would, no doubt, ensure the purity of the gas, but inhaling it, as he and others did, through a wooden tube held between the lips while the nostrils were closed, he could not exclude atmospheric air.²⁸ The result was, as he says, that he could breathe the gas for rather more than four minutes, but never so long as five.²⁹ It scarcely need be said that it is as im-

possible to inhale pure nitrous oxide for four minutes as it is to inhale pure nitrogen for that long period.

It is very noteworthy that when care is taken to exclude atmospheric air during the inhalation of the two gases—nitrous oxide and pure nitrogen—the average time required to produce anæsthesia is practically the same; so when the prolongation of the inhalation of nitrous oxide by Sir H. Davy and of nitrogen at the Middlesex affords conclusive proof that air or oxygen has not been excluded the increase of time required to produce anæsthesia was nearly equal in the two sets of cases. If Sir Humphry Davy could have inhaled, or seen others inhale, nitrous oxide unmixed with atmospheric air his account of the physiological effects would have been very different from that which he published at the beginning of the century, and the general use of the gas as an anæsthetic might not have been so long delayed.

I have proved that the inhalation of nitrogen, either pure or mixed with a small proportion of oxygen, rapidly causes anæsthesia not distinguishable from that which results from the inhalation of nitrous oxide. It now remains for anyone who still believes that azotic nitrous oxide, apart from its power of displacing the vitalising oxygen, exerts a specific anæsthetic influence to set forth the reasons for his belief; and in doing this he will, of course, bear in mind that it is worse than useless, it is seriously misleading to institute a comparison between nitrous oxide anæsthesia and the results of mechanical closure of the trachea.

THE THEORY OF CHOLERAIC COLLAPSE.

The right interpretation of the phenomena of apnoea, whether a result of the mechanical exclusion of air from the lungs or of the inspiration of an azotic gas, such as nitrous oxide or nitrogen, affords much assistance towards the solution of some most interesting and important pathological problems, and amongst these the pathology of the collapse stage of cholera is by far the most interesting. The condition of the heart's cavities which I have before described as occurring in the final stage of apnoea is identical with that which is found after death in the collapse stage of cholera.

We are indebted to the late Dr. Edmund Parkes³⁰ for the first accurate description of the condition of the heart and lungs after death in the stage of collapse. The following is a brief summary of the appearances, which he describes at great length. "The right side of the heart and the pulmonary arteries were generally filled, and in some cases distended with blood; the left side and the aorta were generally empty or contained only a small quantity of dark blood. The inference that was drawn from the state of the cavities in the greater number of cases was that the right side had continued to receive blood till, in some cases, it became full and even distended, while the left side received little or no blood, but had continued to contract upon the last drop of blood which had entered it.....In the lungs there was a remarkable want of air (that is, after the chest was opened); in the most rapid cases a want also of blood in the minute texture, causing the lung to collapse when the chest is opened and its weight to be diminished in a very great degree.....The conditions of the heart and lungs seem to point out unequivocally that in cholera the blood does not pass (freely) through the lungs." The only explanation of the condition which Dr. Parkes could suggest was that "some change in the fibrine" of the blood caused an impediment or arrest in the pulmonary capillaries (p. 113).

Dr. Parkes's book was published in 1847, some years therefore before the researches of Bernard, Brown-Séquard, and others had made known the function of the vasomotor nerves and the arterioles.

In a book on *Epidemic Diarrhœa and Cholera* which I published in 1855, I expressed my belief that the arrest of the pulmonary circulation is a result of contraction of the arterioles excited by the poisoned blood, and I adduced many facts and arguments in support of that theory. The condition of the heart and lungs is identical with that after death by apnoea, the difference in the order of events being this, that in cholera there is a primary asphyxia, that is, an arrest of the blood before it has reached the pulmonary capillaries, and a secondary apnoea, the result of the blood not being brought into contact with the air in the cells of the lung; on the other hand, when death is caused by privation of air

there is a primary apnoea. The blood which enters the capillaries not meeting with atmospheric air in the pulmonary cells, asphyxia—arrested circulation—is a secondary result. The term "cholera asphyxia" therefore, which was used by Scot, Bell, and other writers in India, was more strictly accurate than its authors were aware of. They no doubt meant to convey the idea that cholera was a suffocative disease, but the physiology of their time would not enable them to see the precise manner in which the aëration of the blood was prevented.

That Dr. Parkes was ready to accept my explanation of the phenomena which he was the first to describe appears from the following sentences in a letter which I received from him in November, 1865. He had been reading some papers of mine which were published in the *BRITISH MEDICAL JOURNAL*, and he wrote: "I feel pretty sure that you are right in attaching great importance to the difficulty of passage of blood through the lungs, which I always fancied depended on some condition of the blood itself, but I see that you attribute great effect to contraction of the arteries. This is certainly a very important point to work out, and seems to have evidence in its favour, at any rate that it is a co-operating cause. From what I have seen of your papers it seems to me that they will be a very important addition to our knowledge of cholera."

The pathology of cholera is too large a subject to be treated in detail in this paper. In the chapter "On the Pathology and Treatment of Cholera," in my *Medical Lectures and Essays*, the whole subject is fully discussed, and all reasonable and unreasonable objections to my doctrines have been, I venture to say, fairly met and refuted. In that chapter (p. 124) I have referred to the effect of the subcutaneous injection of the sulphate of atropine in doses of from $\frac{1}{5}$ to $\frac{3}{5}$ of a grain, as described by Dr. Saunders in the *American Practitioner* (July, 1873). "In some cases the relief afforded was astonishing; the skin grew warm, the pulse rose, the surface previously shrivelled and clammy assumed its natural condition, and in some instances the patient slept soundly for several hours." The explanation is to be found in the removal of the impediment to the pulmonary circulation by the paralysing effect of the atropine on the arterioles. In the same chapter I have shown that the marvellous temporary effect of a hot saline injection into a vein is to be explained partly by the dilution of the poisoned blood, but mainly by the high temperature of the injected liquor causing relaxation of the pulmonary arterioles. The cases recorded by Indian practitioners, in which patients in a state of almost pulseless collapse have obtained astonishing relief by venesection, are explicable by the disengagement of the over-distended right cavities of the heart. The condition of the heart's cavities is the same as in the final stage of apnoea, in which, as we have before seen, venesection increases the contractile power of the heart, or even restores its contractions after they have ceased.

The time is not far distant when it will be universally acknowledged that the hypothesis which attributes the collapse stage of cholera to the drain of liquid from the blood (the dehydration theory) is inconsistent with the facts, while in its application to practice it has been one of the most disastrous that ever gained general acceptance.

VARIATIONS OF THE PULSE; AND THE CARDIO-VASCULAR CHANGES IN BRIGHT'S DISEASE.

The two physical forces upon which mainly depend the volume and strength of the pulse are the contraction of the left ventricle at one extremity and that of the arterioles at the other. But another important factor is that of the volume of blood in the systemic arteries. For instance, the small and feeble pulse during the collapse stage of cholera is a result of the scanty stream of blood which passes through the lungs to the left side of the heart, the ventricle being also weakened by the defective supply of blood to the coronary arteries. The rapid changes in the character of the pulse during the successive stages of the various forms of apnoea I have already described and explained.

When, as a result of disease of the respiratory organs, imperfectly aërated blood enters the systemic arteries, an unwary practitioner may misinterpret the full and strong pulse

as a favourable sign, whereas it is a result of increasing de-oxydation of the blood.

Very valuable information may be obtained from an intelligent observation of the pulse in the various forms and stages of Bright's disease. In cases of acute nephritis with a scanty secretion of urine and consequent uræmia, the pulse is usually full and tense, a result of contraction of the systemic arterioles excited by the impure blood. The fulness and tension pass away when the excretory function of the kidneys has been restored.

The long-continued uræmia which results from chronic degeneration of the kidneys, and especially from the contracted granular kidney, is associated with very remarkable changes in the circulatory system. The radial pulse is very full and tense, the large arteries are thickened and tortuous, and there are the physical signs of hypertrophy of the left ventricle of the heart. Some years since I was led to search for the cause of this condition by the following considerations: Hypertrophy of the left ventricle without disease of the valves or of the large arteries—a fact made known by Dr. Bright in the first volume of the *Guy's Hospital Reports*—is probably a result, as he suggested, of some impediment "in the minute subdivisions of the vascular system." Reasoning from analogy, I thought it probable that the impediment is caused by the contraction of the arterioles excited by the impure blood, and further, that their long-continued overaction would result in hypertrophy of their walls, corresponding with the hypertrophy of the left ventricle. This led me to search for and to find the hypertrophy which I had anticipated in the arterioles of every tissue that I examined—in the kidneys, intestines, skin, muscles, and pia mater. The prolonged overaction of the arterioles registers itself in a conspicuous hypertrophy of their muscular walls.³¹

No kymograph could afford a more certain indication of excessive contraction of the arterioles, with resulting high arterial tension.

My description of the thickened arterioles, after passing through the inevitable ordeal of misrepresentation and ridicule, is now admitted to have been correct. The change in question is an example of true hypertrophy—overgrowth of a tissue without change of texture—not to be confounded with lardaceous degeneration, the irregular thickening which results from any form of arteritis, or the so-called "hyalin-fibroid change." The tortuosity and thickening of the larger arteries in the advanced stage of chronic renal disease are results of the excessive strain to which they are subjected between the hypertrophied left ventricle and the resisting arterioles.

The coexistence of cardiac hypertrophy with high arterial tension is very significant of renal degeneration in an advanced stage.

Some writers appear to assume that high arterial tension implies an increase of pressure within the capillaries, but this is an erroneous assumption. In fact it is obvious that if the increased resistance caused by the contracting arterioles be not compensated by increased propulsion from the heart, the intracapillary pressure would be diminished, but it is probable that the opposing forces are so nicely balanced that the pressure in the capillaries is, in most cases, unchanged.

RAYNAUD'S DISEASE.

A remarkable form of arrested circulation, which was originally described by Dr. Maurice Raynaud in his treatise *De l'Asphyrie locale et de la Gangrène symétrique des Extrémités* (Paris, 1862) is, by general consent, attributed to excessive contraction of systemic arterioles in localised regions. Raynaud's original treatise and his later researches, which were published in the *Archives générales de Médecine*, 1874, vol. 1, have been translated by Dr. Thomas Barlow. This translation, together with a valuable appendix by the translator, forms part of a volume of selected monographs which was published in 1888 by the New Sydenham Society. Referring to that volume for the detailed history of this remarkable affection, I can now do no more than indicate very briefly its general characters.

The essential features of Raynaud's disease consist in a local arrest of the circulation, with consequent coldness, blueness, and often dry gangrene of extreme parts, such as

the fingers, the toes, the tip of the nose, and the ears. There is also a remarkable tendency to symmetry in the parts affected, so that the disease usually, though not quite constantly, implicates corresponding parts on the two sides. Thus the upper or lower limbs, or all four together, may be bilaterally affected. In some cases the local gangrene is associated with that form of recurring hæmaturia which is called intermittent hæmoglobinuria.

This affection differs from senile gangrene in the fact that there is no structural change discoverable in the walls of the arteries, and in a large proportion of cases the patients are below 30 years of age.

Referring to M. Claude Bernard's researches into the action of the muscular-walled arterioles, Dr. Raynaud says: "The physiology of the circulation has been enriched in late years by one of the most beautiful discoveries of the century." Yet, with a confusing inaccuracy of language, he occasionally speaks of contraction of the capillaries, when from the context it is evident that he is referring to the arterioles. I venture also to criticise his use of the terms "local syncope" and "local asphyxia." The arrested circulation, he says, "can be compared to nothing better than syncope, in which the action of the heart is momentarily suspended" (p. 30). But, surely, to compare the failure of circulation resulting from suspension of the action of the heart with an arrest which is acknowledged to be caused by arterial contraction is to confound phenomena which are essentially different. The proper term by which to designate the arrested circulation in Raynaud's disease is local asphyxia, a result of arterial contraction. Raynaud, however, referring to the presence of unoxxygenised venous blood in the implicated tissues, calls this "local asphyxia." The correct term is local apnoea.

The word "asphyxia" has become so closely associated in men's minds with the idea of suffocation that, in spite of its obvious etymological meaning, it is frequently misapplied even in scientific treatises. This misapplication has the further disadvantage of leaving no term by which to distinguish an arrest of the circulation and of the pulse from apnoea (suffocation).

I hope to have established my position that the influence of the vasomotor nerves and the arterioles forms a connecting link between the diverse pathological conditions which are the subject of this communication, but there are many other clinical and pathological facts to the interpretation of which a similar influence is believed to be more or less applicable. M. Vulpian³² in the second volume of his work before referred to, has given an admirable summary of many pathological conditions, apparently associated with vasomotor nervous influence.

NOTES AND REFERENCES.

- ¹⁹ *Anæsthetics and their Administration*, p. 114. ²⁰ *Op. cit.*, p. 106. ²¹ *Op. cit.*, p. 130. ²² *Op. cit.*, p. 269. ²³ *Op. cit.*, p. 51. ²⁴ See Dr. Hewitt's account of this administration. *Op. cit.*, p. 268. ²⁵ Dr. Hewitt (*op. cit.*, p. 106) gives the maximum time of inhalation of nitrous oxide in 40 cases as 70 secs., the minimum 25 secs., and the average 51 secs. ²⁶ See the *Life of John Snow, M.D.*, by Sir Benjamin W. Richardson, M.D., F.R.S., introductory to Dr. Snow's work on *Chloroform and Other Anæsthetics*, p. xiv. Also my lecture on the Pathology of Coma and Anæsthesia, *Medical Lectures and Essays*, p. 290. ²⁷ *BRITISH MEDICAL JOURNAL*, 1868, 1, p. 593. ²⁸ In cases where the form of the mouth prevented the lips from being accurately closed on the breathing tube he used a tin plate conical mouth-piece fixed to the cheeks and accurately adapted to the lips. It would be scarcely possible to make such an arrangement airtight. ²⁹ *Collected Works*, vol. iii. p. 273. ³⁰ *On Asiatic or Algide Cholera*. ³¹ See *Med. Chir. Trans.*, vol. li; and *Medical Lectures and Essays*, p. 694. ³² *Leçons sur l'Appareil Vasomoteur*.

THE Association of American Physicians will hold its ninth annual meeting at Washington on May 29th and three following days. The programme of the public meetings includes a number of subjects of general interest.

METEOROLOGY AND HYGIENE.—Mr. G. J. Symons, F.R.S., gave the first of a series of six lectures on Meteorology in relation to Hygiene at the Parkes Museum on April 23rd. The course has been arranged by the Royal Meteorological Society and the Sanitary Institute acting together, and is designed to serve as a practical introduction to the study of weather and climate in relation to health and disease. Mr. Symons dealt with the instruments and with the method of making and recording observations. It was illustrated by photographs shown by limelight and by specimens of the various forms of apparatus in use at a meteorological station.