with the urinary constituents. I do not presume to determine whether this condition is precisely that known under the designation uræmia. In ordinary uræmia the process of secretion in the kidney is interfered with, and the products which ought to be removed accumulate. In rupture of the bladder the products have been secreted, and may have undergone considerable chemical change in the process. The absorption of these altered products may well produce a different complex of symptoms from that of a typical uræmia. Perhaps it would be more convenient to speak of urinary poisoning as the condition in these cases.

In considering the cause of death the condition of the peritoneum is not to be ignored altogether. The presence of a fluid of a different constitution from that of the normal peritoneal fluid may well have an influence on the membrane, and may affect its nervous and vascular arrangements. The symptoms of the patient indicate this in many cases, so that the diagnosis of peritonitis may be made, as in our second case. This must, however, be a subsidiary element, as the experiments of Wegner seem to prove that large quantities of urine may be infused into the abdomen without producing any pronounced symptoms.

REFERENCES.

- ¹ *Rupture of the Urinary Bladder*, by Walter Rivington, F.R.C.S. Eng., London, 1884, p. 12. ² *Ferraton, Des Ruptures Intrapéritonéales de la Vessie*, Thèse de Paris, 1883. ³ *Loc. cit.*, p. 9. ⁴ Wegner, *Experimentelles zur Lehre von der Ovariectomie*, *Langenbeck's Arch.*, XX. ⁵ Grawitz, *Charité Annalen*, 1886, p. 809. ⁶ Waterhouse, *Experimentelle Untersuchungen über Peritonitis*, *Virch. Arch.*, 1890, Bd. 119, p. 342. ⁷ *Loc. cit.*, p. 42. ⁸ Hamilton, *BRITISH MEDICAL JOURNAL*, 1883, I, 1166. ⁹ Macewen, *Lancet*, 1873, I, 448.

A NOTE ON CHLOROSIS IN THE MALE.

By C. F. MARTIN, M.D.,

Assistant Demonstrator in Pathology in the McGill University, Montreal.

CHLOROSIS is almost universally regarded as a disease which, if not peculiar to women, is at least of importance merely as it concerns the female sex—an idea which is likewise borne out by the name so often applied to it, morbus virgineus. Many authors, indeed, deny utterly its existence in members of the male sex, while others, though recognising its rare occurrence in men, have regarded it as a condition present only in connection with other feminine attributes, and existing thus only in men whose general characteristics and constitution render them effeminate. Thus Eichhorst, while denying that chlorosis is exclusively confined to women, and insisting on its occasional occurrence in men, considers that male chlorotics are always slender individuals, of feminine build, and frequently engaged in effeminate occupations, such as tailoring, etc.

In a recent article in the *BRITISH MEDICAL JOURNAL* Dr. Lloyd Jones states that in a research covering more than eight years he has been unable to discover a single case of chlorosis in the male. From the article mentioned it is inferred that these observations were founded merely on estimations of the specific gravity of the blood as a basis for testing the chlorotic condition. While doubtless there may be a definite relation between these two features, nevertheless the distinct relative decrease in the hæmoglobin has, during these last years, been considered as the crucial test for chlorosis, and on this altered relationship—first, I believe, insisted upon by Gruber—Dr. Osler lays special stress.

Regarding then chlorosis as an essential anæmia whose main characteristics are a greenish sallow tint of the skin, accompanied by a definite relative diminution of the hæmoglobin in the blood, we may say that such statements as those of the authors above mentioned fail to coincide with observations made by us during the past few months in Montreal, and hence the mention of a few cases which have led to the conclusion that, whether or not a true chlorosis be at times present in males, it may at least be said that occasionally in men a condition is found very closely approximating the chlorotic condition.

In estimating the hæmoglobin, a Fleischl hæmometer was employed, the instrument having been previously thoroughly tested on normal subjects, and due allowance made for the slight inaccuracies found.

CASE I.—J. S., a doctor of medicine aged 24, had observed that for several months a gradually increasing pallor and sallowness had deve-

loped, accompanied by gastro-intestinal symptoms, of which the most prominent were prolonged and obstinate constipation, anorexia, and fulness after eating. In addition he suffered greatly from headaches and severe intercostal neuralgia. The patient was of a decidedly neurotic temperament, had a personal history of hard work and prolonged studying at night. The family history was good. On physical examination there was found no evidence of organic disease. Estimation of the blood however revealed on an average 5,300,000 red corpuscles, white cells 1 to 400, and hæmoglobin 68 per cent. Treatment by arsenic was recommended and carried out. The patient was greatly improved thereby. Inasmuch as he left town soon after, there was no possibility of making further investigations, though letters received indicate a complete restoration of health.

CASE II.—H. G., a merchant, aged 27, having a history of prolonged dyspeptic disorders, was observed to have a markedly anæmic appearance; he had epigastric pain and fulness after meals, flatulence and marked constipation. Mental depression, almost approaching true neurasthenia, was superadded. Neither the personal nor family history could afford any light on his condition. Physical examination showed a normal state of all the organs, while the estimation of the blood gave the following result: Red cells 5,300,000, white cells about 1 to 350, hæmoglobin 72 per cent., this being the average of several examinations. Under treatment with arsenic and iron rapid improvement resulted, and when examined about two months later the patient's general condition was practically normal, and the blood examination showed the hæmoglobin to be 96 per cent.

CASE III.—E. H., a medical student, aged about 25, was likewise examined, inasmuch as a distinct greenish sallow complexion gave some indication of abnormal blood condition. The patient was of a nervous disposition, but had no complaint other than constipation. Physical examination of the organs was negative. During the examination of the blood the patient became very faint, almost losing consciousness. Red corpuscles numbered 6,030,000, hæmoglobin 77 per cent.

CASE IV.—M. S. This case resembled in nearly all its features Case No. II. An average of several blood examinations revealed 4,800,000 red cells, hæmoglobin 77 per cent. A soft systolic murmur was held at the base.

In these various cases presented, a blood condition has been obtained indicating a disease identical with the chlorosis described by many authors, and inasmuch as no other cause can be discovered with which to associate that anæmia, there seems good reason to believe that in them we have found an ordinary chlorosis. Although in no case there has been an extreme diminution of the hæmoglobin, yet it may be presumed that, as in these, so in many men, from some cause as yet not fully ascertained, an anæmia may be found whose main characteristic is a diminution of the hæmoglobin.

While being unable to contribute in any way to the subject of its etiology, yet that the condition described does sometimes exist in a mild form in males is at least worthy of mention, and that it may be perhaps more frequent than is generally supposed seems borne out by the experiences we have here met with during the past few months.

A REMARKABLE CASE OF INTUSSUSCEPTION.

By E. M. H. O'CONNOR, L.R.C.P.I., ETC.,

Market Deeping, Lincolnshire.

On May 1st, 1893, I was called in to see a lad, aged 13 years. I found him to present all the symptoms of acute peritonitis. He had had two or three well-marked rigors. He complained of vomiting, retching, and intense shooting pain in the abdomen. He lay on his back, legs drawn up; the face looked anxious; tongue small, red, and dry; intense thirst; respirations 50 per minute; temperature 104°; pulse 138, hard and wiry. The abdomen was distended to a marked degree. He could not bear the slightest touch without calling out in agony.

I got the following history of the attack: Eight days previous to my visit the patient had had a severe wetting, and was put to bed when he got home, a warm drink being administered. He got up next morning apparently none the worse, and went about as usual for four days. His bowels, however, had not been moved since the wetting. A dose of castor oil was then administered, but, having no effect, the friends, after waiting two days longer, gave him half an ounce of mag. sulph. This caused him a great deal of pain, so that several times "he cried out," but there was no evacuation.

On the evening before I was called in two "strong" purgative pills had been given, and I was informed that he had been much worse since he had taken them, and that the bowels had not moved.

I ordered turpentine stupes to be applied to the abdomen. The only food was to consist of milk in small quantities,