months, and differences in the numbers and formation of the bacilli noted and compared with cases not treated with creasote—this plan was however abandoned because the number of bacilli bore little or no relation to the patient's health, and no distinction could be drawn, either in number or morphological appearance, in the bacilli of those cases treated by creasote and those taking other drugs.

The next method adopted was that recommended by Professor Delépine, namely, that of injecting tuberculous sputum into the leg of the guinea-pig. A regular sequence of lesions is the consequence. The glands are affected in order, then the abdominal viscera, finally the lungs. At the end of about three weeks the animal succumbs. Such is the normal course of things. Anything that interferes with the virulence of the organism should interfere with the rate of growth of the bacillus, and the virulence of the bacillus can be gauged by the extent of the lesion produced.

To take the first series of cases (that is, those taking creasote by inhalation alone), injections into guinea-pigs were made before any treatment was begun, and again after the patient had been two months under treatment by creasote in this form. No difference was observed either in the time the guinea-pigs lived or in the extent of the lesion, in either case whether before or after treatment with creasote. Twelve such cases were tested with no result as to diminution of virulence.

In the second method of administration of the drug, that by the mouth, the results were much more satisfactory. Even in small doses (π ij, t. d.) after two months' treatment the guinea-pig lived seven weeks, instead of dying at the end of the third. A second specimen of sputum from the same case was injected into the leg of a guinea-pig, and the animal killed on the fifteenth day—this before creasote treatment was begun. All its organs were found affected with tuberculosis. A second inoculation made after two months' treatment gave the signs of tuberculous infection only in the popliteal, inguinal, and lumbar glands, where the animal was killed on the fifteenth day. As the dose of creasote administered internally was increased, so the virulence of the poison was proportionately diminished. Thus, in the case of a patient taking 6 minims of creasote three times a day, the guinea-pig lived nine weeks after the second inoculation; in another case in which the dose amounted to π x twice a day, the animal lived twelve weeks. Twelve cases were experimented on, and certainly in eleven of these it was abundantly proved that the greater the dose the greater was the diminution of the virulence of the poison; the animals, however, invariably died of tuberculosis in the end.

I have only three cases treated by the third method—that of the creasote chamber. In the first case, with inoculations made in the usual way, after treatment for two weeks by this method, the inoculated animals lived eleven weeks. A guinea-pig killed on the fifteenth day showed tuberculosis only at the seat of inoculation and in the popliteal and inguinal glands. In the second case, after two months' treatment by this method, the guinea-pig lived fifteen weeks. In the third case the animal died of septicæmia. Certain supplementary experiments were made. Two guinea-pigs were inoculated with tuberculous sputum in one leg. Into the other 10 minims of creasote were injected. This was followed by the injection of 5 minims twice a week. A third animal was used as a control experiment. One guinea-pig succumbed at the end of a week from cellulitis due to the creasote. Tubercle bacilli were found only at the seat of inoculation. The second pig lived sixteen days, and then succumbed to cellulitis. Here, again, bacilli were found at the seat of inoculation, and nowhere else. The control animal killed on the sixteenth day had extensive affection of the glands, spleen, and liver. These experiments were repeated twice with practically the same results, only that the doses of creasote were smaller, and the animals lived longer.

A guinea-pig was injected with tuberculous sputum, and the disease allowed to take its course up to the fifteenth day. Creasote was injected in doses of 5 minims every other day. In four such cases the drug had no effect whatever, the animals dying tuberculous about the twentieth day. If, however, creasote injections were begun at the end of seven days, it was possible (three cases) to keep the animal alive

for a month or six weeks; the cause of death was cellulitis from the creasote, the tuberculous lesions not having extended beyond the limits of the lumbar glands.

To sum up the conclusions arrived at:

(1) In the first series of cases—those that were taking creasote simply as an inhalation, in addition to such drugs as cod-liver oil and the hypophosphites—no effect on the virulence of the disease was noted.

(2) In the second series, in which creasote was administered by the mouth, in doses ranging from 2 to 12 minims three times a day, though, when the smaller doses were given the diminution of the virulence was slight, yet, when the larger amounts were reached, there was an extremely marked diminution in virulence.

(3) In the third series it may be pointed out that, though it is impossible to dogmatise from such a small number of cases, the animals lived longer than in any other instance.

(4) Creasote injected under the skin in tuberculous guinea-pigs, provided the disease were not too far advanced, had a markedly restraining effect on the poison.

In conclusion, I must express my indebtedness to the staff of the Victoria Park Hospital for their kindness in allowing me to use material taken from their cases in the wards, and especially to Dr. Arnold Chaplin for the use of tuberculous sputum from his patients treated in the creasote chamber.

The PRESIDENT (Dr. Woodhead) agreed that it might be that the virulence was attenuated. Professor Delépine had made similar observations, and Dr. Wood and he had also obtained similar results, but using heat instead of creasote as the attenuating agent.

SECTION OF DERMATOLOGY.

A. J. HARRISON, M.B., President.

A DISCUSSION ON THE MANAGEMENT OF ECZEMA.

I.—MALCOLM MORRIS, F.R.C.S. Ed.,

Surgeon Department of Diseases of the Skin, St. Mary's Hospital.

In opening a discussion, I take it that what is required of the person to whom that honourable task is entrusted is not an exhaustive address on the subject proposed, but rather the suggestion of a few questions to set the wheels of debate in motion. I must in the first place congratulate the Secretaries of the Section on the choice of a subject so far removed from what may be called therapeutic sensationalism, and so practical, as the management of eczema. It has been said that every true-born Briton thinks the capacity of dealing with the most obscure problems of theology a part of his intellectual birthright. In much the same spirit every medical fledgling just escaped from the academic nest will undertake the management of a case of eczema with a light heart. Experience may be trusted to teach him a better sense of our limitations in this direction, for of eczema it may be truly said that familiarity breeds the very opposite of contempt in those who have to treat it.

The time at my disposal will allow only of the discussion of principles, not of details—still less of reference to the methods and results of others. I propose therefore, in as summary a manner as possible, to present the general results of my own experience in the treatment of eczema, inviting special attention to the points as to which a comparison of experience seems to be most desirable. If the conclusions which I have to submit to you have little of novelty, they are at any rate the outcome of a tolerably large experience, and represent what appears to me to be the good therapeutic grain left after a pretty thorough sifting of the numerous remedies which have at one time or another been recommended for eczema.

As rational treatment must be based on a true conception of the process which has to be dealt with, a preliminary word or two as to the pathology of eczema may not be out of place. Opinions still differ as to whether the disease is constitutional or local in origin. My own view is that the

constitutional element is in most cases altogether secondary, the lesion being chiefly the result of parasitic irritation. A constitutional dyscrasia—such as gout, glycosuria, kidney disease, rheumatism, rickets, or scrofula—no doubt often has a serious modifying influence on the local process, intensifying it, and preventing or retarding recovery, but the *causa causans* is, I believe, in the majority of cases the action of micro-organisms. I am also of opinion that seborrhoea is a very common precursor of eczema, for which, in fact, it prepares the way by making the skin vulnerable by microbes. For a demonstration of the close relationship which exists between seborrhoea and eczema we are largely indebted to Unna, but I think that distinguished dermatologist goes too far when he teaches that eczema is always and in all cases the result of microbic action.¹ That seborrhoeic eczema is always parasitic in origin I freely admit, but I do not believe that all eczemas are seborrhoeic. I have often seen the disease arise on perfectly healthy skin, apparently as the result of some disturbance of the nervous system, caused, it may be, by mental shock, strain, or depression, by the action of cold, or by reflex irritation from the uterus, stomach, intestine, etc. Even in cases in which disordered innervation seems to be the starting point, I think it probable that the parasitic factor plays a leading part in the process; the weakening or suspension of nervous control—in other words, the inhibition of trophic influence, in certain parts of the skin creating weak points through which the micro-organisms normally present on the surface of the skin can penetrate into the deeper layers.

Whatever differences of opinion there may be as to the pathogeny of the disease, all will agree that the management of eczema must in every case be carefully adapted to the special indications present. There is no general therapeutic formula applicable to all cases of eczema; it is not the abstract name but the concrete disease that has to be treated, and not the disease alone, but the patient, with his peculiarities of structure and constitution, behind it.

Before touching on the points that appear to me to be open to discussion, I think it well to clear the ground of a question which, though fundamental in its practical importance, is to my mind so far settled as to be outside the domain of controversy. I assume that in all cases eczematous lesions should be cured as rapidly and as thoroughly as the therapeutic resources at our command will allow. Some appear to look upon a patch of eczema as a kind of overflow pipe for the escape of "depraved humours," which would otherwise—to borrow the language of popular pathology—"settle on" the lungs or other internal organs. I confess I looked upon this as an absolute superstition till I found that the notion still survives in quarters not open to the least suspicion of scientific heterodoxy. All I can say is that it is absolutely opposed to my own experience. In every case of eczema that comes before me I do my best to heal the disease as quickly as I can, and of the numerous cases which I have treated in not one have I seen the slightest ill-effect follow the cure of eczematous lesions.

I. The first point which I propose for your consideration is: *Are internal remedies required in eczema? If they are, what are the precise indications which should guide us in the use of them?*

My own view, stated broadly, is that as a general rule the less internal medication there is in eczema the better. If a constitutional dyscrasia underlie the skin affection it must of course be treated on the ordinary principles of medicine; if the neurotic element be pronounced remedies directed to the restoration of the nervous tone are obviously indicated; if the patient be broken down by want of sleep and the exhaustion consequent upon prolonged suffering, sedatives and general tonics will be required; similarly if there be disorder of the digestive apparatus this must be rectified by appropriate measures. Where, however, there are no definite indications of the kind that have been referred to, I am strongly of opinion that internal medication is not only useless, but positively injurious by disordering the digestion. In an ordinary chronic eczema, where the patient's health appears to be unaffected, and there is no reason to suspect constitutional dyscrasia, I trust entirely to local treat-

ment. Where internal remedies are given the selection of the particular drugs employed must be governed by certain definite indications. When the lesions are acutely inflammatory in character, I find antimony most valuable. I begin by giving πx to xij of the *vinum antimoniale*, repeating the dose in an hour, and, if need be, again two hours later. The interval between the administrations is gradually increased, the dose being at the same time reduced till one of μvj is reached. This should be given thrice in the twenty-four hours until a distinct subsidence in the intensity of the inflammation becomes manifest. The indication for antimony is the presence of arterial tension; on the other hand, depression is a positive contraindication of this drug. I am aware that a freer use of antimony has been recommended by some practitioners, but my own experience has not led me to see any necessity for larger doses than those which have been mentioned. If a neurotic element is clearly present in the case, sedatives and nerve tonics must be combined with local treatment. In such cases the sheet anchor is opium, under the influence of which the patient should be kept almost continuously until the nerve storm subsides. In such cases I have often found it of the greatest advantage to begin the treatment with a full dose of calomel, and afterwards the constipating effect of the opium should be counteracted by the administration of a saline aperient, such as Carlsbad salts, Friedrichshall, etc., every morning. If opium disagrees, some other sedative, such as sulphonal, must be substituted. The indication for the use of sedatives is great nervous excitement, accompanied by sleeplessness.

If, on the other hand, the neurotic element shows itself in the form of depression nerve tonics are indicated; of these the most useful on the whole in my hands has been quinine. This may advantageously be combined with opium. If there be much discharge belladonna may usefully be combined with the quinine. Phosphorus is another drug of great value in counteracting nervous depression in these cases, and strychnine also often does good. Arsenic, though undoubtedly useful in some cases, is far from being a specific in eczema, and if injudiciously used will do harm instead of good. The indication for its use is deficiency of nerve force combined with absence of acute inflammation in the lesions. If the eruption be of an actively inflammatory type arsenic is positively contra-indicated. If the disease process shows a marked tendency to frequent exacerbations, ergotin may be useful by the control which it exercises over the vasomotor apparatus.

Malnutrition, weakness, and anæmia are indications for cod-liver oil, and general tonic treatment on ordinary principles. Iron, however, is contraindicated by the presence of acute inflammation; in such cases it adds fuel to the flame. Of course, if any definite source of peripheral irritation is discovered it must, if possible, be removed. In women menstrual derangement or uterine disease must be remedied, and hysteria or the disturbances incident to the climacteric period must be combated by such remedies as musk, valerian, etc.

II. The second point which seems to me specially to call for discussion is the *influence of diet in eczema*. I may frankly state that my own view is that diet has no influence at all except indirectly. Thus, if a constitutional condition such as gout or diabetes underlie the eczematous process, of course the dietetic restrictions indicated in the circumstances are called for. Again, if the lesions are of an acutely inflammatory type, the diet should be limited in quantity, and bland and non-stimulating in quality. Apart from these indications, I have seen no reason to think that restrictions of diet had any good effect; on the contrary, I have seen a "lowering" regimen do positive harm by weakening the patient. The true principle in the management of eczema, as far as diet is concerned, is to let the patient take such food as he finds most suitable to his digestive powers. Anything, however, which disorders the gastro-intestinal tract, or which causes acidity, insomnia, flatulence, palpitation, or vasomotor disturbance must be avoided. The same principle applies to drinks; but here, it is almost needless to add, strict moderation in quantity is necessary. It is only in acute forms that beer is contraindicated, as are all other beverages likely to cause flushing. I place no restrictions on my patients in the use of tea or coffee unless these beverages are contra-

¹ On the Nature and Treatment of Eczema, *Brit. Journ. Derm.*, 1890, p. 231.

indicated by definite ill-effects on the digestion or nervous system. In short, as far as diet is concerned in eczema, I am disposed to say with Bacon in his essay *Of Regiment of Health*, "A man's owne observation, what he findes good of and what he findes hurt of, is the best physicke to preserve health."

III. The third question which I submit for discussion is, *What are the principles on which the local treatment of eczema should be carried out?* It is impossible here to go into the details of the local treatment of eczema of different forms and in different situations; this is the less necessary as my methods have been fully described elsewhere. Speaking generally, I treat every case as if it were of parasitic origin. Even if micro-organisms are not the exciting agents in the causation of the process in all cases, parasitic irritation always comes into play sooner or later as a secondary factor, and requires to be appropriately dealt with. The objects which I aim at in the local treatment of eczema are, first, to destroy micro-organisms; secondly, to protect the inflamed surface from the air and from further microbic invasion; thirdly, to soothe irritation. The general principles which guide me in the application of local remedies are, first, to temper the strength of the remedy to the tolerance of the patient's skin; and, secondly, to keep the remedy continuously in contact with the affected parts. I therefore begin with a very weak parasiticide, and feel my way towards the use of stronger applications, with the most watchful care not to cause any irritation. The best remedy for local use in dry, chronic eczema, especially of seborrhoeic origin, in my experience, is sulphur, and next to that I place resorcin. Both these drugs have this special advantage, that they not only destroy the micro-organisms on the surface of the skin, but give rise to exfoliation of the horny layer, and bring away with it the microbes which have penetrated to the deeper parts of the epidermis. I use at first an ointment composed of 10 grains of precipitated sulphur, or the same quantity of resorcin in 3j of zinc ointment or some other soothing application; the amount of the antiparasitic agent is increased very gradually. When the inflammation is acute, ichthyol is particularly useful, as it is a vascular sedative as well as a parasiticide. Other antiparasitic remedies which are of service as applications in eczema are salicylic acid, white precipitate, boracic and carbolic acid. I use them with cooling "creams" as vehicles when there is much irritation. For the continuous application of local remedies in limited areas of chronic character I have found nothing so useful as the plaster mulls devised by Unna, those I find most generally useful being carbolic acid and mercury, and oxide of zinc.

When discharge is profuse I wash the part with a weak solution of boracic acid, afterwards drying it with muslin bags containing starch and powdered boracic acid, or with flour mixed with a small quantity of the same antiseptic. If itching is troublesome, I find weak lotions of carbolic acid or tar most beneficial. For very persistent chronic eczema of the flexures I have frequently had good results from chrysarobin.

I may sum up the principles on which the local treatment of eczema should be conducted as follows: Soothe when the inflammatory process is acute; stimulate when it is chronic; and in either case keep the parts under the continuous influence of antiseptics and parasiticides, of a strength carefully regulated in accordance with the intensity of the disease and the sensitiveness of the skin.

IV. A fourth point on which discussion is likely to be useful is, *How is the tendency to recurrence of eczema in those predisposed to the disease to be overcome? Is there any plan of alternative treatment by which this result can be brought about?*

Change of climate is often productive of good in this direction, especially if accompanied by complete rest of mind as well as body. I know of no special indications in respect of climate except that eczema, being a catarrhal disease, it will be well to avoid climates which are apt to produce catarrh of any kind. A too bracing climate—especially places where the north-east wind holds sway—should be avoided. The same may be said of sea air. As regards altitude, there is no general rule; as in the case of climate, it is largely a matter of idiosyncrasy. The true principle is to discover what suits the patient. After all, it is the change that is the most

powerful alternative. Sea bathing very often produces a recurrence of the disease, but it sometimes undoubtedly counteracts the tendency to relapse.

As to waters, in estimating their usefulness, the effect of the change of scene and surroundings which a visit to a spa involves must be taken into account, as well as the chemical constitution of the springs. In my experience they are beneficial chiefly, if not so solely, in obstinate chronic cases. In selecting a spring for a particular case we have not much in the way of definite indications to guide us. The alternative influence of spas in eczema is a large subject, which in itself might well occupy the whole time allotted for this discussion. I can only touch on it here in the most cursory way, leaving it to others of large experience in this particular direction to give us the benefit of the knowledge so gained. On the whole, the sulphur springs—especially those of Harrogate and Strathpeffer—are best for external use; if there be a marked tendency to acute inflammation they are apt to aggravate the disease. The internal use of sulphur waters is particularly indicated in cases in which gout or rheumatism underlies the skin affection. Chalybeate and arsenical waters are useful in cases in which general tonics are indicated. The St. Victor Spring at Royat is one of the richest both in iron and in arsenic, and is therefore particularly useful for eczema in anæmic patients. Of the purely arsenical waters, the most useful in eczema are those of La Bourboule and Levico. The alkaline waters of Vichy and Ems have not seemed to me to be of any particular use in cases of eczema even in gouty subjects. In many cases the waters of Bath do good. The springs which I have mentioned are those which in my own hands have proved of greater service in cases of eczema.

I may sum up my own conclusions as to the utility of springs in the following propositions:

1. No spring known to me has any specific action on eczema.
2. Such virtue as sulphur waters externally applied possess is due chiefly to the parasiticide action and partly to the temperature at which they are applied.
3. Such virtue as sulphur, chalybeate, or arsenical waters possess when taken internally is due to their action in quickening the processes of metabolism, increasing the number of red corpuscles in the blood, and giving tone to the nervous system.

II.—E. D. MAPOTHEE, M.D., F.R.C.S.I., London.

Dr. MAPOTHEE referred to the management of one form of eczema—the oozing, postvesicular, especially of the adult and the aged. Having always held that the uric acid diathesis was causative of this condition, he had used for the last two years that wondrous solvent of the urates—piperazin—with most satisfactory results. The only other drug employed was a blue pill about once weekly, which either acted as a cholagogue, or an eliminant of uric acid. Copious drinking of pure water, such as the distilled water sold under various trade names, promoted diuresis and relieved the constipation so usual. When carbonated it was more agreeable, and lithia was often a useful addition. Of dietary articles, fresh fruits and unboiled vegetables were of great use. As to local measures, he thought ablation of the part several times daily, and of the whole skin as by a sponge bath once a day, essential. In Dublin and Glasgow the supplied water was practically pure, holding some 4 grains of solids to the gallon, and in London distilled water was delivered at 6d. per gallon. Ablution was needed to remove the irritating and probably toxic materials which oozed out. Of course the water should be tepid, and the drying should be done with the greatest gentleness; but after a sponge bath a good rubbing of the safe regions was serviceable. The continuous application of water often brought out a rash, but this was substitutive of eczema, often curative, and at any rate, like drug rashes, would cease on the removal of the cause. To refer to one most common site—the leg—he usually applied a single layer of old linen and a cheese-cloth bandage, to be moistened and renewed at least twice daily. When the case was acute, horizontal rest was needed, but inactivity promoted the constitutional condition which he thought causative, and passive exercise in the open air was always admissible. In a few cases connected with varicose veins, rubber bandages were useful. For

the initial step—the removal of crusts—he had found spongipiline soaked in water, sometimes with a little soft soap dissolved in it, efficient. Although he was against sealing up the discharging surface, he acknowledged that for the night, to prevent itching, vaseline with a little ammoniated mercury, should be smeared over the surface.

III.—A. S. MYRTLE, M.D.,
Harrogate.

DR. MYRTLE congratulated Mr. Morris on his interesting address, and was pleased to find that his experience as regards the management of eczema was in accordance with his own. He had long felt that internal remedies were, in the majority of cases, not only superfluous but injurious. Every variety could be dealt with successfully by local measures, and most of these belonged to the coal-tar series. He also considered any strict system of diet uncalled for. He had been unable to trace any connection between gout, uric acid, and eczema, although when it appeared in combination with diathetic conditions treatment must necessarily be directed to lessen or remove such. Whilst agreeing so far with Mr. Morris Dr. Myrtle could not accept his proposition that eczema was caused by the presence of any bacillus or micro-organism whatever. He maintained that in the vast majority of cases its origin was neurotic. Somehow the delicate nerve filaments became irritated, lost their healthy tone; alteration in cutaneous cell life followed, and eczema was developed. A patient suffering from an acute attack of eczema got perfectly well, was dismissed cured, in a few hours or days he was as bad, if not worse, than ever; he attributed his relapse to a chill, error in diet, or bother of one kind or another. A gentleman who had suffered from eczema of a most virulent type in both hands, complicated with sluggish liver for years, got perfectly well under his care. Before leaving he dined with him. When dinner was half through his hands began to burn and itch, the skin became tense, brick red; in half an hour it was in a state of active exudation, and from the tips of the fingers to the wrists, the disease raged again. Children treated for scabies by the free inunction of sulphur ointment at night and soft soap and hot water next morning, quickly got rid of the itch, but no sooner was this cured than in some cases acute eczema appeared spreading from the fingers up the arms to the face and head. Was it to be supposed that whilst they annihilated the itch parasite they created myriads of microbes of another sort? About three weeks ago a gentleman covered with eczema from head to foot had consulted him. He had been in this plight for two months. He (the speaker) asked if he could assign any cause for the outbreak; his answer was "The telephone." Some time before eczema appeared, he put a telephone in his office. He was engaged in most important and very difficult calculations, and when his mind was deeply engaged, ring went the bell, away went his thoughts, and he believed this annoyance proved the exciting cause of his skin trouble. In very obstinate circumscribed eczema looking like lupus, he had found nothing act so satisfactorily as a solution potassa fusa. He painted the patches over with a 15, 30, or 45 grain solution to 1 ounce of distilled water; three or four applications were generally sufficient, but the pain for a short time was intense. He never attempted the sealing up of eczema during its exudative stage, and had seen much harm result from this method. Mr. Morris mentioned Harrogate, and he (the speaker) could truthfully add that he had seen the most splendid results from the proper use of its waters and baths in cases where every other plan of treatment had failed, even a protracted course at foreign springs. This was all the more likely to occur when eczema was complicated with hepatic and dyspeptic troubles.

IV.—R. L. BOWLES, M.D., F.R.C.P.,
London.

DR. BOWLES raised the question whether the cure of eczema by external remedies did not depend more upon the hydrocarbon vehicles in which the numerous remedies were mixed than upon the remedies which were credited with the cure. He was led to this suggestion by the successful results on very bad cases of chronic eczema, lichen ruber, and psoriasis treated with oleum deeline alone; this oil was thin and pene-

trating in its effects, and Dr. Bowles thought it possible that if eczema depended on a microbe, it might be destroyed by the oil depriving it of the oxygen necessary to its existence. He referred also to the successful use in many such cases of coloured lotions and ointments in the hands of the late Mr. Startin. These might have owed some of their benefits to exclusion of the irritating rays of light. In this view Dr. Unna appeared to agree with him.

V.—DR. UNNA.
Hamburg.

DR. UNNA said he showed in 1881 that the influence of sulphur in eczema might do harm as well as good. The value of his plasters had been called in question by Dr. Myrtle, but the cause of failure was often their improper use. The microbic theory of eczema had met with great opposition, but was steadily gaining ground in England as well as Germany. He had convinced himself of its truth by a research extending over many years. He claimed that ichthyol had a powerful effect on eczema when given internally.

VI.—STEPHEN MACKENZIE, M.D., F.R.C.P.,
Physician to the London Hospital.

DR. MACKENZIE agreed entirely with Mr. Malcolm Morris that eczema was in nearly all cases a local disease, and therefore to be treated by local measures only. He agreed also with him that diet was of little or no importance in eczema *quâ* eczema, but when there was dyspepsia or any other condition requiring treatment dieting might be desirable on that account. As regarded the principle of treatment, he believed strongly that the general principle was to soothe the inflammation and heal the surface. The next point was to render the skin aseptic. He attached greater importance to the first than the second. As regards the microbic theory of eczema, though he had been in the habit of disregarding it in practice whilst inclining to believe it, he was quite open to conviction by Dr. Unna that eczema was of parasitic origin.

VII.—HORACE JEFFERIES, M.R.C.S.,
Bolton.

MR. JEFFERIES cited the case of a child who developed acute eczema about the eruptive period of the first teeth, which, being cured by local and internal remedies, was followed by an epileptic fit, the first of which increased with age, and lasted till death at the age of 14. He asked the question. Could this eczema be regarded as a nervous explosion, and as such conservative, and was the rapid curing of the case in any way connected with the epilepsy?

VIII.—FREDERICK PEARSE, F.R.C.S.,
Southsea.

MR. PEARSE said that a principle of treatment which had not been mentioned in the discussion was that based on the fact that alkaline applications would check alkaline secretions, and acid applications acid secretions. While some cases of apparent eczema would only recover under some form of parasitocides, a large number were relieved and cured by alkaline applications. In acute cases with copious weeping surface, very diluted alkaline solutions were to be recommended, the strength being gradually increased according to the less acuteness or chronic nature of the affection.

IX.—JOHN ROBERTS, M.D.,
Chester.

DR. ROBERTS observed that ointments frequently aggravated both the pain and the inflammatory condition. Fomentations were most useful, and soothing in cases of acute local eczema. A weak solution of potassa fusa was the best remedy he had used in chronic eczema of the ears. Oleum deeline was most useful in chronic cases of eczema, but should never be applied during the early stages.

ACCORDING to the *Australasian Medical Gazette*, 131 coroners' inquests have been held on infants in Sydney and its suburbs during the last five years, nearly all of who were illegitimate or "not known." In the great majority of cases the verdict returned was "Found Dead: Wilful Murder by Persons Unknown."