is elaborated which is transmitted to the blood. And here we are reminded of certain appearances formerly alluded to occurring in lower forms, notably the case of the myxine in which secretion particles seemed to be passed into the blood. In the case of the human subject, should this obser-

Fig. 8.—Commencing cirrhosis of medulla; hæmatoxylin and eosin; shows infiltration of leucocytes in clusters and branches; gland tissue undergoing atrophy. $\times 120$.

vation be justified, it can hardly be a matter of speculation as to the importance of the secretion so formed to the animal economy; more especially, it would minister to the blood and have a far reaching effect on the composition of that fluid, imparting to it essential qualities and so enabling it possibly to neutralise and render inert poisonous substances, a conclusion which the examination of the blood of the adrenal vein would seem to indicate. That this substance has also, directly or indirectly, the control of normal colour within certain limits is likewise a legitimate consideration. And, finally, its arrest would no doubt form one of the principal etiological factors in Addison's disease, in which the blood clots with difficulty and the red corpuscles refuse to form rouleaux.

EXPERIMENTAL RESULTS: CONCLUSIONS.

Stilling¹² has found that extirpation of one adrenal is followed by compensatory hypertrophy of the other. He also found that extirpation of the adult organs was followed by a hypertrophy of "accessory" suprarenals situated in the neighbourhood of the sympathetic ganglia. The possibility of such a compensatory action must be considered in cases of Addison's disease.

Tizzoni's¹³ well-known experiments on the effects of extirpation of the gland go to show that a morbid alteration in the blood is at the bottom of the cachexia. Bronzing of skin occurs in rabbits, and vascular lesions of great intensity occur in the central nervous system, consisting in injection and hæmorrhage into the grey matter of the cord and exudation into the ventricles of the brain. Branched villous outgrowths with great leucocytic infiltration are likewise found in the latter situation. Latterly areas of softening occur throughout both the brain and cord. Tizzoni regards these lesions as due primarily to a morbid alteration in the blood, affecting the composition of the cerebro-spinal fluid. An analogy is found in the vessels of the cutis, in many branches

of which thrombi have been found by Riehl and other observers in cases of morbus Addisonii.

The thyroid gland is an appendage of the respiratory system, and the study of its evolution shows that its original function is essentially concerned in the gaseous interchanges of the blood and tissues, "the phenomena following its destruction by disease or experiment being at bottom a disturbance of the gaseous metabolism and mal-assimilation of oxygen by the bodily tissues" (Andriezen). The pituitary gland is an appendage of the central nervous system, and, as the same observer points out,¹⁴ it is originally engaged in providing a secretion carried by oxygen-bearing water, which passes through its duct to the central nervous system, while its destruction is followed by a nutritional failure in the nerve centres. Of the suprarenal gland it may in like manner be said that it is an appendage of the renal-vascular system, being at the same time also in intimate connection with portions of the nervous system. Its functions, so far as the present investigation indicates, are (1) the destruction of certain effete products of metabolism which are of the nature of ptomaines, and (2) the elaboration of a secretion which is absolutely essential to the blood (hæmopoietic). Its destruction is followed by an auto-intoxication and a profound alteration in the chemistry of the blood, which entails, amongst other things, degenerative or nutritional changes in the nervous and digestive systems, and derangement of the the colour-regulating metabolism.

[I have again to express my best thanks to Dr. Thos. Reid, who has executed some of the photographs.]

THE BEAT-TRANSFER: AN EASY TEST FOR TRUE AND SPURIOUS PULSE TENSION.

By WILLIAM EWART, M.D.CANTAB., F.R.C.P.,
Physician to St. George's Hospital, and to the Belgrave Hospital for Children.

In cases where free communication is kept up between the radial and the ulnar pulses through the palmar arch, the strong pulsation which continues to be felt even when heavy pressure is made on the radial artery by the finger, is apt to convey an erroneous impression that the vessel still resists compression at a time when its channel has already suffered complete obliteration. The usual means for avoiding this mistake is to stop the anastomotic wave by the pressure of a finger of the other hand applied at a point nearer the wrist. A much simpler method is the one which I have recently described, and which is carried out with a single finger applied to the pulse in the ordinary way.

Illustrating the two positions of the finger applied to the pulse: I, pointing to the hand; II, pointing to the elbow of the subject; R, the head of the radius.

The method will be best understood by trying the experiments indicated in the figure. This mode of demonstration first occurred to me a few days ago (too late for insertion in my book on *The Pulse Sensations and Tactile Sphygmology*). It was so successful in making matters plain to others that its publication may be of use.

1. If the forefinger (the nail of which must be short) be firmly applied to any radial pulse with the position I, a strong beat will be felt by the pulp of the finger, which faces towards the subject's elbow, whilst no pulsation will be perceived by the

¹² *Revue de Médecine*, Paris, October, 1891.

¹³ *Arch. Ital. di Biol.*, Turin, 1894, v. n. 3 vi and 1888 x

¹⁴ *BRITISH MEDICAL JOURNAL*, January 13th, 1894.

extreme finger tip immediately below the nail, this portion of the finger facing towards the palm.

2. The position of the finger is then reversed to II, heavy pressure being this time avoided. The beat will be found to have changed sides, and to have migrated from the proximal to the distal side of the finger. Free pulsation is still felt by the finger pad in its new position, and very little or no pulsation is experienced by the finger tip.

3. As soon, however, as pressure is increased, a fresh change occurs—the distal beat disappears, and the tip of the finger (pointing towards the elbow) becomes conscious of a strong pulsation on its proximal side. This shifting or transfer of the beat is the sign which can be used as a reliable test that the artery has been successfully obliterated. The method, of which this is merely a practical demonstration, consists in keeping a single finger applied to the pulse in the usual position, and in gradually increasing its pressure till the change in question is brought about. The mechanism of this striking phenomenon, which is quite independent of the anastomotic pulsation properly so-called, is fully described elsewhere, and cannot be discussed in this brief communication.

OPENING OF THE MEDICAL SCHOOLS : INTRODUCTORY ADDRESSES.

ST. GEORGE'S HOSPITAL.

MENTAL TRAINING IN MEDICAL STUDY.

By ISAMBARD OWEN, M.D.,
Dean of the Medical School.

GENTLEMEN.—The honour of greeting you at the opening of a new academic year has fallen this time to me, and custom demands that I should begin with some brief reference to the changes that have taken place among us during the year that is past.

[After a review of the leading events of the year in connection with the hospital and medical school, Dr. Owen proceeded:]—

To some of you who are here, to-day will mark an epoch in your educational career. You will for the first time find yourselves working not merely to win academic renown, but to fit yourselves for the discharge of serious and responsible duties. It may even be that for the first time you will find yourselves turning from the study of abstractions to the consideration of material things. To you in particular tradition assigns me the flattering but puzzling task of addressing a few words of counsel on the subject of your future duties and the qualifications needed for performing them. A puzzling task, because the subject is so vast and all its parts so closely connected that it is difficult to select any one for separate treatment. But time being short and occasion urgent, I will pass without comment over the moral qualities—earnestness, steadfastness, courtesy, and probity—which medical life demands, and over the physical discipline required to fit yourselves for the marvellous handiwork of modern operative surgery, and endeavour to speak of some matters within my more immediate cognisance connected with the mental training that goes to the making of a physician.

I speak advisedly of the physician's training, because, above all, I wish you to bear in mind that training as well as learning is a necessary part of medical education. You are not preparing yourselves only to pass examinations or in prospect of a life of lettered ease. You have to do as well as to know; and medical doing, like all other doing, requires a special cultivation of the faculties by which it must needs be exercised. Mere learning, however profound, can no more enable you to practise medicine than to command an army in the field.

I will ask you in passing to accept the word "physician" in a wider sense than that to which ordinary English custom restricts it. The line that once separated physicians and surgeons has now no existence in law, and is limited to the special sphere of consulting practice. The two professions have been united in one, and every candidate is required to possess the qualifications of both. But Parliament, in con-

solidating the profession of medicine, unaccountably omitted to accord its members any common title by which they may be generally known. The law knows us under the cumbrous style of "registered medical practitioners"; in the services we are "surgeons"; to the newspapers "medical men," or, sometimes, "medical gentlemen"; and the general public, taking refuge in a piece of time-honoured slang, persistently calls us "doctors." In the United States common consent has extended a more general term to cover a more special one; and the sensible American habit of calling medical practitioners generically "physicians" is one I think we should do well to borrow.

The practice of physic, in the eyes of the public, consists in the application of remedies to the cure or relief of disease. Though this is indeed our ultimate aim, the actual application of the remedies bears some such relation to the entirety of the physician's work as the visible part of a modern battleship does to the mass of metal floating under water. You will not proceed far in your studies before you discover that nine-tenths of his task is most often done before he utters a word of advice or writes the first line of a prescription; and that the power of discriminating diseased conditions, to the casual eye a simple and easy matter, is in reality the aim of the greater part of the lengthy course of study you have to go through.

Diseases are disturbances of organs of the body. The organs of the body are many, the modes and causes of disturbance at least equally numerous. There are parasites from without and new growths from within, inflammations and structural changes, and countless derangements of function of less tangible origin, but not always of less serious import. Few of these disordered conditions can be the subject of direct observation. The organs they affect are for the most part concealed from actual view, and direct exploration is a measure that can be used only in the gravest emergencies. We have for the most part to form our judgment from a combination of observations of a difficult and delicate character, observations of subjective phenomena made through the medium of the patient's mind and our own skill in questioning, observations of minute changes in the condition of the visible surface, observations of involutions of the surface by means of optical apparatus, observations of the results of physical, chemical, and physiological tests; microscopic observations of disrupted fragments, and observations of a series of fine acoustic and tactile phenomena and reactions elicited and appreciated by means of the trained fingers or various artificial appliances.

To but few observed phenomena, subjective or objective, does a single value attach. It is only by a consideration of their varying combinations that we can arrive at an opinion as to their cause, or tell with certainty if the patient's state be grave or trivial, one of temporary disturbance, specific infection, or incurable organic change. But until such a judgment is formed, to apply a remedy is to cast a stone at a venture. Rational treatment can only be based upon the exact "diagnosis," as our esoteric term has it, of disease. His observations made, the physician has therefore to frame such a theory of the case, or theories, if more than one be possible, as, in the light of his accumulated knowledge, shall accord with and explain the entire body of the facts. He must verify his theory, or discriminate between them, if there be more than one, by such further present or subsequent observations as may be feasible, and his final conclusion he ultimately brings to a practical test, by laying down a plan of treatment and abiding the result. Such, in broad outline, is the physician's method. It is not peculiar to medicine; it is essentially the ordinary method of physical investigation. Between the science and the art of medicine there is no hard and fast line, for the art mainly consists in the application of scientific method to a particular set of problems, and the science is but a wider application of the art.

I am anxious that you should attach some definite meaning to the words "science" and "scientific," which will encounter you like haunting spirits at every stage of your career; spirits of good, if rationally understood; if not, will-o'-the-wisps to lure you astray. Around these words has gathered in our day a certain halo of intellectual grandeur, so much that the mere use of them in an argument creates in