OSTEOPOROSIS OF THE CRANIAL VAULT.

By GEORGE WHERRY, M.C.CANTAB., F.R.C.S.

University Lecturer in Surgery, Cambridge.

EXCEPT in cases of osteitis deformans, there have been no clinical records of symmetrically-growing heads in adults such as might furnish examples of osteoporosis. The thickened calvaria of our museums are usually without history, and though Sir James Paget, by his epoch-making account of osteitis deformans has enabled some of these to be classified, there remain many gaps in our knowledge which can only be filled by clinical accounts.

The accompanying figure is from the photograph of a carpenter, aged 56, who came under my care in Addenbrooke's Hospital, suffering from fracture of the lower third of the tibia. The bone was broken in two places, about 3 inches apart, with a rounded swelling between the breaks, the fibula being sound. The swelling is tender rather than painful. He had been laid up several months with a "rheumatic" knee, and three months ago broke the leg in getting off a night stool, and has borne no weight upon it since.

There is evidently a sarcomatous growth in the bone, resulting in the fractures from such an inadequate cause.

There is well-marked evidence of arthritis deformans in both knee-joints, the lipping of the articular ends of the bones being characteristic. There is no thickening or curving of the shafts of any long bones in his body.

The spine is bent and stiff in the dorsal region, but the neck and loins are fairly free from stiffness.

The head is large, but retains its symmetry, and merely gives the man an intellectual appearance, without any deformity. It measures above the ears in circumference 26 inches, and has been growing for years; he thinks as many as ten years. His

hats have been obtained of increasing size, until they had to be made for him. His hatters—Golding and Son, Newmarket—write to me as follows:

We have supplied him with hats about three or four years, and during that time we have noticed that his head has been growing larger. The first hat or two that he had we were able to supply from our stock (size $7\frac{1}{2}$, the largest size we stock), and then the next he wanted $7\frac{3}{4}$, and the next $7\frac{1}{2}$, and we had then to get a $7\frac{3}{4}$ made for him; the last one he had from us was $7\frac{3}{4}$ full; this of course is a tremendous size, the largest we have come across in our experience. We are rather interested in this man ourselves, as we have often remarked that we thought he must have some disease of the head for it to grow to such an extent, especially as our experience is just the opposite; if heads vary at all with our customers we find that they decrease instead of increase.

Hatters' scale of sizes: $7\frac{1}{2}$, head measures 22 in.; $7\frac{3}{4}$, 22 in.; $7\frac{1}{2}$, 23 in.; $7\frac{3}{4}$, 24 in.; $7\frac{1}{2}$ full, 24 in.

This patient first came under my notice as an out-patient in the eye department in July, 1890. He was suffering from atrophy of the optic nerves apparently from neuritis. There was albumen in the urine, but there were no retinal hemorrhages. He was a considerable smoker, but there were not evidences of tobacco amblyopia. His tobacco was stopped, and he was treated with cod-liver oil, iron, and digitalis during twelve months. His vision improved in this period from counting fingers at 4' to V = $\frac{2}{3}$, and reading J 2. He has greyish-white atrophic discs. He has become very deaf of late from chronic catarrh of the tympanum. The drum-heads are easily seen, and there is no exostosis or bony deformity of the aural canal, but the optic nerve affection may have been due to alterations in the optic foramina.

He is now, and has always been mentally quite alert and intelligent, and, except for the infirmities mentioned, has had good health, and never suffered from headaches. He has been a cheerful well-occupied man all his life.

From the fact that unusually large heads required hats to be made for them, I was led to elicit from Messrs. Lincoln and Bennett, hatters, the following information on the subject, with a diagram showing a head of unusual size with no evidence of want of shape or symmetry; the circumference of this gentleman's head increased $3\frac{3}{4}$ inches in fourteen years. The final measurement was twenty-seven inches.

Measurements and dates were taken.

1870, Sept. 16th,	23 inches.
1872, July 1st,	23 $\frac{1}{2}$ "
1873, May 22nd,	24 $\frac{1}{2}$ full.
1874, Nov. 16th,	24 $\frac{1}{2}$ bare.
1875, Feb. 26th,	24 $\frac{1}{2}$ full.
1877, Jan. 22nd,	24 $\frac{1}{2}$ "
1878, April 10th,	24 $\frac{1}{2}$ "
1878, Sept. 18th,	24 $\frac{1}{2}$ full.
1880, July 5th,	24 $\frac{1}{2}$ "
1881, Jan. 10th,	25 $\frac{1}{2}$ "
1882, May 31st,	25 $\frac{3}{4}$ inches.
1884, March 26th,	25 $\frac{1}{2}$ "
" May 26th,	26 $\frac{1}{2}$ "
" Oct. 20th,	27 "

This gentleman died, aged about 70, soon after the last measurement. He used to walk with a stoop both of head and body, and always appeared glad to sit down. He was slightly lame in one leg. The remarkable increase in the

growth of the head at the end of life is noteworthy.

In another case of enlargement of the head, the gentleman stooped and walked heavily in the same way. He died aged about 80. His head, always large, began to grow in 1872, between which date and June, 1878 (shortly before he died), the increase in circumference was three-quarters of an inch, the actual size being $26\frac{1}{4}$ inches.

Both these are probably cases of osteitis deformans. Sir James Paget notes in his typical case that his patient wore a shako in 1844 measuring $22\frac{1}{2}$ inches inside; in 1876 his hat measured $27\frac{1}{4}$ inches inside; but "the changes in shape and size in both the limbs and the head were arrested, or increased only imperceptibly, in the last three or four years of

life," in decided contrast to the rate of growth as recorded by Messrs. Lincoln and Bennett.

At present, without more evidence, my case can hardly be pronounced one of osteitis deformans, and yet the enlarging head occurs in that disease, and is not found associated with arthritis deformans; also new growth is often found with osteitis deformans, as in several of Sir James Paget's case. In Mr. Lunn's case there was a sarcoma in the pelvis. The entire skeleton is in the St. Thomas's Museum, and is beautifully figured in *St. Thomas's Reports*, 1883, and described by Professor Stewart.

As rarefying osteitis is the chief feature of the disease in osteitis deformans, this would afford also the best explanation of the symmetrically enlarging cranial vault in my patient.

ABDOMINAL SARCOMA INFILTRATING THE ABDOMINAL WALL.

By J. WARD COUSINS, M.D., F.R.C.S.,

Senior Surgeon to the Royal Portsmouth Hospital, and the South-East Hants Eye and Ear Infirmary.

M. A., a married woman, aged 27, was admitted into the Royal Portsmouth Hospital during September, 1891. The

abdominal swelling had been noticed twelve months. Her general health had previously been good, and the catamenia

regular. On examination the surface of the abdomen appeared regular and full, and a well-marked and elastic tumour could be readily defined, which extended upwards 3 inches above the umbilicus. There was no trace of fluctuation. The pelvic cavity was free, the uterus was drawn upwards, and the os felt soft and healthy. The circumference at the umbilicus measured 32 inches, and the distance from the symphysis to the ensiform cartilage 17 inches. An exploratory operation was performed on October 1st. The abdomen was found filled with an omental tumour, which was intimately associated with the internal organs. Its surface was highly vascular. A small portion of the growth was removed, but microscopic examination revealed only normal omental structure.

Shortly after the patient left the hospital in fair health, and returned to her situation. She stated that the operation had given her much relief.

In July, 1893, she was admitted again under my care. The abdomen had greatly increased in size. The circumference at the umbilicus measured 38 inches, and midway between this line and the symphysis it reached 45 inches. She appeared much emaciated, and her features had the pinched expression which has often been described as the "facies uterina." For some months she had been unable to follow any occupation. The abdominal surface was deeply pigmented, especially around the scar, and the cicatrix was completely infiltrated with new growth, and measured 3 inches from side to side. In many places it presented a nodular appearance, and the infiltration had even penetrated through the suture scars, and from these pedunculated masses of new growth were protruding.

The accompanying photograph was taken at the hospital a few days since, and I am much indebted to my colleague Mr. Newby for his kind assistance.

MEMORANDA: MEDICAL, SURGICAL, OBSTETRICAL, THERAPEUTICAL, PATHOLOGICAL, Etc.

CASE OF VIPER BITE.¹

C.B., aged 16, while playing with a viper, was bitten in the ball of the thumb at 5.10 p.m. The wound was sucked immediately, ammonia applied, and a ligature tied tightly around the wrist.

He was seen by me at 6 p.m. The fang marks, about 1 inch apart, were distinct; the hand was much swollen below the seat of ligature, but the patient felt perfectly well. Three minims of liquor ammoniæ fortis, diluted, were injected at the seat of injury, and 2 drachms of sal volatile given internally. At 7 p.m. there was so much œdema that it was thought wise to loosen the ligature. The œdema spread rapidly up the arm to the shoulder. As the patient was now feeling faint, 2 ounces of brandy were administered in tea, and he felt better; the pulse was very irregular. At 8 p.m. he felt very sleepy, and said later that at this time he felt that he did not care what happened to him. The pupils were enormously dilated, the pulse most irregular—peculiarly so at times, 20 beats or more with extreme rapidity, then long intermission followed by a few slow beats. A teaspoonful of Wyeth's meat juice, and half a tumbler of champagne, were given. A few minutes later there were rigors, followed by vomiting about 9 p.m. Hot-water bottles were applied to the lower extremities, and the patient was put to bed, where he remained in a very drowsy condition, but the heart became more regular. By 8 p.m. a patch of blackened tissue, about 2 inches in diameter, was found at the seat of injury. About 11 p.m. there was a sharp attack of diarrhœa, and at 11.30 he was fairly comfortable.

At 2 a.m. his mother roused me, as he was rather restless and excited. The pulse was rapid, and he seemed feverish, but without any rise in temperature. The drowsiness had passed away. I now gave him 25 minims of chlorodyne, after which he had a good night's sleep, and in the morning felt

¹ Read at a meeting of the Isle of Wight District of the Southern Branch of the British Medical Association.