A MALARIA CHART.

[WITH COLOURED PLATE.]

By PATRICK MANSON, M.D., M.R.C.P., LL.D.,

Physician, Seamen's Hospital, Greenwich.

THE value of the temperature chart and of similar graphic methods of projecting clinical observations is universally admitted. Such methods greatly facilitate the recording, and still more the comprehension and comparison of the facts in the clinical tableau. The "malaria chart" is an attempt to apply the same principle, so far as the nature of the subject will permit, to the recording in cases of malarial fever of the types and varying phases of the malaria organism along with, and in comparison with, the fluctuations of body temperature. The scientific and practical value of such a chart is obvious. It is believed that after a little practice these combined records will be as easily made and read as those of temperature alone, and it is hoped that they will be of assistance in unravelling some of the many difficulties and obscurities at present connected with the malaria question.

So far as concerns the figures of the malaria parasite given in the coloured plate, the compiler lays no claim to originality. These are taken from the works of Laveran, Golgi, Marchiafava, Bignami, and Mannaberg, and were selected as being probably the best illustrations of the malaria parasite we possess.

In the plate the various forms of the malaria organism have been to a certain extent classified, grouped, and graduated according to their appearances, not necessarily according to their assumed biological relationship. This has been done in such a way that a figure may be easily found corresponding very closely, in type if not in detail, to any parasitic form which may be encountered in blood under examination. Provision by letter and number is made for rapidly and accurately indicating this on the chart, and afterwards of re-identifying it in the plate. These letters and numbers, written in their appropriate place on the temperature chart, constitute the record.

It may be objected to the plate that there is a useless multiplication of certain of the forms of the malaria organism. The compiler thinks, however, that in the present state of our knowledge about this parasite it would have been unwise to leave out unnecessarily, and at the risk of missing what may prove useful even in a small degree, one faithfully drawn figure. Should the observer, as is probable, desire to add to the number of figures, there are several unoccupied spaces which may be used for this purpose. The figures are classified as follows:

Columns A and B contain simple forms of unpigmented intra-corpuseular bodies; columns C and D, the ringed forms of intra-corpuseular bodies; E and F, principally the pigmented and larger forms of the latter; G and H, pigmented forms of the simple intra-corpuseular bodies; I and J, larger forms of the latter graduated according to size; K and L, intra-corpuseular, segmented sporulating forms; M, the latter free and breaking up; N, O, P, Q, include the crescent and flagellated forms and the bodies supposed to be connected with these, besides others which cannot be easily classified with the foregoing. The three figures at the foot of B column are somewhat diagrammatic representations of pigmented leucocytes; those at the bottom of D are extra-corpuseular forms, the central one, D 11, representing the "swarming" corpuscle; those at the bottom of F represent, according to Laveran, dead forms. L 11, M 8, 9, 10, represent free forms perhaps in a degenerated condition; M 11 and 12 are free parasites; and O 11 and 12, degenerated forms.

It is not supposed that under ordinary conditions of practice the medical observer could afford the time to make microscopical examinations of the blood every four or six hours, but he may be able, at all events in hospital practice, to make a daily or occasional examination. Such a series of fragmentary and more or less irregular examinations need not be wasted if systematically recorded in the "malaria chart."

DIRECTIONS.

Preparation and Examination of Blood for the Malarial Organism.—Cleanse very carefully with alcohol and ether several slips and thin cover glasses, and cover them up from the dust. Wash one of the patient's finger tips with soap and

water, and afterwards with ether, and dry carefully. Ligature the end of the finger in the usual way, and prick the congested pad with a fine, clean needle. Wipe off the first drop of blood which exudes, being careful to leave the skin quite dry, so that subsequent drops shall not "run." Squeezing the pricked finger pad gently between finger and thumb, express a second and smaller droplet of blood from the puncture. This droplet ought not to exceed in size the head of a large pin. Touch the apex of the droplet with the centre of a cover glass, and immediately lay this on a slip. The blood will now run out between slip and cover glass in an exceedingly delicate film, in which, after a few minutes, the red corpuscles will be found to be each of them perfectly isolated, and lying flat on their sides. Prepare several such slides. Reject all slides in which the corpuscles in any considerable proportion are disposed in rouleaux, or are heaped up upon each other.

Perfect cleanliness of finger and slides, minuteness of the droplet of blood, thinness of cover glass, and a certain quickness of manipulation, are the best guarantees for success in obtaining the flat disposition of the blood corpuscles *absolutely* indispensable.

Examine the slides so prepared with a twelfth immersion lens, and in not too bright an illumination. Scrutinise the interior of every corpuscle in the field, looking in them for specks of black pigment surrounded by a pale, hyaline, slightly or markedly amoeboid substance; also for smaller, pale, unpigmented, hyaline, and more actively amoeboid bodies in the same situation. These are the intra-corpuseular and commoner forms of the malaria parasite, and are always present in malarial fevers which have not been treated by quinine. The crescent and flagellated forms and the pigmented leucocyte—although the two former are more rarely encountered, and only in certain cases—are much more easily recognised.

If no parasitic form be found in the first field pass to a second, a third, and so on, devoting at least half an hour to the examination of the slide before pronouncing definitely in a negative sense on the presence of the parasite.

Registration of Malaria Organisms.—When a parasite or pigmented leucocyte is found search in the coloured plate for a corresponding form, and ascertain the letter and figure indicating the vertical and horizontal lines in which it occurs. Note these on the temperature chart (for example, thus—A 1, or B 7, or G 3, etc., as the case may be) in the highest compartment of the column corresponding to the day and hour at which the examination is made. This done, search for a second parasite; a second being found, identify, and note it also, in the same way on the second space of the same column, and so on until ten or more parasites have been found and recorded. It is important that every parasite found should be noted; there must be no selection of parasites, otherwise the proportion in which the different forms occur will not be accurately represented. A note of the number of fields examined should also be made in the space provided for the purpose at the head of the column.

It is recommended not to register more than fifteen or twenty consecutively-found parasites, as this number is sufficient to convey a fair idea of the proportions in which the prevailing forms occur. It is also recommended that that part of the column which is above the 98.4° line be devoted to the noting of these, space being saved by a tally mark of some sort below the registration of any form which is found two or more times. When an idea of the prevailing forms has been arrived at, and the types noted, the observer may proceed to hunt for rarer forms; these might be registered in the spaces below the 98.4° line in the appropriate column.

Temperature, etc., are to be recorded in the chart in the usual way.

In the absence of the special form of chart the ordinary temperature chart may be used for the registration of parasites.

The charts accompanying the plate are not divided into days, as is usual in temperature charts. This is left for the observer himself to do, so that he can adapt his charts to a four, six, or twelve-hour record, as may suit his purpose and convenience.

The object of registering the number of fields examined is to find out approximately the proportion of parasites to cor-

A. B. Aged 40.

TERTIAN; MALARIAL CACHEXIA.

puscles. This implies that the observer has already ascertained the number of corpuscles which in properly-prepared slides an average field of his particular microscope contains; and that in looking for the malaria organism he selects average fields for observation.

Great amoeboid activity of the parasites or other striking and unusual phenomena in connection with them can be indicated on the chart by appropriate signs.

CLUBS AND FRIENDLY SOCIETIES IN THE SOUTH OF IRELAND.

A VERY serious difference has arisen between the medical practitioners of Cork and the various clubs and friendly societies of that district. It would appear that the medical men in Cork have drafted a series of rules by which the tenure of club appointments is to be regulated, and that they have unanimously pledged themselves to accept such appointments under no other conditions.

The matter had been referred to the Medical Society and then to the South of Ireland Branch of the British Medical Association, and finally the entire profession unanimously adopted the present rules. These have been submitted by the Branch to the various friendly societies of the district, who have as unanimously repudiated them, so that a very critical state of affairs is likely to arise.

It would appear that the main complaints of the medical men arise from (1) the smallness of the fee at present paid in some of the societies; (2) the absence of a wage limit. But there are several other points on which differences have arisen. The first of the suggested rules runs as follows: Scale of remuneration to be in proportion to member's income, and, when this cannot be ascertained, to valuation of dwelling house. The second is: Minimum yearly fee 7s. 6d., if member's income is under £100 per annum, or valuation of dwelling house under £7 10s. per annum; minimum yearly fee, 15s., if member's income is under £200 per annum, or valuation of dwelling-house under £15 per annum. No one with an income above £200 per annum to be admitted. This scale is for medical attendance only. There are other rules regulating the maximum distance of attendance, fees for midwifery, operations, consultations, etc. The above scale of remuneration seems to include attendance upon wives and families so long as they are dependent on the member.

There seems little in these rules of which working men can properly complain. They seem, in fact, very moderate, and are such as might with advantage be adopted throughout the greater part of the United Kingdom. Few will be ready to believe that a working man in the receipt of a pound a week is unable to pay twopence per week for his medical attendance, and fewer, we should imagine, will contend that those earning higher wages ought not to pay proportionately higher rates. Operations, again, requiring anæsthetics, and midwifery have always been considered extras, so that it is difficult to see what there is unreasonable in the demands of the South of Ireland Branch of the Association. Probably the violent opposition which seems to have been aroused has been caused chiefly by fear engendered by the business-like and well-organised action of the profession in Cork. These associations have so long had their medical officers in complete subjection to them, that such evidence of a determination to put an end to this unseemly condition of affairs is doubly displeasing, especially when they dread very much the possibility of its success.

Surely it is high time to put an end to the present disgracefully low rate of club payments. When the cost of medical education is considered, and compared with what it was in the past, and when it is further noted that in spite of the enormous increase of wages during the last twenty years, the ordinary club rate of remuneration has been quite stationary, if it has not in some cases even retrograded, can any argue that some reform is not urgently needed? And as to the wage limit, it certainly is a scandal and disgrace that people in comfortable circumstances should be permitted to rush into these clubs and get medical attendance at such a rate.

We quite understand the feeling of the friendly societies, which seem to consider that any wage limit in regard to their

members would tend to cut out of their associations their best, their most thrifty, and their most successful members. It is no doubt a subject of satisfaction, and even pride, to retain among their members those who have got on well and risen to better positions, and it clearly is a matter of profit, seeing that these members rarely claim sick pay. If they rarely asked for the doctor's attendance, it would not matter, but unfortunately that is not the case. These better-class subscribers, it would appear, are a cause of a great part of the friction, and we cannot but feel that the most equitable compromise would be that while they should retain all other rights of membership, medical attendance should not be claimable by those whose income is above a certain limit. No exception can fairly be taken to the limit suggested. By this plan no member need ever be turned out of a club; the only change would be that, when sufficiently prosperous not to require it, medical attendance would cease.

On the whole, the scheme, as proposed by the Cork Branch, is an exceedingly moderate one, and in no way threatens, as loudly declaimed by its opponents, the existence of genuine friendly societies; nevertheless, there can be little doubt the Cork friendly societies will fight the matter to the end; and, to do this, will endeavour to attract other medical men from a distance to take the place of the local practitioners.

If this should be so, we would advise the friendly societies to think well before adopting so suicidal a course. Accustomed as they have been to the professional assistance of doctors of good standing, they probably fail to recognise the advantage which they derive from being attended by such men, and do not at once see the grave evils which are apt to follow when large numbers are put under the care of those who are exclusively engaged in cheap work. We would also advise all members of the profession to think carefully over the situation before accepting such seductive offers as may be made to tempt them to undersell their brethren. Fees are not everything. Position and independence counts for much in the happiness of life, and if practitioners whose work only partly lies among these clubs find their position under present circumstances intolerable, what will it be to those who will be bound to them exclusively?

We should then advise the Cork friendly societies to come to terms with their ordinary medical advisers rather than import strangers. Even doctors are wary nowadays; whatever salaries are offered, they will naturally inquire about the work, and when they find that their future masters consider twopence per week far too much for attendance on a family, most of them will, we hope, hesitate to accept such a position.

There can be little doubt that few reputable practitioners will be likely to present themselves at Cork in the teeth of the whole body of the local profession. The class of practitioner they will obtain, even if successful, will be very inferior to those they have been accustomed to, and they might with advantage consider in the doctor's behalf an aphorism they are ready enough to consider on their own—namely, a fair day's work is worth a fair day's wage.

There can be little doubt that most members of the Association, and of the profession generally, will thoroughly sympathise with the bold efforts of this Irish Branch to inaugurate an important reform, and one which vitally affects the interests of the mass of the profession.

MEDICO-PSYCHOLOGICAL ASSOCIATION OF GREAT BRITAIN AND IRELAND.—Examinations for the certificates in psychological medicine of this Association will be held in London, Edinburgh, Glasgow, Aberdeen, and Dublin on December 20th. Candidates intending to present themselves for the examination should give fourteen days' notice to the Registrar, Dr. Spence, Burntwood Asylum, Lichfield.

THE SCOTTISH COMMISSION ON INEBRIATES.—The Departmental Commission on the treatment of habitual offenders, vagrants, beggars, and inebriates, has, under the chairmanship of Sir Charles Cameron, held sittings in Glasgow and Edinburgh. Dr. Farquharson, M.P., a Commissioner, and Dr. J. F. Sutherland, the Secretary, have been also present. Among the witnesses, whose evidence has been given in private, have been Dr. Yellowlees, Professor W. T. Gairdner, and Dr. Alexander Peddie.