Fig. 3.

Fig. 5

For transfusion of blood (Fig. 7) the same apparatus is used, modified by the addition of a connecting tube with cannula attached for the giver of the blood.

After a little blood has passed from the giver to the receiver, saline fluid is infused into the giver's veins to take the place of the blood lost. Then more blood is transfused, the loss of which is again made good by the substitution of saline fluid. By this plan the giver scarcely misses the loss of some 14 or 16 ounces of blood, and does not faint during the operation.

A CASE OF BULLET WOUND OF THE PREGNANT UTERUS.

By ARTHUR W. PRICHARD, M.R.C.S.,
Senior Surgeon to the Bristol Royal Infirmary.

THE following case, in my opinion, will be of interest. It is the only case that I can find recorded of a bullet penetrating the pregnant uterus and intestines, and though it ended fatally, yet for nearly a week it gave us hopes that the patient would escape with her life. The facts are these:

On the morning of Thursday, January 23rd, the patient, Mrs. W., aged 28, who was about seven months advanced in pregnancy, awoke at 7.30, kissed her husband, and smoothed his face with her hand, but did not speak to him. He got out of bed, and immediately shot her with a revolver in the abdomen. She rushed from the room, and as she did so she heard and saw another shot, by which her husband killed himself.

She was admitted into the Bristol Royal Infirmary about 10 A.M. in a collapsed condition, and the usual restoratives were at once applied. I saw her at 1 P.M. There was then considerable reaction, and she was able to answer questions as to the occurrence. The entrance wound of the bullet was 3 inches above and 2 inches in front of the right anterior superior spine; there was little sign of hæmorrhage, and no great resonant distension of the abdomen. The fetal heart was easily heard. A probe introduced would not pass through the muscles of the parietes, and as there was no distinct evidence that the bullet had penetrated the abdominal cavity I resolved to wait a few hours before operation.

About 6 P.M. the patient's pulse was quickening, distension was increasing, and Mr. Thomas Carwardine, the house surgeon, to whom I am greatly indebted for the unremitting care which he gave to the patient throughout the case, found a bruised spot under the skin on the opposite side of the body, half-an-inch above and 5 inches behind the left anterior superior spine, and extracted the bullet under cocaine. He then summoned a consultation.

At 8.30 I made a median incision of the parietes from the umbilicus to 2 inches above the pubes, and could easily see the entrance and exit wounds on the sides of the uterus. I had to prolong the incision 2 inches higher to enable me to put my hand round the uterus, and, having put on a temporary rope *écraseur*, performed Cæsarean section. The dead fœtus and placenta, both shot through, were removed without hæmorrhage, and, seeing the danger of leaving a uterus with two holes in it in the abdomen, I substituted a wire clamp for the rope *écraseur*, and removed the uterus 1 inch

above it, fixing the pedicle outside the abdomen at the lower part of the wound by hysterectomy pins. So far the operation had not lasted long. On examining the intestines I found three coils of small intestine perforated, the wounds of entrance and exit being markedly dissimilar. I closed each of these six apertures by a "purse string" silk suture, Mr. Carwardine tucking in the protruding mucous membrane in the exit wounds as I tied the knot. An artery in the mesentery that was shot through and bleeding was ligatured, and then I emptied the peritoneal cavity of blood and clot, having carefully washed every part with boracic lotion. A big glass drainage tube with a gauze drain in it was put in Douglas's pouch, and I sutured and dressed the abdominal wound, leaving the drainage tube next to the pedicle. The operation took one hour and a-half.

Everything seemed to be going fairly well for the next three or four days. There was some vomiting, but little abdominal distension. She was fed *per rectum* entirely for two days, and a turpentine enema on the second day, repeated several times afterwards, relieved the bowels of flatus. A natural stool without any sign of blood was passed on the fourth day, and this was followed by slight diarrhoea. A brownish vaginal discharge came on. She did not complain of great pain, and when it came on $\frac{1}{4}$ gr. of morphine hypodermically gave her several hours' sleep. After the fourth day she took stimulants—brandy and champagne, thin gruel, Brand's essence of peptonised cocoa by the mouth, and until the afternoon before her death I had reason to hope for her recovery. The glass drainage tube was pumped out two or three times a day, and sanguineous fluid, which became purulent later, to the amount of 3 or 4 drachms each dressing, was drawn out. The pedicle was kept dusted with boracic powder.

A change for the worse came on suddenly on the afternoon of the 29th, and she died collapsed in the evening just twelve hours under the week after she had been shot.

At the necropsy we found that death was due to purulent peritonitis, which was not over the whole of the abdomen but chiefly in three or four foci, the worst being round some extravasated blood that lay inside the wound of exit. I very much regret that I did not at the time of operation excise and drain both entrance and exit wounds. It was satisfactory to find that all the holes in the intestines were completely sealed. The pedicle was also in a very satisfactory shrivelling condition. The entrance wound in the foetus was just to the right of the spine, about the neck of the eighth or ninth rib, and the exit wound in the costal cartilages of the same side; the shot thus passed through the pleura, diaphragm, and liver.

A CRETIN TREATED BY THYROID EXTRACT.

By WM. RUSHTON PARKER, M.A., M.D. CANTAB.,

Honorary Surgeon to the Kendal Hospital.

M. E. K. was born on March 1st, 1888, her mother—who suffers from hare-lip and cleft palate—having two months previously suffered a great shock from the rapid death of one of her boys from croup. Otherwise there is nothing noteworthy in the family history—no goitre, no consanguinity, and the four grandparents living to a good age. During infancy it was noticed that she differed from other children in being always very still and quiet, in sleeping inordinately, and in never crying. Up to 6½ years of age she remained very stunted in growth, propped up all day on a chair, being unable to sit up or stand, usually very stolid, and sleeping indefinitely, breathing very noisily, and unable to recognise people or to speak. She was always cold, liking to be near the fire. Her actions were very slow, so that she would take a full minute to raise her arm when asked to shake hands. Her bowels were extremely costive, there being usually only two or three very painful motions in a month.

On November 24th, 1894, the child was photographed, being held up for the occasion by her elder sister. At that time

Photograph taken November 24th, 1894.

her face was swollen, so that her eyes were often scarcely visible; her lips were swollen and livid; her tongue was swollen and livid, and commonly protruding; the teeth were black and stumpy; the mouth seemed always full of phlegm, and the breath smelt; the bridge of the nose was sunken and the tip flat; the neck was thick, the thyroid gland quite

imperceptible; there were soft swellings above the collar-bones and similar masses outside the nipples; the limbs were short and stumpy, the belly was swollen, and an umbilical hernia protruded to the size of a walnut. The hair of the head was not noticeably scanty or coarse.

A 5-grain thyroid tabloid was given every weekday in one dose for the succeeding six months, during which time the swelling gradually vanished from the tongue, then from the face, then from the body and limbs, so that in a few months her mother thought she felt quite a stone lighter, although she had grown some inches in height.

On May 10th, 1895, she was again photographed, looking very much like any ordinary child of two years of age; she

Photograph taken November 28th, 1895.

had then been able to sit upright for about two months, and could stand with the help of a chair; her bowels had long ago become quite regular, and the hernia long since disappeared.

In June the tabloids were increased gradually to 10 in a week—namely, one daily and an extra half on alternate days, and from October onwards she took 2 every day.

In July she could relish a meal like any other child; two lower and one upper permanent incisor had appeared. She enjoyed sitting out in the sun all through the summer, and was fond of rolling about on the hearthrug.

By October she was full of fun and activity, and quite affectionate; she could toddle about with a very little help; she had a large appetite; she was quite quick and intelligent in shaking hands, recognising people, etc.; she played and squabbled like any normal child.

On November 28th, 1895, after twelve months' treatment, she was photographed once more, having very much the appearance of any healthy child from 2 to 3 years of age, but being unable to talk—with the exception of saying a very few simple words like "dad," "mam," etc.—and having several healthy permanent incisors, most of the old black stumpy teeth having vanished.

The thyroid treatment caused no unpleasant symptoms whatever, so that it was never discontinued or the dose reduced; but the impression throughout has been that the child was taking just about as much as it could tolerate, being occasionally on the verge of diarrhoea and feverishness.

PRESENTATION.—Dr. James Bauchop has been presented by the members of the Hollinwood Ambulance Class with a silver-mounted ebony walking stick engraved with a suitable inscription.