hold to the opinion he has expressed. Fortunately science rests upon other grounds besides those upon which he has based his attack on my work.

Naturally the question that has been raised, namely, the question of the pre-existence, as such, of the carbohydrate obtained by the treatment of proteid matter with potash, had to be considered in my original inquiry upon the subject. I had been labouring under the idea that the carbohydrate obtained in the potash method of analysis came from free glycogen simply existing in contact with the proteid, and it was only in the course of work that results presented themselves which led me to question this idea, and to have ground before me suggesting that the carbohydrate found was derived from the proteid operated upon.

I learnt that the amount of carbohydrate obtainable was influenced both by the length of time during which the potash was in contact with the proteid substance and by the strength of the potash solution used. Now if the carbohydrate had existed in a free form the act of solution ought alone to have placed it in a position for the whole of it at once to be recognisable. What, it has to be said, is actually observable is that more carbohydrate is obtainable under lengthened exposure to the action of potash, and by boiling with a strong solution of potash, than when the period of contact is shorter, and when a weak solution is employed. The only conclusion legitimately to be drawn from these facts is that the carbohydrate has its origin in the breaking down of the proteid matter. Sugar, of course, does not enter into the question as a source of the carbohydrate. In the first place, the carbohydrate obtained by the agency of potash is an amylose carbohydrate; and, in the second place, sugar would be destroyed in the process employed.

As regards the carbohydrate obtained, it may be said with absolute certainty that it is not glycogen. Glycogen is precipitable by alcohol of moderate strength; 60 per cent. secures its complete precipitation. The body obtained from proteid is, on the other hand, only precipitable by spirit of much greater strength. Glycogen is readily convertible into sugar by boiling with sulphuric acid of 2 per cent. strength. Not so, however, with the cleavage carbohydrate; this requires prolonged boiling with sulphuric acid of 10 per cent. strength to obtain any marked amount of conversion.

(To be continued.)

VAGINAL HYSTERECTOMY FOR MYOMA OF THE UTERUS.¹

By ARCHIBALD DONALD, M.D., C.M.,

Obstetric Physician, Manchester Royal Infirmary; Surgeon to St. Mary's Hospital, Manchester.

VAGINAL hysterectomy for fibroid tumours has been practised extensively in France and in Germany for several years, but as yet the operation has made little advance in England. So far as I am able to ascertain, the only case recorded by an English gynaecologist is one reported by Dr. Heywood Smith in the *British Gynaecological Journal* of February, 1894.

I wish here to record four cases in which I have removed the uterus by the vagina on account of fibromyoma, and to add a few remarks on the indications, advantages and limits of the operation.

CASE I.—S. S., single, aged 37, admitted to St. Mary's Hospital, Manchester, June 8th, 1892. For two and a-half years has suffered from menorrhagia and metrorrhagia. During the last six months has had several attacks of serious flooding. There is a copious watery intermenstrual discharge. Patient is very anæmic, with hæmic murmurs and blanched face. On abdominal palpation a large smooth rounded mass was felt rising 3 inches above the pubes, very firm and hard. The cervix was small and nulliparous, but through the anterior fornix the rounded tumour was felt, and was recognised as the enlarged body of the uterus. On bimanual examination the tumour was estimated to be about the size of a four-months' pregnant uterus. Sound passed 6 inches. Owing to the weak condition of the patient I hesitated to perform abdominal section; but, knowing how well such patients bear vaginal operations, I determined to try to remove the tumour by vaginal hysterectomy.

Operation, July 8th, 1892.—Preliminary ligature of vaginal mucous membrane above line of incision. Then bladder was separated from uterus, pouch of Douglas opened, broad ligaments tied and divided alternately until uterus was quite freed from attachments. Owing to its bulk it would not pass through aperture in vaginal vault. The cervix was pulled well down with volsella forceps, the anterior uterine wall slit right up

with scissors, and the tumour cut into and diminished by knife and scissors. The uterus was then drawn through as an elongated sausage-shaped mass. The anterior and posterior peritoneal margins of the gap in the pelvic floor were then united in the middle by a fine silk suture and the raw edges of the anterior and posterior vaginal walls were treated in the same way. The gaps at each side, through which the ligatures tying the broad ligaments passed, were loosely filled with iodoform gauze.

The patient made a rapid and uninterrupted recovery. Temperature ranged from 99° to 102° during first week, but after that gradually came down to normal. Patient was able to take nourishment from the first, had no pain, and slept well. Wound was dressed daily (gauze renewed and vagina douched). Ligatures removed on the tenth day. Patient discharged August 11th. Wound in vaginal vault firmly healed.

After-History.—Patient presented herself at hospital two months after discharge. Her appearance was so much improved that it was difficult to recognise her. She is now perfectly well and strong.

CASE II.—S. A. D., aged 27, married, no children, sent by Dr. Laslett, Farnworth, to St. Mary's Hospital, Manchester, and admitted October 5th, 1892. For nine or ten years has suffered from severe dysmenorrhœa, a constant pelvic aching, bearing down, and occasionally pains shooting down right leg. For two or three days each month patient is quite incapacitated. At times she suffers from difficulty in passing water. No menorrhagia or metrorrhagia. On vaginal examination the cervix was found to be long and conical, os small, and the body of the uterus was felt (retroverted) through the posterior fornix. On its left was a small hard nodule about the size of a walnut, and on the right, bulging into the lateral fornix, was a large, irregular, very hard body, closely attached to the wall of the uterus.

Vaginal Hysterectomy performed November 9th, 1892.—The steps of the operation were similar to those described in the previous case, except that as the tumour was not so large the uterus was removed entire. There were two intramural fibroids, the larger of which was the size of an orange, and had grown between the folds of the right broad ligament. The patient made a rapid recovery; highest temperature 101.4°. Ligatures removed on November 23rd. Patient discharged December 3rd, wound being soundly healed. After-progress was quite satisfactory; there was immediate and complete relief from her symptoms. Patient is now in excellent health.

CASE III.—A. B., a widow, aged 45, admitted into St. Mary's Hospital, Manchester, May 27th, 1893, suffered for the last four years from excessive uterine hæmorrhage, which of late has been so profuse as to render her a complete invalid. In the intervals between the attacks of hæmorrhage there is continuous watery discharge. On examination the body of the uterus was found to be very hard and almost symmetrically enlarged; the upper border was felt about 3 inches above the pubes and over the right Poupert's ligament. The size of the enlarged uterus was estimated on careful bimanual examination to be rather bigger than a foetal head at term. The cervix was not expanded by the tumour—the cervical canal being narrow and long.

Operation, June 14th, 1893.—The uterus was removed *per vaginam* exactly as in Case I. Its removal was impossible until the anterior uterine wall had been incised and the greater portion of the tumour—which grew on the posterior wall—had been removed piecemeal with scissors.

The patient made an excellent recovery; the temperature only once touched 100°, and the pulse never ran above 100. She went home on July 8th, twenty-four days after operation.

After-history has been completely satisfactory; she is now in very good health and able to work in the mill.

CASE IV.—Mrs. P., married, aged 38, nulliparous, consulted me on June 28th, 1893, on account of pain in the left iliac region and bearing down. She also suffered from menorrhagia—periods lasting ten days—and dysmenorrhœa. Duration of symptoms two years. On examination under chloroform the uterus was found antverted and of normal size. Two tumours were felt *per vaginam*—one behind the uterus in the pouch of Douglas; the other, which was fluctuant and much the larger, in the left postero-lateral quarter of the pelvis.

Abdominal Section, July 15th, 1893.—On opening the abdomen the swelling on the left was found to be a dilated Fallopian tube, elongated, tortuous, much distended at the distal extremity, and containing about six ounces of pus. It was removed along with its ovary. The other tumour, lower down in the pouch of Douglas, was found to be a fibroid, the size of a walnut, growing from the posterior wall of the uterus.

Patient improved rapidly after operation—gained weight and strength; but periods still continued to be rather profuse, and dysmenorrhœa persisted. On June 13th, 1894, she was sent to me by Dr. Dochard, of Whitefield, on account of the menorrhagia, which had again become very troublesome. The tumour in the pouch of Douglas was now found to be much increased in size; it was about as large as a cricket ball, and was incarcerated in the pelvis. Patient had much pain in the back and occasionally suffered from urinary difficulties. She was very anæmic.

Vaginal Hysterectomy, June 28th, 1894.—After tying the greater portion of the broad ligaments the tumour was diminished in size with scissors and the uterus was removed.

Patient made an uninterrupted recovery, and is now in very good health.

SUMMARY OF CASES.

The indications for the operation in Cases I and III were the prolonged and excessive hæmorrhage and continual leucorrhœa. The patients had been drained of blood, and had all the symptoms of extreme anæmia. In Case II the operation was undertaken on account of pressure symptoms and dysmenorrhœa caused by an intramural fibroid, which had developed in the folds of the broad ligament. In Case IV the rapidity of the growth of the tumour—which could be very definitely determined owing to the previous abdominal section—in addition to menorrhagia and pressure symptoms gave ample reason for operating.

As regards the size of the tumour this varied in the different

¹ Read in the Section of Obstetrics and Gynaecology at the Annual Meeting of the British Medical Association, held in London, July-August, 1895.

cases from that of an orange to that of a four months' pregnant uterus. In three of the four cases the uterus had to be diminished in size before it could be removed through the aperture in the vaginal vault. In the larger tumours neither the dimensions nor the weight could be accurately ascertained after operation on account of the mutilation which was necessary. All the tumours were specimens of intramural fibroid, but in Cases I and III they had developed more towards the uterine cavity, while in Cases II and IV they bulged out the peritoneal aspect of the uterus. The cases practically divide themselves into two classes:

(1) Intramural fibroid of moderate size, with a considerable amount of its periphery in contact with the lining membrane of the uterus, causing excessive hæmorrhage.

(2) Intramural fibroid developing towards the outer surface of the uterus, and more or less incarcerated, causing distress from pressure on pelvic tissues and organs.

REMARKS.

A few years ago the only method of dealing with such growths, apart from palliative treatment, was by oöphorectomy. With the improvement in the technique of vaginal hysterectomy, this latter method now presents itself as an alternative to oöphorectomy, and it was between these two operations that I had to make my choice.

In comparing these operations, the first question we have to ask concerning each is, What is the chance of immediate death? The mortality of oöphorectomy for fibroid is variously estimated at from 2 to 10 per cent. The latest statistics I can find are contained in an article by Hermes, in the *Archiv. f. Gynäk.* vol. xlviii, part I:

Kaltenbach ...	48	cases with 2 deaths.
Wiedow ...	66	" "
Leopold ...	35	" 4 "
Hofmeier ...	40	" 2 "
Hofmeier ...	15	" 2 "
Olshausen ...	13	" 4 "
Olshausen ...	14	" 4 "
Freund ...	23	" 2 "
Fehling ...	30	" 3 "

Total ... 312 " 28 " = 9 per cent.

In estimating the mortality of vaginal hysterectomy for fibroid, we have to go to foreign statistics. The largest list of cases I have seen is given by M. Landau who publishes the following figures:²

Richelot ...	39	" 1 death.
Doyen ...	28	" 1 "
Jacobs ...	22	" 0 "
Péan ...	300	" 6 deaths.

Total ... 389 " 8 "

The percentage mortality of 389 cases therefore was a little over 2 per cent. This statement probably gives too favourable a view of the mortality. Leopold³ in 1890 published his own statistics: in 35 oöphorectomies for fibroid there were 4 deaths (about 11 per cent.); in 21 vaginal hysterectomies there were 3 deaths (14 per cent.). In considering these statistics it is to be remembered that in all operations the mortality diminishes with the experience of the operator, and that vaginal hysterectomy is comparatively a new operation. We are therefore justified in claiming for vaginal hysterectomy a mortality at least as low as that shown by oöphorectomy; indeed the balance of evidence is decidedly in favour of the former. There are one or two reasons why we might conclude *a priori* that the vaginal operation would show a comparatively small mortality, in spite of the fact that it is generally a prolonged and apparently a trying method. In the first place, there is a notable absence, or comparatively small degree, of shock immediately after vaginal operations, even in those which involve the opening of the peritoneal cavity. This will be recognised by all who have seen the quiet, and even restful, condition in which a patient is often found a few hours after vaginal hysterectomy for cancer. There is none of the restlessness and anxiety which is so common a sequence of the most favourable laparotomy. This absence of shock is probably explained by the fact that the intestines are not touched, and that they are not exposed in any but the smallest extent to the external air. For the same reason there is much less liability to abdominal sepsis

or ileus after the vaginal operation than after abdominal section. The complete drainage which is obtained in the former operation may also act as a factor in the prevention of sepsis. It is for the two reasons just stated that abdominal hysterectomy offers distinct advantages in cases in which the patient has been drained of blood by long-continued hæmorrhage. In this class of cases the simplest laparotomy becomes a critical procedure.

Apart from the advantage in immediate mortality which I believe belongs to vaginal hysterectomy as compared with oöphorectomy, there are minor advantages belonging to the former both in the nature of the convalescence from operation and in the ultimate results. The one is a radical operation; the other, if not exactly palliative, is at least only an indirect method of cure. The cure in the one case is manifest within a few weeks of the operation; in the other the patient has generally months if not years of considerable discomfort, during which the tumour is slowly diminishing; nor is permanent cure after recovery from oöphorectomy always a certainty, a fact which is admitted by its strongest partisans. Finally, the absence of an abdominal scar is a distinct advantage on the side of vaginal hysterectomy. It is not easy to estimate the frequency with which ventral hernia occurs after oöphorectomy, but there is no doubt that, even with the greatest precautions, ventral hernia does occasionally follow abdominal section, and that when it does occur it is a source of real discomfort if not of actual danger to the patient.

In spite of the advantages which vaginal hysterectomy offers in certain cases of fibroid tumours, it is not an operation which is likely to be very extensively practised in this country, for two reasons: (1) It is only applicable to a limited number of cases of fibroid tumour, in which the symptoms call urgently for operative interference. In all cases in which the fibroid projects well into the uterine cavity, the proper treatment is to dilate or incise the cervix and to enucleate the tumour. Further, only tumours of a moderate size can be removed by vaginal hysterectomy. It is true that this limit has been considerably extended by experience and by the method of removing the tumour in pieces. But nevertheless, if the tumour rises above the pubes to the extent of more than 3 or 4 inches, or is of such volume that it will not pass through the pelvic brim, the case is not a suitable one for extirpation by the vagina. (2) The technical difficulties are at times so great as to tax one's patience and perseverance very severely, and for this reason it is not likely to become popular.

ON THE TREATMENT OF PUERPERAL ECLAMPSIA OCCURRING DURING PREGNANCY, AND PRESENTING NO SIGNS OF LABOUR.¹

By JOSEPH GRIFFITHS SWAYNE, M.D.LOND.,
Consulting Obstetric Physician to the Bristol General Hospital.

In the older handbooks of midwifery, such as I used when a student, puerperal convulsions were divided into three classes—namely, the hysterical, the epileptic, and the apoplectic. It is difficult to understand why such a classification was ever made. There is only one of these three that has any claim to be considered a distinct disease *sui generis*, and that is the epileptic form, or puerperal eclampsia as it is more correctly termed. The other two are merely attacks of hysteria or apoplexy which happen to come on during pregnancy or parturition. But, in my opinion, it would be much more rational to divide cases of puerperal eclampsia into three kinds, according to the time when the attack begins, whether before, during, or after labour; for the prognosis and treatment differ widely in each of these varieties. My own statistics of puerperal eclampsia abundantly prove this. Since I have been in practice (during a period of more than fifty years) I have met with 36 cases of puerperal eclampsia. Of these, 11 came on during pregnancy (before any signs of labour were present), 20 during labour, and 5 after labour. Now in the 11 that occurred before labour, 6

² *Arch. f. Gynäk.*, vol. xlvii, part i.

³ *Arch. f. Gynäk.*, vol. xxxviii, part i.

¹ Read in the Section of Obstetrics and Gynecology at the Annual Meeting of the British Medical Association held in London, July-August, 1895.