given to the hair and scalp, and the leaving behind of all clothes which he had worn up to that time.

Coming back for a moment to the question of vaccination and revaccination, it was ascertained that Nos. 6 and 7, the first infected by No. 5, had not been vaccinated, and no vaccination marks were discoverable in the first four cases. It is interesting to note the comparatively large number of cases attacked who were over 40 years of age, some of which were somewhat severe, whilst the younger people who had been vaccinated in infancy had comparatively mild attacks. Of the 21 cases here recorded, 13 were 40 years old and upwards, 5 of them being over 60. Some of the attacks were so mild that, had we not been on the small-pox track, the rash would probably have been overlooked; this has a bearing on the probable spread of the disease from mild undetected cases.

The Vagaries of Small-pox.—It is to be noted that in the first batch of cases (1 to 4) although about 100 inmates were exposed to the risk of infection, and revaccination was carried out to a limited extent only, there was no further development of the disease. Why? It is impossible to say. Then, why should it select No. 15 to act as a new source of infection in the house, and pass over some thirty or more in the infected wing of the hospital? And why should No. 16, after being exposed to infection from January 7th, not be attacked till February 5th? Again, it is not unreasonable to suppose that of all those exposed to infection from No. 15, so many would take it, and there an end. No. Four were infected by this man, and then the last removed acts as a fresh centre of infection amongst a body of men who had been quarantined, and who had already been exposed to infection for nearly a month, and No. 21, who refused revaccination, is attacked.

Recognising the great protective influences of vaccination and revaccination, are there not some other influences at work in successive epidemics, and if so, are they meteorological, virulence of poison, or what? Just ten years ago there was an outbreak of small-pox at the workhouse. A man was admitted to the hospital on February 16th, 1886, and the next day developed the disease and was removed. The next case cropped up in the house, some distance away, on March 3rd, and on March 17th a third case occurred in a man who slept in the next bed to the first case (which he only occupied for three hours) being twenty-nine days after his removal. There were no further cases. There were no cases in the town so far as could be ascertained by inquiring of the medical officer of health. Why should we have had so few cases then when, so far as I remember, no special precautions were taken beyond removal of patients affected?

Precautions Taken.—It is a happy incident that so few inmates, out of 800, have been attacked, and that the disease was confined entirely to the male side of the house. Revaccination was carried out as far as possible, but this was not all that was done. So soon as a suspicious case occurred, after his removal the bedding, etc., was burnt, and the particular part of the building where the case had cropped up was immediately quarantined for the requisite period, and no inmates allowed to go out except such as had been revaccinated.

Besides the cases here reported there were several suspicious cases which were isolated and revaccinated. In all the revaccination was successful. One other case of small-pox was caught in the tramp ward, and at once isolated and then sent to the small-pox hospital.

NOTES ON A CASE OF UTERINE CYSTIC FIBROMA.¹

By JOHN HOSACK FRASER, M.B., M.R.C.P. EDIN.,
Bridge of Allan, N.B.

The following is a summary of the essential facts of the case:

Mrs. A., aged 67, was first seen in 1887, when on examination an abdominal tumour was found. She had always enjoyed very good health up to June, 1870, when she had a miscarriage at the third month, followed by pelvic inflammation. In June, 1880, she had another miscarriage followed by flooding. After this, as she continued to feel weak for some

months, and to increase in size, she thought she must be pregnant, and imagined she felt quickening. The swelling, however, felt hard, and not like the pregnant uterus. Menstruation at this time was perfectly regular, and continued so until the age of 60, when it ceased. As she was anxious about her condition, she consulted Dr. Thomas Keith, who informed her that she had a tumour. As it did not give her much trouble she had no further advice until 1885, when she underwent a course of treatment by ergot. This had the effect of diminishing the size of the tumour somewhat. In October, 1887, she again saw Dr. Keith, who advised treatment by electrolysis. Before this treatment was begun, however, the tumour was aspirated, with the result that 18 pints of blood-stained fluid were withdrawn. The idea of employing electrolysis was now abandoned, but no more radical operation was advised, as the patient was a bad subject for any such treatment. With the exception of some bronchial attacks, she remained fairly well until May, 1891. On the 30th of the month, owing to the further increase in size it was found necessary to tap her again. On this occasion 19 pints of fluid were removed. It had a dark red colour, and deposited a considerable amount of sediment. Under the microscope the latter showed blood corpuscles, pigment, and fatty granules. Both Drysdale's cells and columnar epithelium were absent. The fluid coagulated spontaneously on standing.

On September 3rd, 1891, tapping was again necessary. After this tapping the patient suffered a good deal from vomiting, and had pyrexia. On January 11th, 1892, when she was tapped for the fourth time, the fluid was found to contain pus, besides blood, fatty granules, and micrococci. After this she improved a good deal in health, and the temperature fell to normal. Up to May, 1895, the patient was tapped altogether on twenty-two occasions. After the tenth tapping she had some bronchitis and pyrexia. From the twelfth to the twenty-second tapping the average interval was eight weeks.

On May 25th, 1895, the patient was seized with a severe rigor, and became greatly collapsed, with a temperature of 103°. Perforation of the cyst wall was diagnosed, and the tumour again tapped, 9 pints 10 ounces of fluid being removed. On the following day dulness, tubular breathing, fine crepitations, increased vocal fremitus, and resonance were found at the bases of both lungs. The patient was very restless, suffering from cough, shortness of breath, and frequent attacks of vomiting. The pulse on May 27th was 126, respirations 40. During the next few days the temperature varied between 100° and 103°, the pulse was weak and at times intermittent, from 120—150, and the respiration rose to 56 per minute. Marked distension of the abdomen and tympanites set in on the 28th. After this the condition became rapidly worse, and she died on June 1st.

Post-mortem Examination.—The body is well nourished. There is a considerable quantity of subcutaneous fat. *Thorax.*—The left pleural cavity contains 1½ pint of blood-stained serum. The left lung is congested at its base, and the bronchi are inflamed. The right lung is adherent; some of the adhesions are old, but most of them are recent. The right pleural cavity contains half a pint of blood-stained serum. The

¹ Read in the Section of Obstetrics and Gynaecology at the Annual Meeting of the British Medical Association held in London, July-August, 1895.

middle and lower lobes of the right lung are in a state of red hepatisation. The pericardium contains 2 ounces of blood-stained serum. The heart is healthy, but loaded with fat. The valves are all competent and healthy. *Abdomen.*—The stomach is dilated, and in a state of chronic catarrh with superficial ulcers. The liver is enlarged and fatty. The spleen is normal. The kidneys are fatty. A large cyst occupies a considerable part of the abdominal cavity. It is firmly adherent anteriorly to the abdominal wall, the adhesions in some parts being cartilaginous in density. The mesentery and bowels are adherent to the cyst. The lymphatic glands, retroperitoneal and mesenteric, are enlarged and indurated. The cyst arises from the fundus of the uterus by a narrow pedicle. The solid part of it is on the right side. The cyst contains a reddish-brown fluid which measures 30 ounces. The entire tumour has been separated from the surrounding parts, and removed along with the uterus and its appendages. Fluid, identical with the contents of the cyst, was found in the peritoneal cavity. The wall of the cyst has undergone fatty degeneration, and posteriorly it is very attenuated; and at the upper and posterior part of the wall, where the thinning is most marked, there is a small perforation about the size of a sixpence. The cyst is unilocular. There is a small cyst about the size of a pigeon's egg in the left ovary.

REMARKS.—A uterine cystic fibroma is so rare that I need make no apology for bringing this case before the Section. Let me briefly refer to the history of this very remarkable and most interesting case, and, in the first place, the sexual history of the patient and her family deserves to be noted. The patient had a miscarriage at 52, and did not cease to menstruate until she was 60. Her mother had a miscarriage at the sixth month when she was 54, and the patient's sister menstruated regularly until she was 54. The patient enjoyed good health up to 1870, when she had a miscarriage, which was followed by inflammation. From that time she seems never to have regained her former health. In June, 1880, ten years later, she had another miscarriage, which was attended with "flooding," and it is to the results of this that she attributes her subsequent troubles. Some months after this miscarriage, she observed that the abdomen was increasing in size, and for a time she thought she was pregnant, but, as the swelling was hard, and did not increase at the same rate and in the same proportions as a gravid uterus, and, moreover, as menstruation was occurring regularly, she suspected that she was the subject of a growth. About that time she consulted Dr. Thomas Keith, who told her that she was suffering from a uterine fibroid tumour. There can be no doubt that the tumour was at first an ordinary fibroid, which, in the course of years, became cystic. In 1885 she was treated with ergot, and under its administration the tumour contracted considerably—a point of some diagnostic value. About this period she was examined by a specialist, who told her that she had an ovarian tumour. It was not until 1887 that the cystic nature of the growth was discovered. It was in October of that year that she was tapped for the first time, when fluid like venous blood was removed from the cyst. A microscopical examination of the fluid was made, when the absence of Drysdale's cells and columnar epithelium helped to confirm the diagnosis. Any slight doubt that may have been previously entertained as to whether the cyst was uterine or ovarian was now removed. Beyond causing her some discomfort, the tumour gave her no further trouble until the month of May, 1891, when, owing to extreme tension of the abdominal wall and interference with the respiration and the action of the heart, she had to be relieved by tapping for a second time. The instrument which was used on this, as well as on all subsequent occasions, was Mathieu's aspirator. It is a very remarkable circumstance that there should have been such a long interval as three years and seven months between the first and second tapplings. From the second tapping onwards it was found necessary to relieve her, at first at intervals of a few months, and latterly every eight weeks. It was in January, 1892, at the fourth tapping, that the fluid was for the first time observed to contain pus, which was present on all subsequent occasions. As the pus was sweet I resolved to continue the tapplings as long as it remained so, and in the event of its becoming putrid to open the cyst and drain it. From the fifteenth tapping, which was performed in April, 1894, until the twenty-first, which took place in April, 1895, the cyst underwent a steady diminution in size, and the amount of fluid became less at every tapping, falling in the course of the last twelve months from 12 pints to 8 pints. This is an important fact, and it gave me some reason to hope, more especially as the patient was gaining

flesh and strength, that I might eventually effect a cure by emptying the cyst periodically. As to the acute attack of septicæmia, which brought about a fatal issue, I feel certain that it was caused by the perforation of the cyst.

ON THE MENSURATION OF THE DEFORMITY OF HIP DISEASE.¹

By A. B. JUDSON, M.D.,

Orthopædic Surgeon to the Out-patient Department of the
New York Hospital.

THE object of the present paper is to facilitate methods of recognising, measuring, and recording the degrees of deformity of hip disease. A great advance in the study of this subject was made in 1877 by Mr. Howard Marsh. I reproduce in Figs. 1 to 5 the original figures with which he illustrated

Fig. 1.

Fig. 2.

Fig. 3.

Fig. 4.

Fig. 5.

his first lecture on Hip Disease at the Hospital for Sick Children in London. These figures, modified in various ways, have reappeared many times in the literature of this subject, and to one clinically familiar with these deformities they do not require explanation. Mr. Marsh's lecture contains a complete exposition of the mechanics of the deformity of hip disease, and it will not be easy to add anything to what he has said.² I have, however, constructed the movable models represented in Figs. 6 to 10, in order to make this difficult subject more easy of comprehension.

We appreciate at a glance the clinical significance of abduction or adduction when combined with fixation. Abduction means apparent lengthening, and adduction apparent shortening. Abduction and lengthening, adduction and shortening—flexion and lordosis—are to us mentally interchangeable terms, but it may not have occurred to us all that these factors of deformity may be easily measured and recorded, and that we may thus better appreciate the importance of affording relief and the merits of whatever method we may adopt to remove or prevent deformity. The dolls in Figs. 6 and 7 were arranged and photographed several years ago. They show graphically the degrees of motion and deformity, but they cannot, from the immobility of their vertebral joints, show the effect of the disease on the length of the limb or the curve of the spine. These effects are, however,

¹ Presented at the ninth session of the American Orthopædic Association, Chicago, 1895.

² BRITISH MEDICAL JOURNAL, July 14th, 1877, pp. 37-39.