state of calcareous degeneration in the opened up parametrium. I stuffed a large sponge into the capsule, sewed its edges together, and as on withdrawing the sponge I found that the oozing had ceased I tied the sutures and did not introduce a drainage tube. The capsule fell to the bottom of the pelvis. Convalescence was rapid. The tumour was a small cyst of the ovary bearing some secondary cysts. The left appendages were healthy. The pain was due in this case to the pressure of the tense cyst against its capsule, which is much less yielding than the layers of the mesosalpinx. Its tenseness allowed of diagnosis. In the preceding case, No. iv, the thin-walled parovarian cyst had burrowed downward, just as the ovarian cyst had in this case, No. v, but in iv there was no pain and the fornix seemed free. The cyst was rather flaccid, hence not so easy to detect by touch as in v, nor so likely to be painful. I suspect that it is easier for a parovarian cyst developed between the layers of the mesosalpinx to burrow lower in the broad ligament than for a true ovarian cyst to push itself downwards.

CASE VI presented a great peculiarity. The patient was a domestic servant, aged 25, and single. A small cyst filled the lower part of the abdomen and intestine lay in front; the fundus of the uterus could be felt in the right groin, its cavity measured $\frac{3}{4}$ inches, the cervix lying high. No part of the cyst came down into the pelvis. I operated on May 1st, 1895, assisted by Dr. Hubert Roberts. On tapping a pint and 6 ounces of ovarian fluid escaped. The surface of the tumour was white. On inspecting the tapping hole two distinct layers could be seen, as though there were a true capsule. I placed my hand down behind the supposed capsule and pushed out the entire cyst, with a normal pedicle, the mesosalpinx and the ovarian ligament running on to the "capsule." There were no adhesions, and the opposite (right) ovary was healthy. Convalescence was uninterrupted. The "capsule" was simply the ovary itself (Fig. 7) distended by the cyst within its substance. The true cyst wall was connected with the ovarian capsule by very loose connective tissue. Had the union been close and firm, that would have been the usual condition in ovarian cystoma, and no two separate layers would have been seen on inspecting the trocar wound. On the other hand, had the capsule of ovarian tissue been adherent to pelvic structures, the cyst might have been deliberately enucleated from the ovary itself; the appearance after enucleation would then have been puzzling.

Pawlik, in his instructive essay already quoted, writes of other conditions where capsulation is simulated to the confusion of the operator. He refers to cases (not very rare in my experience) where the lower part of the front of the cyst adheres firmly to the posterior layer of the broad ligament. In another instance a large pyosalpinx lay in front of an encysted peritoneal effusion or peritoneal cyst. He speaks also of a large hydrosalpinx. A cystic tube is encapsuled, as the mesosalpinx is of necessity opened up, till the tube touches the ovary. Unless the relations are much obscured by inflammation there is little difficulty in recognising a tumour of this kind. After tapping its relation to the uterus becomes manifest. The ovary is often flattened down and atrophied beneath the hydrosalpinx; hence the operator is apt to overlook it, and to take the cyst for a parovarian, or more likely a tubo-ovarian tumour, as a large hydrosalpinx is always constricted at a certain point. Fortunately this mistake is not of high gravity to the patient, who seems to suffer little if the ovary be left behind.

Such an error, however, is more serious to the operator if he fail to recognise the truth after the operation. In this communication, I trust, that I have clearly demonstrated how important it is that the operator should know something about the anatomy and pathology of capsules. All my six cases here related recovered, and out of a large number of ovariectomies where enucleation of parovarian or ovarian cysts was undertaken, only one patient succumbed. She had been long subject to bronchitis and emphysema, and was suffering at the time of the operation from a subacute attack of the pulmonary affection greatly aggravated by distension of the abdomen. As the abdominal swelling caused great dyspnoea, I operated. There was a parovarian cyst, which, as in Case iv, burrowed downwards as well as between the folds of the mesosalpinx. Enucleation was troublesome; unfortunately there proved to be an ovarian tumour on the opposite side, with twisted pedicle, hæmorrhage into its cavity, and strong adhesions to the parietal peritoneum in the left loin, and to diseased omentum. The operation was of necessity prolonged. Unfortunately, in spite of all precautions, the pulmonary trouble increased, coughing became incessant, and the patient died on the fifth day. Death had little to do with the enucleation required in this case; the removal of the opposite tumour caused much more disturbance of parts and expenditure of time.

If the surgeon knows that he has to do with a capsule, he can proceed to enucleation freely, but deliberately, as inside the capsule he cannot damage pelvic structures. There is no space for me to discuss the correct treatment of a true capsule; as a rule, but not always, it may be treated as in Case v; but fixation to the lower angle of the wound and

drainage of the cavity of the capsule is usually needed if the enucleated tumour is in a suppurating condition. If the surgeon persists in doing ovariectomy, without any pathological knowledge, he must not be surprised if he loses cases and finds after death that he has torn up the peritoneum of Douglas's pouch, lacerated a ureter or large vein in the wall of the pelvis, mistaken intestine adherent to and stretched on the cyst for the tube in a true capsule, or committed some other deplorable error.

NOTES ON THE PERMANENT SUBCUTANEOUS METHOD OF WIRING THE PATELLA FOR FRESH TRANSVERSE FRACTURE.

By ARTHUR E. BARKER, F.R.C.S.,

Professor of the Principles and Practice of Surgery and Professor of Clinical Surgery at University College, and Surgeon to University College Hospital.

IN February, 1894, I ventured to publish in the BRITISH MEDICAL JOURNAL a new method of suture for fractured patella which had given me excellent results, and seemed worthy of further trial. Experience since then has convinced me that the procedure has its place among our other methods and has much to recommend it as a substitute for some of the latter. And from what I have heard from surgical friends at home and abroad this view appears to be gaining ground.

Since my first operation by this method I have employed it in every case of fresh fracture of the patella admitted into my wards at University College Hospital. The standing order is that I shall be sent for as soon as the patient is admitted, and that in the meantime he is to be prepared for operation. This means, as a rule, that the patella is wired within the first twelve hours after fracture; one or two have been left a little longer.

The cases thus treated have turned out so well that there seems no reason to depart from this as a routine practice. It has proved itself safe, simple, and time saving, both to the patient and surgeon. And though I have myself employed in the past many appliances for the treatment of these fractures without operation, and later have operated on them with enthusiasm by the open method, I do not think I have ever seen better results than those achieved by this very simple procedure.

But to get the very best results certain measures must be adopted in the general treatment, and certain details carefully attended to in the application of the suture, which, perhaps, I did not sufficiently emphasise in my first paper. To these I should like now to be allowed to refer as an answer to many questions put to me, and to correct some slight errors into which a few have fallen, who have done me the honour to employ this method upon my recommendation.

In the first place this operation is only suitable to recent fractures. This would appear self-evident to most people, and yet I was consulted some time ago by a gentleman who had broken his patella months before in Scotland, and who had recovered with bad fibrous union, who had been told that I could fix and hold the fragments together by a subcutaneous wire stitch without opening the joint. When I assured him that in his case after so long an interval no operation short of opening the joint and freshening the surfaces of the fragments before wiring them could possibly bring about good firm union, he looked very disappointed, as he had been incorrectly informed that this could be done subcutaneously by my method, and that the wire would take the place of the bony union. This method he was perfectly willing to submit to, but he declined altogether to have the joint opened for the larger operation.

In the next place the operation should be done as soon as possible after the injury. This generally means within a few hours. Such a rule only carries out the principles applied to the treatment of clean wounds and fractures elsewhere, that is the bringing together of their fresh surfaces as soon as possible, with a view to primary union. If the operation is delayed many hours there is a probability of the formation of clots about the broken surfaces and in various parts of the joints which may give a little trouble. On the other hand if the operation is done while the blood in the

joint is still quite fluid, the latter runs out readily when the punctures are made and leaves the broken surfaces clean.

Again, if there be any periosteal fibres overhanging the fractured surfaces, they are easier to dislodge within the first few hours than later when lymph is beginning to be poured out, and to fasten them down.

The next question of principle is the immediate employment of passive, and even slight active, movement after the operation. In my earlier cases I fixed the limb for a few days on a splint. This I now think to be quite superfluous and even harmful, except in the case of violent patients—for example, those who are drunken or suffering from delirium tremens, etc. Some antiseptic pads are now simply applied over the punctures, and over these a layer of wool and a bandage, and the patient is told as soon as he is conscious that he can move his leg about in the bed as much as he likes if it is not painful. And, in addition to this, slight passive flexion is produced daily by the patient or surgeon with the hand behind the knee. With a good strong wire round the fragments this is quite safe, and as the wounds are only punctures, they are not likely to be injured in any way by the trifling movement. I hold this early movement to be of the utmost value as preventing atrophy of muscle or bone, encouraging the normal circulation, and also as preventing stiffening and adhesions.

Then another most important measure to be adopted immediately after operation is massage. This is begun the day after operation, and continued for a couple of weeks. The effect is to cause absorption of any blood left in the joint and in the tissues around, and to keep up the nutrition of all the structures involved. It has been proved over and over again that some cases of transverse fracture of the patella, where fibrous expansion around the fragments has not been much torn, recover with very useful limbs under massage alone, though of course with fibrous union, and after some loss of time. But I know of some surgeons who, having formerly been most enthusiastic for the treatment by massage alone, have lately adopted my method of suture in addition to it, as being more rational. That is to say, that to bring the fractured surfaces together at once in the fresh state, and to secure them thus by simple but effectual means, is better than to leave them apart, no matter how beneficial massage may be in removing blood and promoting nutrition.

But this question of the removal of blood is also most important. It should be effectually done by squeezing the fluid out through the punctures which are made into the joint. And if those who advocate massage alone as a treatment for these fractures claim that its chief merit lies in its rapid removal of the effused blood, it is quite clear that this can be more completely effected at once through our punctures, and in these aseptic days quite as safely.

The material used for suture has been in all my cases (with two exceptions) stout, soft, silver wire, as thick as a No. 1 silver catheter (English scale). Silk was twice employed in deference to a suggestion that wire running along the joint aspect of the patella might be more injurious than silk. But I question whether this is true; and even if it were the advantage would, I think, be counterbalanced by the greater difficulty in obtaining a close tight suture with silk than with wire. For though the first tie of the silk can be made easily and quite securely, the second tie for the knot is apt to loosen the first, and then the fragments are not held firmly together.

But, again, the way in which the wire is closed upon the fragments is of great importance. When the two limbs of the loop embracing both fragments are brought out of the lower puncture they should be crossed, so that the posterior one is carried upwards and to the left of the anterior one, which is pulled directly downwards. When now the posterior one is drawn upwards in the axis of the femur by the surgeon's left hand (Fig. 4) and the anterior downwards in the axis of the tibia, the fragments are forced together very intimately. Then the operator's left hand (above) with the posterior wire is carried across to right, and (below) right with the anterior wire to his left, so making a twist which fixes the loop round the fragments securely. This twist is then repeated three times before the ends are cut off, leaving ends half an inch long. The twist of the wires is then pressed down through the puncture, and the ends are opened out

transversely and buried in the soft parts over the ligamentum patellæ.

The wire I have always employed is of soft silver. In thickness it is just equal to a No. 1 silver English catheter. The exact thickness by wire scale I do not know, but this measurement is accurate enough. A piece of this about a foot long should be straightened, and then softened by heating in a spirit flame.

And now as to the adjustment of the fragments. Before the wire is introduced the operator takes each fragment between the finger and thumb of opposite hands. He always stands on the patient's right side. He then proceeds to rub the fractured surfaces one against the other, which in fresh fractures is easy enough. Thus rubbing them from side to side he is soon conscious that he has dislodged all blood from between them, or clot and fibrous material, and that they grate one against the other like dry lumps of sugar. When he gets this sensation he is ready to introduce the wire. Then while he is preparing to close the wire on the fragments as above his assistant repeats the rubbing process, and finally brings the fractured surfaces accurately together, with special care that the external and internal edges correspond, and that there is no rotation of either on its vertical axis. The closing of the wires keeps them in the same plane, one passing over the front and one over the back of the fragments. Then the wires are twisted as above. Care should be taken that the first twist should not be too sharp lest the wire be snapped. This occurred in one case with a thinner wire, and I had to introduce a second wire. The second and third twists will tighten the first.

And now let me once more briefly describe the steps of the operation, bearing in mind the emphasis to be laid upon the matters of principle and detail mentioned above.

First, everything about the field of operation and the instruments, etc., is made aseptic. Then the lower fragment is steadied between the operator's left finger and thumb, while a narrow-bladed knife is thrust exactly through the middle of the upper attachment of the patellar ligament, with its edge upwards cutting on the bone. When the blade has entered the joint it is withdrawn, still cutting on the lower edge of the lower fragment, and enlarging the skin wound upwards to the extent of two-thirds of an inch. Through this opening a special long stout-handled needle, without any wire (Fig. 1), is now thrust into the joint behind both frag-

Fig. 1.—Needle specially designed to carry a thick wire. The eye is drilled obliquely, and should receive only a little loop on the end of the wire; this little loop should be made previously. *Vide* Figs. 2 and 3.

ments and is made to pierce the tendon of the quadriceps close to the upper edge of the upper fragment, exactly in the middle line of the fragment. When its point appears under the skin it is cut down upon with the knife, whose edge should now be turned downwards, and enter the joint beside the needle, cutting upon the upper edge of the upper fragment, so as to leave nothing between the needle and the bone. A stout silver wire (b) is now threaded through the eye of the needle (Fig. 2), and the latter is withdrawn and then unthreaded. The empty needle is then again thrust through the same lower opening, but this time in front of and close to both fragments until it emerges through the upper puncture. Here it is threaded with the other end of the wire (c), and is withdrawn (Fig. 3). Both ends of the wire now emerge by the lower opening. I then thread each through holes in the centre of short metal bars (Fig. 4) for the convenience of pulling, and fix their respective ends to the bars by a second transverse threading. Then the posterior wire is drawn upwards and the anterior downwards as described above (Fig. 4). When the fragments are adjusted the bars enable the surgeon to make a powerful even strain on the wires, and twist them as above accurately. While the wires are *in situ*, but before they are twisted, all

Fig. 2.—Needle (a) introduced behind the fragments, and receiving one end (b) of the silver wire (b, c).

the blood in the joint should be squeezed out by pressure with the hands and kneading. The whole operation should not take more than about five minutes from beginning to end.

Now I do not wish to say anything against the open method, which I have practised freely myself, and with suc-

Fig. 3.—Needle (a) passed in front of the fragments, and receiving the other end (c) of the silver wire (b, c).

cess so far, and which I regard as almost devoid of risk in good hands. But the saving of time by the method now described is most important, if nothing else is. To the surgeon it means much; to the patient the deliverance from prolonged

Fig. 4.—Wire in position round fragments and threaded through metal bars. The lower and posterior wire runs upwards to the left of the upper ready for twisting.

anæsthesia means much more. Two of my patients were over seventy, and to their lungs and general condition the difference between five minutes and fifty or more was probably immense.

Again, the absence of a more or less fixed scar on the front of the patella, and the short confinement to bed, were of great importance for obvious reasons.

That the wires so applied brings the fragments easily into good position I have proved in three cases where it was necessary to open the joint widely for old fractures, etc. The fragments came accurately into the same plane and close together when the wires were twisted as above. Moreover, the fragments were not weakened by being bored through. And that the presence of the wire on the femoral aspect of the patella was productive of no inconvenience these cases also proved. A permanent buried wire is a very different thing to a wire passed temporarily round the fragments and tied over a pad as in Kocher's operation. Here there are two punctures through which the wires enter the joint, and these are left open for a fortnight or three weeks until the wire is removed, the wire being tightened from time to time. All this means risk, and the wire is removed at a time when we know that the fragments are still capable of separating from one another. I have operated by the method now described thirteen times. All these patients have recovered without a single hitch, and have had excellent results.

On the whole I prefer this method to any other, as the routine treatment of recent transverse fractures of the patella, especially since massage has become part of the regular after-treatment, together with immediate passive and light active movement of the joint.

Other surgeons operating by this method have had just as good results, some of which I have seen and many of which I have heard of.

Patients treated by this method who at the end of six or eight weeks are walking about with a firm limb and good flexion and extension, and with apparently bony union, offer a very marked contrast to those treated by splints who at the end of two or three months have much stiffness in the joints and have to be most careful about flexion for many months more lest the bond of union be stretched and great weakness result. Many of these patients treated by splints we know are obliged to wear a support for the rest of their lives, and some recover with permanently stiff joints, as was the case

with a man of whom I have heard to-day, and of others of whom I have read or heard.

And if this procedure is contrasted with the open method, even if we were to admit that there was an equal absence of risk in both the saving of time and stress of a long operation is great and well worth considering. As to the adjustment of the fragments, there is no more difficulty in one than the other, and as to passive movement and massage, they can be begun at once after the subcutaneous method, while with the open method the healing of the wound must be first allowed to take place.

I have lately made a needle specially for this operation, which is a great improvement on the old ovarian pedicle needle formerly used. It is safer not to break, and its eye is so drilled as to carry the loop of stout wire much more easily than the older needle. It is represented in Fig. 1 as copied from my own manufacture by Krohne and Sesemann, Down Brothers, and Meyer and Meltzer. In using it I find it convenient to make a little short loop on each end of the silver wire before the operation is begun. If this is done the end can be hooked in the needle at once, and the loop is buried in the needle, and offers no resistance to the withdrawal of the latter, which a large loop of thick wire would do.

It should be remembered that patients who have once broken their patella are liable, whether operated on or not, to attacks of synovitis on over-exertion of the joint. If such an attack takes place some months after the bone has been wired it might suggest to the inexperienced that the wire was giving trouble. But this is not so, as I have seen in two or three cases. With a little rest the condition is quite relieved. One can hardly expect after so severe an injury as that which fractures the patella, that there should be in all cases a complete re-establishment of the vasomotor balance in the circulation of the synovial membrane.

A SUCCESSFUL CASE OF PYLORECTOMY.

By JAS. A. ADAMS, M.D., F.F.P.S.G.,
Surgeon to the Royal Infirmary, Glasgow.

Excision of the pylorus has not been very successful in Scotland, if one may judge from the published records. Indeed, I have not been able to find any account of a patient who has survived the operation many days. In publishing this case I hope to bring before surgeons the desirability of performing the radical operation, in preference to the palliative and at best temporary procedure, in which the growth remains untouched. Such operations give at most a few months', often less, relief to the patient, and by improving his condition hasten the growth of the tumour, which ultimately kills him.

It may be urged that this short time is worth more to the patient than his chance of recovery from the major operation; and that is true as the statistics stand at present; but as our results are much worse than those of some Continental operators, the presumption is that our methods are at fault. It therefore behoves us to improve these methods so as to obtain success in what is, after all, the more scientific operation, and if the history of this case stimulates others to its performance, its publication will have been warranted.

On July 1st, 1895, I was consulted by Mrs. C. regarding an abdominal growth she had noticed about six months previously. She stated that her age was 56 years, although her emaciated appearance made her look at least fifteen years older. She did not complain of vomiting, but stated she felt constant nausea, with disinclination for food, and had lost flesh very rapidly. She was exceedingly feeble.

The growth, which was the size of a small orange, could be moved freely, and the stomach was much dilated. On July 19th, assisted by Drs. McGregor, McPherson, and Findlay, I made a straight central incision, 3 inches long, above the umbilicus, and drew out the growth, which proved to be a carcinomatous tumour of the pylorus. The swelling was not adherent to the other viscera, and I was unable to detect any enlarged glands under the liver, or in the vicinity of the lesser omentum, or pancreas.

The tumour was dragged well out of the wound, and the abdominal cavity protected with natural sponges, wrung out

of warm boracic solution. An assistant grasped the stomach and duodenal sides of the growth, between his forefinger and thumb, compressing the viscera to prevent the escape of any contents. The knife was then passed lightly round the pylorus, at a distance of 1 inch from the edge of the growth, dividing only peritoneal and muscular coats. These were reflected, about an eighth of an inch, in the direction away from the growth, and the mucous membrane divided with scissors, as far from the tumour as possible. Immediately prior to division of the mucous membrane, the portion to be excised was clamped. Several vessels in the mesentery were ligatured with catgut, and the wound well swabbed with warm boracic solution. The stomach cavity was flushed out with the same solution, and a few fragments of food removed.

The upper portion of the stomach wound was stitched with a continuous catgut Lembert suture, until the lumen corresponded to that of the duodenal opening. The assistant, grasping the two wounds with forefinger and thumb, brought them into apposition, and they were stitched up with a similar suture, which only included the serous and muscular coats. This part of the operation was greatly facilitated by the previous free removal of the mucous membrane from the edge of the wound. The peritoneum was carefully stitched with catgut, in the lesser curvature of the stomach, and a flap of omentum lightly stitched over the gastro-duodenal wound. The abdominal cavity was not flushed. The parietal wound was closed with buried and superficial sutures. The operation took exactly sixty minutes, and chloroform was used.

For thirty-six hours nothing was administered by the stomach, but the patient was permitted to wash her mouth with warm water containing a few drops of brandy. Rectal feeding was commenced at once, two ounces of milk and half an ounce of brandy being given every four hours. From the 19th to the 21st the patient vomited twice, but this ceased after administration of 10 grains of bismuth powder.

On the 22nd, during a period of twelve hours, she had 1 ounce of Valentine's meat juice; on the 23rd, 4½ ounces of Valentine's juice; on the 24th, 7 ounces of raw beef juice, with Valentine's juice; on the 25th, 33 ounces of milk, barley water, raw beef juice, and Coleman's wine. From July 25th until July 30th she had daily an average of 50 ounces of fluid, after which date she was put on ordinary diet. The bowels moved twice on July 25th, once on the 26th, and twice daily until August 13th, when the record was discontinued.

On July 30th the patient became slightly delirious, but this condition immediately ceased on reducing the daily allowance of alcohol. The parietal wound was dressed on the fourteenth day, when the temperature rose for the first and only time from normal to 101.2° F. No cause for the rise was detected, and the temperature fell again to normal in four hours' time. The patient was allowed up on August 21st, and went to the country on September 11th.

Unfortunately, prior to the operation, no record was made of her weight, but she has markedly increased in weight and gained strength until the present date. From the date of operation she has not had a vestige of nausea, flatulence, or eructation of any kind.

Tumour.—The growth, which involved the entire pylorus, was especially prominent on the posterior wall. It was epitheliomatous in character, and occluded the pyloric opening, reducing it to the size of a No. 12 catheter.

REMARKS.—While gastro-enterostomy is an easier operation and less fatal than pylorotomy, and while the immediate effect upon the patient is very striking, it is only palliative, and the longest period that life can be prolonged is about a year. After complete excision of the pylorus the results are very different. Thus Wölfler had a case which lived in comfort for five years and a-quarter, and although the glands in the portal fissure were found carcinomatous, the stomach remained free from disease. Rydygier has also reported a case which lived two years and a-half, and was entirely free from complaint for two years. These cases, however, are excelled by Kocher, who operated five years and a-half ago on a lady who is at the present moment in the enjoyment of perfect health. I am indebted to this latter surgeon for an interesting summary of the cases recorded on the Continent, and for many valuable hints in his recent work on *Operative Surgery*. It will be noted that I did not use clamps unless on