CLINICAL LECTURE

ON A

CASE OF CHYLURIA.

By BYROM BRAMWELL, M.D., F.R.C.P. EDIN., F.R.S. EDIN.,

Physician to the Royal Infirmary, Edinburgh; Lecturer on the Principles and Practice of Medicine in the School of Medicine, Edinburgh; etc.

THIS, gentlemen, is a remarkable specimen of urine. It looks exactly like milk. The condition—chyluria—is rare in this country. I myself have only seen one other instance in which the milky appearance was so marked, though I have met with one or two other slight cases. I therefore take the opportunity of showing you the urine. I have two specimens of the urine. Let us examine their characters more carefully.

The first specimen, which was passed before breakfast, is, you will see, only slightly milky; it has a reddish tint. The second specimen, which was passed after breakfast, is perfectly white; it looks like pure milk. After standing, a thick, cream-like layer collects on the top of the urine. The reaction of the urine is strongly acid; the specific gravity 1024; the urine contains a considerable quantity of albumen and blood, but no sugar. After shaking up a small quantity of the urine with a large quantity of sulphuric ether, the white milky appearance entirely disappears, and a yellowish-coloured fluid, which is turbid, remains below the ether.

On microscopic examination the urine is found to contain a considerable quantity of granular *débris*, a considerable number of corpuscles which look like lymphoid cells, and some large round or oval crystalline-looking bodies which on careful examination appear to be globules of oil to the outer surface of which the granular *débris* adheres, and gives the resemblance of a crystalline deposit.

Dr. Noël Paton kindly examined a specimen of urine, and reported as follow:

Buff yellow, milky; reaction acid; specific gravity (pycnometer) 1019.5. Urea=1.90 per cent.; fats=1.70 per cent.; native proteids=0.97 per cent.; ash=1.03 per cent. Proteids not coagulated on boiling; albumoses or nucleo-proteids are also present. Sugar absent. Another specimen of urine received on October 26th looked exactly like milk; it was alkaline in reaction, the specific gravity 1020; it gave no precipitate with heat, but a dense coagulum with nitric acid; it did not reduce Fehling's solution. On microscopic examination it was found to contain numerous vibrios and a good deal of granular *débris*.

Now these characters show, as the naked eye appearance of the urine suggests, that the case is one of chyluria.

ETIOLOGY.

Before considering the special features of this case in more detail, let me briefly refer to the more important points connected with the etiology of the condition. Cases of chyluria are, as I have already stated, extremely rare in this country. In the great majority of typical cases of chyluria the condition is due to the obstruction of the lymphatic channels by the *filaria sanguinis hominis*. Hence it is that chyluria is rarely met with in this country, and very rarely indeed even then except in persons who have lived in tropical and subtropical countries in which the *filaria sanguinis hominis* abounds. You remember that the disease elephantiasis Arabum has the same distribution and is due to the same cause. Recently I showed you a case of solid oedema which seemed to be due to obstruction of the lymphatics, a condition which, from a clinical point of view, exactly resembles elephantiasis Arabum, and we then discussed the subject of elephantiasis. We saw that elephantiasis Arabum is usually due to the presence of the *filaria* in the blood and tissues.

The *filaria sanguinis hominis*, which is quite microscopical in size, is the embryo form of a long thin worm, which measures about three inches in length, and one-ninetieth of an inch in breadth, and which has been termed the *filaria Bancrofti*, after Dr. Bancroft, of Australia, who first described it. The worm resembles a piece of horsehair, and has been found in the dilated lymphatics. The female produces large quantities of young, which are discharged into the circulation in the form of the *filaria sanguinis hominis*, or rather in the form of the *filaria sanguinis hominis nocturna*; for three varieties of *filaria* have been discovered—one, which only

appears in the blood at night, another only seen in the blood during the day, and another which is seen in the blood persistently both during the day and the night. The *filaria nocturna* is about the diameter of a red blood corpuscle, and about one seventy-fifth of an inch in length. It seems to live in the lymphatic channels, and to migrate into the blood between the hours of 9 P.M. and 6 A.M. The number of *filaria* which the blood may contain is enormous; every single drop of blood may contain several specimens. It has been computed that, in some cases, at least forty millions are present in the blood during the night time, whereas none of this particular variety seem to be present during the day. It is found, however, that if the habits of the patient are reversed as regards day and night the *filaria* also reverses its habit; if the patient is made to sleep during the day, and keep awake and go about through the night, the *filaria* migrates into the blood during the daytime, and disappears at night. The embryo of the *filaria*, or masses of these little organisms, obstruct the lymphatic channels and produce elephantiasis Arabum, lymph scrotum, and chyluria.

In cases of chyluria the dilated lymphatics and lacteals must obviously communicate with some part of the genito-urinary tract.

I further told you that mosquitos seem to be the active agents by which the disease is propagated. The mosquito bites a man or an animal affected with the filarial disease. The *filaria* curls itself round the proboscis of the mosquito, is sucked into the stomach of the mosquito, passes into the tissues of the mosquito, grows and develops there. When the mosquito dies the worm is set free, and, getting into drinking water, is again introduced into the human subject through the stomach and alimentary canal.

It must, however, be remembered that all cases of chyluria do not appear to be due to the *filaria sanguinis hominis*. Cases are certainly met with occasionally in this country in which the patients have never lived in districts where the *filaria sanguinis hominis* abounds, and in which microscopical examination of the blood and of the urine fails to show the presence of these organisms. It seems certain that if the abdominal lymphatics are obstructed in any way, and if a communication is formed between the dilated and varicose lacteal or lymphatic channels and the urinary tract, the urine assumes the characters of chyluria. But, although this is possible, it must be admitted that the vast majority of cases of typical chyluria are due to the presence in the body of the *filaria sanguinis hominis*. Sir William Roberts, in his *Diseases of the Urinary Organs*, states that, of 30 cases which he has collected and analysed, 24 occurred in persons who had lived in hot climates and in districts where the *filaria* was known to occur—in the Mauritius, the Isle of Bourbon, Brazil, the West Indies, or India. Chyluria is common also in Barbadoes, Australia, and China.

It seems probable that in many cases of chyluria the lymphatics in the wall of the bladder, ureter, or pelvis of the kidney are dilated and in a varicose condition. Sir William Roberts has reported a case which supports this theory.¹

In this country the majority of cases occur in males, but this is owing to the fact that men travel much more than women, and that they contract the disease in the countries where the *filaria* is prevalent, and bring it home to this country. In filarious districts women are said to suffer in a somewhat larger proportion than men. Such are the general facts connected with the etiology of the condition.

Let me now briefly detail the clinical features of this particular case. They are as follows:

The patient, a single lady aged 66, was seen with Dr. A. on November 2nd. She was born in the Mauritius, and lived there till she was 30 years of age. She then came to this country, and has resided here ever since. She states most emphatically that she has never had anything like chyluria before. For some years she has suffered from bronchitis, bronchitic asthma, and has not been very strong. For some years past the tongue has been large and sore, but she has never suffered from dyspepsia. Three months ago the feeling of weakness increased, and she began to lose flesh. Her friends noticed that she looked ill. It was about this time (three months ago) that the chyluria commenced; it has continued ever since. The morning urine has frequently been quite clear, but the urine passed in the afternoon and evening has always been milky. The heart was weak, but all the organs appeared to be normal; in particular, there is no evidence of disease in the abdominal cavity.

The urine passed on November 2nd was quite clear and pale, specific gravity 1020, neutral in reaction, and free from albumen, blood, and fat. The blood examined at 11 A.M., and also at 11 P.M., was quite normal in

appearance. There were no filarial embryos. The patient was advised to continue the malto-peptin, which she had been taking for the past three days, and to which Dr. A. attributes the improved condition of the urine; to carefully regulate the diet and bowels; and a tonic containing dilute nitric acid, strychnine, and infusion of gentian was prescribed.

I have had no opportunity of seeing the patient again professionally. She is now quite well. Under the treatment which was adopted the chyluria rapidly disappeared, and has never returned.

THE CHARACTERS OF CHYLOUS URINE.

In some cases of chyluria every specimen of urine is milky; in others the chylous appearance is only seen at intervals or at some particular time of day. In many cases the urine which is first passed in the morning is clear, and it only becomes milky after eating and after exercise; and this seems to be so in the case which I have brought before you. In many cases of chyluria the morning urine is either quite normal or it contains some blood and gelatinous mucus. After eating it assumes the chylous appearance, particularly in the afternoon after dinner. The fact that the milkiness only appears after eating is of course highly suggestive of chyluria. It shows that the fatty material only gets into the urine when the lacteal and lymphatic vessels are distended during the process of digestion.

In our case the morning urine gives, as you have observed, a very definite reaction with guaiacum and ozonic ether; it evidently contains a considerable quantity of blood.

And here let me say that every student nowadays seems to know that guaiacum and ozonic ether are the reagents which we employ at the bedside for detecting the presence of blood in urine—one of the stock questions asked at a clinical examination. But my experience tells me that many students who know that guaiacum and ozonic ether are used in testing for blood seem to have had very little if any practical experience in the use of the test. I base this opinion upon the fact that they do not know the naked-eye appearance of tincture of guaiacum; they have to pick out the tincture of guaiacum by reading the label upon the bottle. Now, a student who has been in the habit of using tincture of guaiacum at the bedside knows it at once by its appearance and colour. When asked to test a case for blood, he at once picks out the tincture of guaiacum, even if the bottle is unlabelled; and I sometimes take care, with the object of seeing if the students who come before me are practically acquainted with urine testing, to turn the bottles containing such reagents as tincture of guaiacum and Fehling's solution, which can be at once picked out by their appearance, so that the labels cannot be seen. I at once see, by this little device, whether the candidate has been accustomed to work with the reagents in the wards. If he has to turn round the bottles to see their labels, I at once know that he has little or no practical experience with the test. The mere fact that a man who does not know the appearance of tincture of guaiacum or of Fehling's solution is able to tell you that these are the tests for blood and sugar, should not, I think, count for much. Candidates often cram up information of this kind, or are coached up in it, just before the examination. They may be able to remember the test for a brief time, and perhaps to use it in some sort of fashion, but often in a way which shows the examiner that they have had little or no experience with it. This is only one illustration of the sort of discrimination which examiners ought, I think, to use in judging of the clinical knowledge of candidates. A correct answer does not necessarily imply that the student knows, understands, and intelligently comprehends the question which has been put to him, and which he has answered correctly. What the examiner wants to do is to try and find out whether the candidate has a real knowledge of the subject, or whether he has merely a superficial (cram) knowledge. By observing the way in which the candidates examine patients, test urine, etc., one can in most instances very soon make up one's mind whether they have a practical knowledge of clinical medicine or not.

As I have already stated, in many cases of chyluria blood and gelatinous mucus are passed, particularly, so far as my limited experience enables me to judge, with the specimens of urine which are not distinctly milky. Now this must have some significance. The presence of blood possibly suggests an eroded condition of the bladder mucous membrane or of some other part of the genito-urinary tract. In many cases, after standing for a time, the chylous urine sets into a gelatinous

mass like blancmange, and the jelly-like mass which is formed sometimes has a distinct pink colour. In this particular specimen no such setting has taken place. The milky appearance of the urine is due to the presence of fat. In many cases the fat is in the form of an emulsion, the fat particles being so small that they cannot be recognised, even by high powers of the microscope. In other cases definite fat globules can be seen under the microscope. That seems to be the case in this particular specimen.

In cases of chyluria the kidney is rarely diseased, the urine very rarely contains tube casts. This is an important point, for it tends, I think, to show that the theory which some authorities have advocated, namely, that in cases of chyluria the fat is separated from the blood by the kidney, is erroneous. If the kidney were separating fat from the blood for long periods of time—and in some cases the chylous condition of the urine persists for years—it is probable that in the course of time the kidney substance would become diseased, and if this were so the evidence of such disease would be present in a considerable proportion of cases of chyluria.

CLINICAL SYMPTOMS.

The clinical symptoms which cases of chyluria present are very various. In some cases there are no symptoms at all. Cases have been known in which the patients have passed chylous urine for years, apparently without any disturbance of the general health or any local symptoms. Sir William Roberts mentions one case in which the patient began to pass chylous urine at the age of 25, and went on passing it until she died, at the age of 73.

In most cases the patients complain of debility; in many cases the patients become emaciated, in some cases markedly so. In some cases they are anæmic. Diarrhoea is by no means uncommon. Occasionally some of the other pathological alterations which result from the presence of the filaria sanguinis hominis in the body (such as elephantiasis of the legs or scrotum) are also present.

TREATMENT.

In the tropical (filarial) common form of the disease the object of treatment is to destroy the filaria, and to remove the lymphatic obstruction. Iodide of potassium, tincture of perchloride of iron, gallic acid, and a decoction of mangrove bark are said to be the most useful remedies. My friend Professor Lawrie of Hyderabad claims to have cured several cases by the internal administration of thymol, 1 grain three or four times a day. Benzoic acid has also been recommended. Boracic acid should, I think, be tried and possibly also corrosive sublimate.

In the non-filarial cases, the treatment must be guided by the supposed cause in each individual case; but it is rarely possible to diagnose the exact cause of the lymphatic obstruction. And in the rare cases in which the cause can be definitely determined, the nature of the lesion (malignant disease, etc.), usually precludes successful operative interference.

These, then, are some of the chief facts relating to this peculiar and interesting condition. As I have already said, it is of great rarity, and hence I have taken the opportunity of showing you the specimens, and of detailing the more important clinical characteristics of the condition.

REFERENCE.

¹ *Urinary and Renal Diseases*, p. 377.

ORAL INSTRUCTION OF THE DEAF AND DUMB.—The annual meeting of the Association for the Oral Instruction of the Deaf and Dumb was held recently at 11, Fitzroy Square, W., when Miss Amy Paget distributed the prizes to the scholars. Mr. L. Van Oven presided, and a short but interesting demonstration of the system of tuition was given. Mr. W. Van Praagh, the Director, read the report, which showed that there were now 23 girls and 22 boys in attendance at the school, and that the training college had been attended by 9 students, 7 of whom had obtained certificates. The Committee regretted that they did not receive sufficient applications from students to be trained, owing to which they were unable to supply all the demands made upon them for teachers.