artificially tuberculous, and here the results were so encouraging that Dr. de Backer determined to try the cure in the human subject.

In order to ascertain the effect of Backérine upon a healthy man, M. Bruhat offered himself for inoculation, and was injected in the flank by Dr. de Backer. At the end of five hours there was marked coldness or rigor, followed by abundant sweating. M. Bruhat went through all the stages of a mild intoxication, owing to the presence of alcohol and carbonic acid in the blood. At the end of three days all reaction had quite ceased, and no harm resulted.

After this, phthisical patients were treated by inoculation, with the result that in early stages the disease has been nearly always arrested. In phthisis of the second stage, with signs of softening, the cure or permanent improvement has been nearly 75 per cent.; in the later stages, the result much depends upon the amount of lung tissue remaining, and the vitality of the patient.

The question naturally arises: How do these ferments act? Are they specific against tuberculosis? The reply to this is negative. These cells come to the support and aid of the leucocytes (the natural phagocytes) and fight side by side with them in destroying the invading bacteria, and to quote M. de Backer's own words, "substitute a healthy, natural fermentation for one that is pathological."

It is not sufficient that the bacillus be only destroyed, for its dead body and its toxins have the power of dissolving the tissues and poisoning the system. The great point is to eliminate the microbe from the system before the bacillary intoxication becomes too much for it. The same principle is seen at work in the industry of wine making. It may happen that many thousand gallons of wine wort have turned sour, owing to the presence of bacilli (acetic, butyric, or lactic) yet all that is necessary to effect a remedy is to introduce a certain quantity of pure ferments into the wort: acetic fermentation is arrested, and the wort brought back to its normal fermentation. In this manner we may conceive the blood to be the wort, "the human must" as M. de Backer calls it, which is disturbed in its functions by Koch's bacillus just as the wine wort is disturbed by the bacillus aceti, and can be restored to its normal condition by a certain quantity of pure ferments.

Against the pathological fermentation in the body by Koch's bacillus no means of arrest has been known until now, but arrest has been effected again and again by means of these pure ferments furnished with proper nutriment.

The great condition of success is that they be absolutely pure, cultivated step by step from single cell, from culture to culture, shielded with all care from possible contact with any other organism, even that of a variety of its own species, carefully oxygenated before its final fermentation, as it will have to live an anaerobic existence in the body. Only young and vigorous cells are selected for the work, and these are furnished with their food supply and stored in specially prepared tubes.

Even if Metchnikoff's doctrine of immunity by phagocytosis be not entirely correct there can be no doubt that it plays a very considerable part, the other part being probably played by the nuclein of the cells. Here again we get an additional reinforcement from the yeast cell whose nuclein yields the same results to analysis as the nucleins of the animal cells.

In regard to cancer, that other terrible scourge of humanity, for which no cure is known, the treatment by pure ferments has given some astonishing results. To quote again from de Backer, "Out of a total of sixty tumours, we have seen eighteen cures lasting over some years. Of these four had been operated on, with rapid recurrence. Twelve cases of very rapid growth also had been very markedly retarded."

It may be asked, In what way do we explain and account for this curative action of pure ferments, there being as yet no evidence to prove that cancer is due to the action of any bacillus? The answer to this may be found in looking at the reasons that led Dr. de Backer to apply this method to such a disease. He found that the guinea-pigs used in a long course of experiments, and which lived together in full liberty during the time that injections of Backérine were being administered to them, by way of testing its innocuity, got fat,

but were generally sterile. In cases where impregnation occurred the foetus did not develop. In some animals that were killed, blighted embryos were found. He then conceived the idea that the sterility was due to "the lessened glycogenous material in the uterus," (a possible cause of sterility in the human being hitherto unsuspected).

It was known that cancer contains a very large proportion of glycogen in its chemical composition, and that this disease occurs in localities in which glycogen is normally present in large quantities (breast, liver, uterus, etc.). Hence the deduction was easy, that if ferments could materially lessen the natural glycogen of the body, they would probably have the power of checking the pathological increase of the same substance, and that the cancer cells which revel and luxuriate in its presence would starve, and the whole tumour shrink. The researches of M. Brault have recently proved that the rapidity of growth in a tumour is in direct proportion to the amount of glycogen contained in it; probably from the fact that the cells are enclosed in a medium too rich in glycogen, and so proliferate luxuriantly without order, symmetry, or selection, and this constitutes a neoplasm.

The whole account of the method, and its application to diseases due to the action of bacteria, may be found in a work by Dr. de Backer in collaboration with M. Bruhat and Dr. Charlier entitled, *Les Ferments Thérapeutiques*, published by the Société des Editions Scientifiques, Paris. A further account of cancer, with cases, is given in a pamphlet by Dr. de Backer, *De la Cancérose et de son Traitement par les Ferments purs*, Imprimerie Daix Frères, Clermont (Oise).

WANDERING OEDEMA.

By W. C. BROWN, M.D.,

Penang.

THE case described by Mr. Masterman, of Damascus, in the BRITISH MEDICAL JOURNAL of April 3rd, is an interesting and typical instance of an affection which, though by no means rare, in the tropics at all events, has attracted comparatively little attention. It cannot, however, be called "a new disease," for several previous observers appear to have noticed this malady, to have assigned to it different positions in nosology, and to have described it under different names. They have generally taken it to be a variety of urticaria; but, in the absence of more evidence as to its causation and pathology, the term "wandering oedema"—descriptive of the principal and characteristic symptom—is a good one, and none of its previous titles appear to fit it so accurately. Owing to the peculiarity of the oedema, the uncertainty as to its progress, and the anxiety as to its suddenly attacking dangerous and vital organs, the malady is an important one, and merits more attention than it has received.

Four cases have come under my own observation, and in all of them the symptoms were practically the same. The onset of wandering oedema is characterised by pain in one or both feet, which is generally referred to the plantar fascia, and the ligamentous structures about the metacarpophalangeal joints. The pain is described as being of a heavy, dull, burning character, and at first simulates very closely the onset of an attack of gout. After an hour or two, however, the pain subsides, and it is noticed that the foot is swelling round the parts where it has been most severe. Relief is, in fact, afforded by effusion. The oedema is sometimes excessive, but is always, at first, local. Later it shows a tendency to spread up the limb, and, coincidentally with its change of position, the lower swelling subsides. By the time the knee is reached the foot is free from oedema. The skin of the parts that have been affected retains a peculiar boggy feeling, with numbness and loss of sensation for some days afterwards. About twelve hours after the feet have been attacked the knuckles and wrists become painful, and the same train of symptoms occurs in the upper limbs. The oedema is, however, generally less marked, and does not appear to reach the elbow so frequently as it extends to the knee in the lower limb. As this, in its turn, subsides, pains are felt in different parts of the body, which are, in the same way, followed by localised oedemas. The scalp, the cheeks, the lips, the skin, and subcutaneous tissue of the abdominal

wall, pendulous parts of the genitalia, seem to be invariably affected. The abdominal viscera and intestinal canal also are prone to attack. Internal pain is followed by indigestion, sometimes by vomiting, and diarrhoea is frequent. This latter, indeed, seems generally to act as a curative agency, and to occur at a period of crisis. In one case that came under my notice there was deep perineal effusion into the perirectal tissue, which was followed by sharp dysentery. The hard and soft palates, the fauces, and pharynx are usually affected, but though the oedema comes so dangerously near the glottis, I have not known an instance of the larynx being attacked. Rapin, in describing what was doubtless a case of this malady as a localised cutaneous disease,¹ pointed out that there was no itchiness. This absence of itchiness in wandering oedema is peculiar, but appears to be invariable; it differentiates the malady widely from urticarial affections, and may be taken as a characteristic symptom.

The duration of the illness is generally about three days, but it is often protracted to a week or even longer. Diarrhoea and copious perspiration are frequently associated with its subsidence, and have doubtless a curative effect, but these symptoms are by no means constant concomitants of the malady.

During the attack the condition of the patient is not characteristic of severe illness. The temperature is but slightly increased, the pulse and heart maintain their normal condition, and there is no albumen in the urine. There is generally gastro-intestinal disturbance from the first—indigestion, and occasionally vomiting. Headache is often severe.

The tendency is always to recovery, though it is obvious that, in view of the danger of oedema attacking the air passages, there must ever be considerable anxiety as to the progress of even the mildest cases. It is probable that a tendency to recurrence is a characteristic of wandering oedema, but the cases which have been recorded are too few to justify a conclusion, and more information on this point is desirable. Sufficient data are not yet before us to justify the acceptance of theories as to the causation of wandering oedema, and ascertained facts are too rare and isolated to establish its pathology.

In a case in which I was able to make a cover-glass preparation of the oedematous fluid a short round-ended bacillus was found. It has a tendency to bipolar staining, but refuses to grow on glycerine agar, blood serum, or potato. It is apparently, like the bacillus of malignant oedema, an anaërobe, but an opportunity for further cultivation experiments has not occurred, and, in the absence of definite knowledge, it would be premature to claim for it a pathogenic power. Microscopic examination of the blood does not reveal the presence of a malarial or other parasite. Localised oedemas are always difficult of explanation. No theory has as yet been accepted as accounting for the localised oedema that follows the bite of a centipede or the sting of a scorpion, though it appears now to be probable that in the oedematous fluid which is secreted there is an antitoxin and curative agency to the poisons.

Two conditions suggest themselves at first sight as being likely factors in the causation of wandering oedema—gout and urticaria; but, as the symptoms develop, it will be seen that neither of them satisfactorily accounts for the various phases of the malady. In none of the cases that are reported or that have come under my own observation has there been a history of gout or have the sequelæ justified the supposition of a gouty diathesis. Urticaria also seems to be ruled out of court by the absence of itchiness, the character of the swelling, and the progress of the symptoms.

In three of the four cases that came under my notice there was a history of mushroom poisoning. Discomfort was felt after eating tinned French mushrooms, and the attack came on within forty-eight hours in one case, and in another in seventy-two hours. In a third case, which was milder, edible mushrooms, grown in the Straits Settlements from English spawn had been eaten a week before; and in the fourth case—that of a Malay lad—though there was no definite evidence as to the mushrooms having been eaten, herbs that had been gathered for the family curry where fungi grow plentifully, had been freely partaken of. These facts would seem to afford presumptive evidence that wandering oedema is the result of fungus poison, though they are

scarcely sufficient in number to be conclusive. In any case, it is apparent that it is to a poison that the symptoms must be attributed, but whether the poison is one which has been introduced from without or has been within the body—a product of perverted metabolism—it is difficult to say. The symptoms are as far removed from the ordinary manifestations of agaricin poisoning as they are from gout.

If it is true that wandering oedema has a tendency to recur, the theory that it is due to a poison formed within the body would be considerably strengthened, while the theory of ingestion would be correspondingly weakened. But the cases that have been reported as recurrent appear doubtful instances of the disease, with the probable exception of Kruger's case,² in which the patient had repeated attacks of oedema, and died suddenly from oedema of the glottis. Some of the cases (at least of those described by Dr. Allan Jamieson in the chapter on Urticaria in his *Diseases of the Skin*, and quoted from Strübing, Quincke, and others as acute circumscribed cutaneous oedema) are doubtless identical with wandering oedema and have nothing in common with urticarial affections. But further reports of cases are necessary before the causation and affinities of this malady can be considered as at all settled; and it is to be hoped, now that attention is being directed to the subject, that these will be forthcoming.

REFERENCES.

- ¹ *Revue Médicale de la Suisse Romande*, xii, 1886. ² *Lancet*, February 9th, 1890.

SKIN ERUPTIONS OCCURRING IN PATIENTS ALREADY SUFFERING FROM INFECTED WOUNDS.

By G. G. STOPFORD TAYLOR, M.D.,

Honorary Physician Liverpool Hospital for Cancer and Skin Diseases.

IN THE BRITISH MEDICAL JOURNAL for 1892, vol. i, p. 861, I published a case of dermatitis of the eczema type beginning in a septic axillary wound, which ran an acute course, and was attended by universal exfoliation. During the last two years I have kept a record of cases in my private and hospital work which I believe originated in a similar way, namely, by the absorption of some morbid matter into the circulation.

CASE I. *Pemphigus Foliaceus*.—Hospital patient: Referred to me by Dr. Shaw, of the South Dispensary, on July 3rd, 1895. It appeared that the patient, a girl, aged 9 years, had sustained an injury to her skull by falling from a swing. On admission to hospital the whole cutaneous surface was intensely red and peeling in large dry flakes, for the most part leaving large circular areas of various sizes. In a few days her mouth became affected, the epithelium wrinkling and peeling off, exposing a raw and bleeding mucosa. On July 15th, distinct but flaccid bullæ appeared on the trunk and extremities. In spite of treatment the child succumbed on July 24th, and a post-mortem examination revealed an old superficial injury of the scalp in the left suboccipital region with the remains of extravasated blood beneath it.

CASE II. *Acute Pemphigus*.—Hospital patient: This eruption followed an injury to the hand from a steel hook in a strong, healthy young butcher. The eruption was universal, and consisted of large and small bullæ with serous contents. When admitted the wound had nearly healed. The patient was discharged cured in about six weeks.

CASE III. *Pemphigoid Eruption*.—Private patient: Seen with Dr. Moyle, of Walton. A bullous eruption of both hands, forearms, and face in a lady who was suffering from a whitlow of the right thumb. The contents of the bullæ were serous, sero-purulent, and in a few instances sanguinolent. Dr. Moyle informed me that the patient made a good recovery.

CASE IV. *Impetigo*.—Hospital patient: Developed an impetiginous eruption on the left side of the face and neck and posterior auricular region, also a pustular folliculitis of extensor surface of left arm while suffering from an onychia of the right forefinger.

CASE V. *Acute Dermatitis*.—Private patient referred by Dr. Leask, of Birkenhead: Acute dermatitis of the eczema type of left hand and forearm after a severely-lacerated wound of palm. The scrotum was also inflamed and irritable. Dr. Leask informed me that there was no further extension of the disease, and the patient made a good recovery.

CASE VI. *Erythematous Dermatitis*.—Hospital patient: A workman presented himself with an eruption of an erythematous and scaly character on the trunk, more particularly on the anterior surface. The penis and scrotum were oedematous and scaling. A few papules were present on the left forearm, and the corresponding axillary glands were enlarged and swollen. He was also suffering from a whitlow of the middle finger of left hand and suppurating of the palm, for which he had been under treatment at a general hospital, and during which the eruption appeared.

CASE VII. *Papular Eczema*.—Hospital patient: While under treatment for a varicose ulcer of the leg, developed a papular eczema on the inside of both legs, thighs, and flexor surface of forearms and elbows.

CASE VIII. *Erythema Multiforme*.—Hospital patient: While suffering from an extensive impetigo of the face, developed an erythema multiforme on the trunk and extremities.