Year.	No.	Initials.	Age.	Sex.	Cases: Duration.	Treatment.	Complications.	Result.
1879	1	A. M.	18	F.	Several years	Purgation (useless), later H ₂ SO ₄	None	Recovery.
1887	2	X.	50	M.	3 months	Arg. nit. sol.	"	"
1890	3	N.	44	F.	3 years	H ₂ SO ₄ , arg. nit. inj.	"	"
"	4	E. M.	16	F.	Some months	H ₂ SO ₄	"	Relieved, lost sight of.
"	5	P.	56	F.	Over 9 years	Various	Sore tongue occasionally, liver possibly enlarged	Chronic, little relief obtained.
1892	6	N. N.	18	F.	Some months	H ₂ SO ₄ , arg. nit.	"	Gradual relief, now fairly well.
1894	7	S.	18	F.	2 years	Various	Inordinate and unhealthy appetite indulged	Gradual recovery.
1895	"	S.	40	F.	Some months	H ₂ SO ₄ occasionally	None	"
"	9	J. E.	50?	F.	Several years	Various	"	Death.
"	10	S. S.	47	M.	"	Untreated, inquest, etc.	Scoliosis, displaced kidney	"
1896 and 1897	11	J. P.	55	M.	2 years	Various	Cancer of sigmoid; perforation	"
"	12	Lady —	—	F.	2 months	H ₂ SO ₄ , milk diet, rest	None	Recovery.
"	13	M. E.	21	F.	Several months	H ₂ SO ₄ occasionally	"	Relief.
"	14	E. G.	47	F.	8 months	Various	Hysteria	Recovery.
"	15	F. P.	32	F.	6 months	H ₂ SO ₄ , milk	Pregnancy	Almost well, under treatment.
"	16	A. P.	30	F.	5 months	"	"	Recovery.
"	17	D. D.	19	F.	10 months	"	"	Recovery, colic supervening.
"	18	P. P.	25	M.	1 year	"	"	Improving.
"	19	M. A. P.	55	F.	Several years	"	"	"

The drug I most rely on is dilute sulphuric acid; 10 minims three or four times a day, with a very little opium and a carminative, generally controls the diarrhœa quickly. When this good effect is produced the drug ought to be continued as required, a loose morning motion being followed by a dose, and should diarrhœa again come on, the mixture should be regularly taken, but never should the first dose be taken before the first motion; the distension then caused is most distressing. If sulphuric acid fails, large doses of catechu, or bismuth and salol, may be tried, and these failing, 10 grains of sodium bicarbonate, with dill water, act as a sedative if nothing else.

In certain cases, in which the descending colon is attacked, the passage of a long tube and injection of a strong solution of nitrate of silver (3ss-3i to 1 pint distilled water warmed), is an admirable remedy; it should be administered with caution, but frequently results in absolute and, I might say, immediate cure, the after-colic being but a minor evil easily combated.

Opium is merely of service during the early stages, or after diarrhœa has been checked. In some cases strychnine acts as a good tonic but the early administration requires caution.

SUCCESSFUL CÆLIOTOMIES FOR AN INTRA-PERITONEAL RUPTURE OF A TUBAL GESTATION AND TWO OTHER UNCOMMON PELVIC TUMOURS.

By J. LYNN THOMAS, F.R.C.S.,

Assistant-Surgeon to the Cardiff Infirmary, and Consulting Surgeon to the Cardiff Provident Dispensary.

THE following cases are worthy of record on account of their individual points of interest; the first one I photographed four hours after removal. It shows the full extent of the loss of tissue over the placenta in the inner third of the Fallopian tube; I skiagraphed the same, and that shows two separate dots, as if ossific centres were already formed. The second case is an undoubted true fibroma of the ovary, and from the investigations of Sir John Williams and Mr. Bland Sutton it is evidently a very rare specimen; its macroscopic appearance is well shown in the photograph. The third case is also an example of a rare condition—the impaction of a parovarian cyst in Douglas's pouch under the reflections of the rectovesical fascia.

CASE I: RUPTURED TUBAL GESTATION.

Mrs. H., aged 27, had been married only eleven weeks, and was fourteen days behind her period, when she was taken suddenly ill with severe abdominal pain as soon as she finished breakfast on May 25th, at 10 A.M. When my friend Dr. Biggs saw her a few hours later he found her in a state of

extreme collapse, with a cold, clammy surface, and obtained a history from her sister that years ago the patient had had gastric ulcer and hæmatemesis. By means of opium, rectal feeding, transfusion, and stimulants the patient was kept hovering between life and death until May 28th, when I saw her, and at that time we advised immediate operation, as she had rallied and became sensible during the last two or three hours, and hæmorrhage had also appeared from the vagina. The anterior vaginal wall was full, resisting, and depressed, more especially on the right side. She now complained of "killing, stabbing pain" on that side, was blanched, had a temperature of 102° F. Until this day the prominent symptoms were vomiting and nausea, abdominal pain, and collapse, with a running and at times imperceptible pulse. The abdomen was distended, and had signs of fluid in its peritoneal cavity; the thighs were slightly flexed. We now agreed that it was a right tubal gestation ruptured into the abdominal cavity, and gave an injection of morphine whilst the instruments were being got ready.

Fig. 1.—A, ovary; B, placenta; C, ampulla; D, fimbriae. Seventy-four hours after the first symptom Dr. Morland

administered ether, and as soon as I got my hand into the abdomen, which was full of blood, I clamped the right broad ligament with a long-curved forceps. The cavity was then rapidly cleared of many handfuls of clots and several tumblersful of fluid blood, when the uterus became visible with the forceps attached close to the ruptured tube. When the forceps was relaxed, blood poured out of the ruptured tube. The tube and ovary were removed close to the uterine cornu, and Fig. 1 (from a photograph) shows the exact size and condition of the parts. The rupture is on the anterior and upper surfaces of the Fallopian tube within $\frac{1}{2}$ inch of the uterus, and the placenta is attached to its inferior and also slightly to its lateral walls. The lumen of the tube on the uterine side does not admit a horsehair, whilst a thick probe is easily passed up to the placenta from the fimbriated extremity.

It is hardly necessary to state that the operation had to be done on the bed, as a removal of the patient would probably have been fatal. For the first forty-eight hours after the operation it required the constant care and skill of Dr. Biggs to combat the various and varying symptoms, and I am quite convinced that but for the timely administration of champagne, of opium, and for the resort to transfusion after the operation, the result would have been different. On the fifth day the patient had fish, and afterwards rapidly got well.

The left tube was normal, and the left ovary contained a large corpus luteum. Sister Yates stated in her note before the operation that on two occasions she thought the patient was dead.

CASE II: LARGE OVARIAN FIBROMA: ELEPHANTIASIS OF THE ABDOMEN.

Fibromata of the ovary are extremely rare according to Tait, Greig Smith, John Williams, Bland Sutton, and others, whilst the series recorded by Mr. H. Briggs¹ are not altogether of the pure type, and consequently are not quite so common as the title of his valuable paper would lead one to conjecture. A woman, aged 42, was sent to me by my friend Dr. Powell (Newcastle Emllyn), who had been compelled to tap her five times to relieve dyspnoea, as she would not submit to abdominal section. The result was that when I saw her in June, 1895, her abdominal wall was of enormous thickness, the skin being coarse, tuberculated, and covered with a

Fig. 2.—Ovarian fibroma. The gap in right half is the place from which microscopic section was made.

foul secretion, looking like an advanced case of elephantiasis. There was a large solid tumour in the right lower half of the abdomen, and a cyst extending up to the ensiform cartilage. The patient's suffering had been agonising for some months before operation. The abdominal wall cut was like the subcutaneous tissue of elephantiasis, and no trace of muscle was seen. The cystic tumour was attached in all directions to intestines, abdominal wall, and bladder, whilst the fibroma itself was comparatively free from adhesions. Recovery was retarded by carboloria and albuminuria. Eighteen months after operation the abdominal wall was quite smooth, there was no tendency to hernia, and the scar was 14 inches long.

The macroscopic appearance of the fibroma is well shown in the photograph reproduced in Fig. 2. It is resting on the collapsed wall of the papillomatous cyst to which it is attached. It cut like a dense fibroma, and looked typical of one. The report of the Clinical Research Association is: "The tumour is a dense fibroma composed of wavy fibres arranged in interlacing bundles." The section was taken from near the periphery of the tumour (shown in the figure on the right hand). It weighed $5\frac{1}{2}$ lbs.

CASE III: PAROVARIAN CYST IMPACTED IN DOUGLAS'S POUCH.

The causes of impaction of cysts in the pelvis are commonly the following: Adhesions, malformation of the pelvis, gravid uterus, and rotations; but in the following case I believe it was due to confinement under the reflections of the recto-vesical fascia and peritoneum.

Miss —, aged 20, had been treated with pessaries for nearly four years for prolapsus uteri. There was a history of difficulty in micturition for six years. Menstruation occurred first at 16, and had been irregular ever since every three or four weeks. There was in Douglas's pouch a firm cystic swelling which moved with the uterus. The uterine cavity was 3 inches long.

On September 8th, 1896, I opened the abdomen in Trendelenburg's position, and found a tumour behind the uterus, the fundus of which was higher than the cyst; I failed to release it until one of my colleagues introduced two fingers into the rectum, when after three or four strong pushes the cyst suddenly darted out into the wound and was easily removed. The cyst was the size of a lady's fist; was parovarian in origin, and had no adhesions; it had two atrophic patches the size of a threepenny bit on two opposing surfaces, probably due to pressure against the lateral walls of the pelvis. It is rather strange that she had had no trouble in defaecation. The fundus uteri was thickened. It is probable from the history that this parovarian cyst commenced at an unusually early age, for Mr. Bland Sutton in his *Surgical Diseases of the Ovaries* states that he has not found one recorded under the age of 16.

REFERENCE.

¹ BRITISH MEDICAL JOURNAL, May 1st, 1897.

REMARKS ON THE TREATMENT OF CUT THROAT,

WITH NOTES OF A CASE TREATED BY IMMEDIATE SUTURE IN LAYERS.

By R. G. HOGARTH, F.R.C.S.ENG.,
Salisbury.

Late House Surgeon, St. Bartholomew's Hospital; and
Senior Resident Medical Officer, General Hospital, Nottingham.

OUT of seven cases I have had to treat, the following is the only one of which I have full notes. Of the others, in one the wound was situated above the hyoid bone, three were through the thyro-hyoid membrane, one through the crico-thyroid membrane, and one through the trachea. Unfortunately the immediate suture treatment was not adopted in all the cases.

J. R., aged 48, was admitted to the General Hospital, Nottingham, on February 12th, 1896, at 10 A.M. The skin was cold and clammy, the pulse rapid and feeble. He was covered with blood and vomit. There was one extensive wound in the neck, deeper on the left side than on the right, opening the larynx and pharynx by dividing the thyro-hyoid membrane. The epiglottis was completely divided at its base, and the middle constrictor of the pharynx was cut on each side. The thyroid cartilages were uninjured. The sterno-mastoid muscles were notched, but the large vessels had escaped injury. The wound gaped a great deal, and through it bubbled with expiration a mixture of saliva, vomit, and blood; by the look of the tissues the wound had been made some hours, and there was only a little oozing of blood, the primary hæmorrhage from the division of the superior thyroid or lingual vessels having stopped. The larynx and pharynx were well sponged out, and made as clean as possible, all hæmorrhage carefully stopped, and the severed tissues of the larynx brought together and fixed in position with chromic catgut. The epiglottis was done first, and came well and easily into position. The air passage was apparently rendered quite tight. The whole wound was then well washed out with biniodide of mercury solution, 1-500 first, then 1-2000, and the cut tissues united layer by layer with chromic catgut, except the skin, which was brought together with horsehair. A small drainage tube was inserted on each side at the ends of the wound, and an antiseptic dressing applied.

At 6 P.M. the temperature was 101°, the pulse 90 and of good volume. He was fairly comfortable. Now and then breathing was noisy. He had morphine $\frac{1}{4}$ gr. early in the day. He coughed very little, but when he tried