I have brought these matters before you so that you may understand some of the reasons of the outcry which is raised against the Army Medical Service. Even with these drawbacks many will find it their interest to go forward for appointments. I by no means blame them. The competition of life is very keen, and I do not advise anyone to hold back. Perhaps I shall be condemned for this statement, but who, I ask, will provide an equally secure position for the young men who follow the counsels of another kind? I know of none; and it seems to be asking too much of our students to stand back, and thereby perhaps sacrifice a life's career. Each knows his own necessities and desires. We will do all in our power, as we have done, to help forward army medical reforms, but we must not speak hardly of those who, in spite of disadvantages, seek entrance into the service. At the same time the authorities must know that many brilliant men are looking elsewhere for a career, and that it depends upon an enlightened policy whether the medical department of the army will be again largely recruited from among the most promising of our young surgeons.

THE ACTION OF THYROID EXTRACT ON THE SKIN AS ILLUSTRATED IN CASES OF ICHTHYOSIS.

By WM. WALTON DON, M.D.,
West Hampstead.

THE following cases illustrate the general action of thyroid extract on cutaneous surfaces, and, at the same time, show its special remedial effects in ichthyotic conditions.

In chronic skin diseases in which the extract has been given with good effect the question is still open whether its use can be approved of as a routine treatment, for it still remains to be demonstrated whether the continued use of the remedy in such diseases would not ultimately prove injurious to the general health. The rapid pulse which the internal administration of thyroid extract always induces must, I think, in the end prove hurtful to the cardiac function, and therefore its too prolonged use is probably to be deprecated. Its remedial effect in skin diseases appears to result from increasing the cutaneous vascularity, thereby stimulating dermal nutrition, and also through diaphoresis getting rid of effete material. Certainly this action on the skin is always assisted by external stimulant applications, such as baths and inunctions.

The action of thyroid extract in ichthyosis seems to be so beneficial that if any constitutional treatment in that disease is attempted it should be given a fair trial. It does not seem to have been much employed in ichthyosis, which is the more remarkable when its peculiar action on the skin is considered.

In both of my cases the results were temporarily excellent, but on leaving off the remedy the diseased condition of the skin showed liability to return.

I have found in administering the extract to patients that they display tolerance of a considerably greater dose after its administration has been continued for some time, and in this fact apparently lies the best chance of successful treatment, because with a continued limited dose only a certain point is reached beyond which no visible improvement apparently results, but on increasing the dose (provided the patient can bear it) improvement, if it is going to take place at all, very shortly appears.

The form in which I have given the remedy was Messrs. Burroughs and Wellcome's 5-grain tabloids, commencing with one tabloid, and after three or four days increasing to three or five daily, according to the effect on the patient. At the same time I gave a cardiac tonic containing nux vomica and digitalis at the commencement of the treatment.

The following notes taken during the treatment of two cases of ichthyosis show in a general way their progress:

G. T., a boy, aged 12 years, had suffered from scalliness of the skin since birth. There was a hereditary history—grandfather had suffered similarly all his life; while a maternal uncle was very rough skinned, and subject to cutaneous fissures. The areas most affected were the back of the arms, flanks, hips, thighs, and shins; but on the body the skin was also scaly, bran-like, and easily brushed off; on the shins and outside calf of leg, the fissures in the skin formed diamond-shaped patches, from

1 inch to $\frac{1}{2}$ inch in size. Face ruddy, covered with light fine scales—like the peeling in scarlet fever. His condition was much aggravated in winter, when cold winds prevailed. At times large patches of epidermis were shed. In spite of washing, the parts affected very quickly became dirty again, and could not be properly cleaned till the epidermis was detached; curiously, he was said to perspire freely—mostly about the head and face. There were very few nutmeg-grater areas, only those on the back and posterior parts of the arms.

September 12th. Pulse 96; temperature 98.6°. Commenced taking two tabloids (= 10 gr. ext.) a day.

September 15th. Pulse 104; temperature normal, but complained of a hot feeling of the skin. To take three tabloids a day (= 15 gr. ext.)

September 19th. Pulse 100; vomited twice on the 17th; temperature 99.8°. Tight feeling, and aching over the eyes.

September 26th. Pulse 140; temperature in axilla and mouth 98°. Felt very well, and no excessive sweating, but skin very hot and red looking in places.

October 3rd. Pulse 96; temperature 99.4°. Vomited on September 29th; and on October 1st and 2nd had considerable diarrhoea and nausea (a fortnight after commencement of treatment).

October 10th. Feeling well. Pulse 122; no sweating; condition of skin slightly improved. To take four tabloids a day (= 20 gr. ext.).

October 14th. Pulse 120; no unpleasant symptoms. To take five tabloids a day (= 25 gr. ext.).

October 17th. Pulse 125; much sweating; appetite very good; and condition of the skin improving.

October 24th. Pulse 122; sweating abated.

October 30th. Pulse 120; sweating again great, but skin rapidly improving.

November 7th. Pulse 120; less sweating, skin smooth on face, arms, and neck, from which the scales can be easily brushed off. Ordered sponging with glycerine and water (1 in 20) to remove scales.

November 14th. Pulse 133. Every portion of previously affected skin now smooth, soft, and supple; but those areas originally most affected present a transparent reddish-blue appearance.

November 21st. Left off the treatment during the past week; scurfy condition slightly returned.

November 28th. Pulse 126. Taking three tabloids (= 15 grs. ext.) a day for past week; the slight scurfiness now gone.

December 5th. Pulse 120; skin quite clear and healthy; no unusual sweating. Treatment discontinued.

December 20th. Pulse 90; skin normal, except on forearms, where it is somewhat coarse-looking.

F. L., girl, aged 4 years. The disease of the skin had existed since she was 9 months old; was worse in cold and better in warm weather; exposed areas most affected; a well-nourished and rosy-cheeked child. I first saw her on June 1st, when recovering from measles. The face was covered with small, fine scales; back and upper arms very rough—like an aggravated "goose skin." The disease worst on shins and knees, the skin over these surfaces covered with patches of greenish-grey detachable scales; between these areas deep cracks ran, nearly at right angles, giving a diamond-shaped appearance. The measles had greatly aggravated the skin disease; although at times previously it had been as bad. Had been treated with glycerine inunction, with slight temporary improvement.

On July 1st the legs were better, with fewer flakes; but the outer surface of the upper arms, back, and thighs felt nutmeg-grater-like, while the shins had an appearance as of chronic dry eczema. Unfortunately, no photograph was taken of the condition of the skin at this period. Pulse 100.

July 3rd. Began treatment; to commence with $\frac{2}{3}$ gr. thyroid extract, increasing to 5 gr. on the 4th, 10 gr. on 5th, and 15 gr. on July 6th.

July 20th. Had continued with 15 gr. daily till July 13th, and then ceased. To recommence remedy.

July 31st. Pulse 130. Arms now greatly improved; the nutmeg-grater touch less, and the small papules thereof could be rubbed off with the finger; fissures on the legs less. The child irritable and sweating greatly, but with good strength. Urine passed in excess.

August 8th. Taken 5 gr. thrice daily since July 20th; although weather colder, general condition much improved; shins show marked improvement; skin on the back and thighs now soft and supple; the face, however, retaining a dry eczematous condition, similar to that often seen in fair-skinned young children; not sweating so much. Glycerine, an ounce, to be added to a small bath, given twice weekly, and the scurf to be removed as much as possible.

August 25th. Very rapid improvement during the last fortnight; skin over shins, back, arms, and thighs nearly normal, but on face still dry and eczematous-looking.

September 1st. Skin quite healthy; treatment discontinued; patient going to the country.

January 4th, 1897. Patient again under observation; the ichthyotic condition has returned, though not so bad as before treatment.

From the various cases in which I have given thyroid extract, and from a careful scrutiny of the bibliography of the subject, I have come to the following conclusions regarding its action on the skin.

There is no doubt it is strongly stimulating by directly increasing the cutaneous circulation, as evidenced in sensations of flushing, hot tingling, and congestive irritation, such as is felt preceding ordinary perspiration.

The increased cutaneous vascular supply apparently results in: (1) Increased nutrition of the skin, hence its probable remedial action in ichthyotic conditions, an effect produced without any necessarily abnormal perspiration; (2) increased action of the cutaneous glands, accelerating excretion of waste products, thus keeping the surface in a supple condition; (3) regrowth of hair, as shown in myxedema and some cases of general alopecia; (4) increased activity of the epi-

dermal layers, causing desquamation of unhealthy epidermis and reproduction of a new covering, as observed in ichthyosis, psoriasis, dry chronic eczema, and also in some cases of myxœdema and cretinism.

Among its general effects I have found that in doses of 15 gr. daily it causes a pulse-rate of 120 to 140 per minute, which in different subjects may be either of high or low tension. But thyroid extract seems also to rapidly exhaust latent cardiac energy, for if the dose be unduly increased, or if the patient is not fully accustomed to it, then the heart soon tires, and the pulse, although still quick, is weakened so as to induce irritability, depression, and tendency to syncope. It is difficult to say whether increased excretory action be due to direct irritation of the secreting glands or from general increased blood pressure; possibly it arises from both.

In neither of the above cases did the temperature rise high; there was indeed a slight rise at the commencement of the treatment, but none in its later stages, although the remedy was given in larger doses, a result probably due to increased metabolism of the tissues, plus insufficient surface cooling, from the skin not acting in a normal manner, for when the skin became healthier there was no appreciable thermometric rise.

The action of thyroid extract on the nervous system in most of my cases has been depressing, patients complaining of feeling debilitated and low; but in a case of myxœdema in which I used it its action was the very opposite.

In the above cases the results are, I think, sufficiently encouraging to warrant a more extensive trial in ichthyosis, which has hitherto proved so intractable to treatment.

SKIN ERUPTIONS AND INFECTED WOUNDS.

By DAVID WALSH, M.D. EDIN.,

Physician, Western Skin Hospital, London.

THE short article by Dr. Stopford Taylor in the *BRITISH MEDICAL JOURNAL* for September 25th, 1897, calls attention to a point of importance not only in dermatology but also in general medicine. Briefly put, he presents a number of cases of dermatitis of varying type associated with septic inflammatory conditions of the skin, either of traumatic or non-traumatic origin. Such rashes, in his opinion, arise from the absorption of some morbid matter into the circulation. As that view tallies exactly with conclusions published some years since by the present writer, a few comments may perhaps have a suggestive or epexegetical value.

First of all there is of course the possible fallacy that the dermatitis may result from direct extension and surface inoculation of virus from the primary skin lesion. However, we find a large number of severe symptomatic rashes, such as those resulting from drugs and exanthems, where no such external inoculation could take place, and which therefore offer a strong analogy to Dr. Taylor's assumed absorption of septic matter. How the irritant is to affect the skin after its entry into the blood is not discussed by the observer in question. The present writer's theory is that the dermatitis results from direct inflammation of the skin by the excretion of a bloodborne irritant. A similar explanation would account for the skin phenomena met with in connection with certain drugs, exanthems, and various septic invasions. Moreover it would have a bearing upon the constantly accompanying injury to other organs of elimination as the bronchi, intestines, and kidneys.

Dr. Taylor mentions a case previously described by him¹ "of dermatitis of the eczema type beginning in a septic axillary wound, which ran an acute course, and was attended by universal exfoliation." His attention has doubtless been drawn to the curious skin eruption in a steer, described by Dr. Eddowes² at the Dermatological Society of Great Britain and Ireland. The beast had a septic operation wound, and was attacked by a severe desquamative dermatitis of the white parts of his body, while, curiously enough, the red were not affected. A white star on the forehead was not involved, a fact that Dr. Eddowes thought due to its not being licked by the animal. At the same time he compared the case with Savill's epidemic dermatitis. Of the latter disease, which presents a most important series of problems to the derma-

tologist, the present writer has spoken as "an epidemic dermatitis where the desquamative lesion has all the appearance of resulting from the cutaneous excretion of an intensely irritant product of obscure origin."³

At the end of his article Dr. Taylor asks⁴: "Do not the foregoing cases suggest to us that many skin affections which have been attributed to vasomotor disturbance and reflex irritation are due to the absorption of micro-organisms or their products either by the skin or alimentary tract?" The same question was asked by the present writer in 1890 at the Birmingham meeting of the British Medical Association in a paper entitled, *Dermatitis as an Excretory Symptom*.⁵ A comparison was then made between various rashes where the common feature was an inflamed skin and the presence of some bloodborne irritant, whether drug or specific virus. Whether or not the true etiology of symptomatic dermatitis may be ultimately found in excretory irritation, it is not a little curious that none of the existing theories are capable of explaining such every-day phenomena. So far as the present writer has been able to test his general conclusion by particular applications, it seems to afford a simple explanation to such varied rashes as those associated with scarlet fever and the rest of the exanthems, antitoxin and tuberculin injection, arsenic, iodine and a host of other drugs, gout, rheumatism, and with septic conditions of the body, local and general. In all these instances there is a marked tendency to vicarious inflammation of eliminating organs other than the skin, a disturbance which it is suggested may have been caused by the passage of an internal irritant through the essential cells of the excretory organs.

REFERENCES.

- ¹ *BRITISH MEDICAL JOURNAL*, 1892, vol. 1, p. 861. ² *British Journal of Dermatology*, p. 54, February, 1895. ³ *Excretory Irritation*, Walsh, 1897, p. 49. ⁴ *BRITISH MEDICAL JOURNAL*, September 25th, 1897, p. 805. ⁵ *Med. Press and Circular*, October 22nd, 1890.

NOTES ON FIFTEEN CASES OF INTUSSUSCEPTION.

By CHARLES P. B. CLUBBE, L.R.C.P. LOND.,
M.R.C.S. ENG.,

Honorary Surgeon to the Prince Alfred Hospital; Honorary Surgeon to the Sydney Hospital for Sick Children; Joint Lecturer on Clinical Surgery to the Sydney University.

In the 15 cases of intussusception that I now put on record, 8 recovered and 7 died. In 3 cases the intussusception was reduced by injection of oil without operation. In this series, then, there were 12 laparotomies, with 5 recoveries. In one of the cases that recovered, 4 inches of the small intestine was resected, and this in a child under 12 months. I have previously reported 7 laparotomies with 4 recoveries. Taking the operations in the two series of cases together, then, the figures are 19 cases, 9 recoveries, 10 deaths. The cases are given in the order in which they occurred.

CASE I.—E. G., aged 6 months, admitted to the Children's Hospital July 10th, 1895. History: Thirty hours ago screamed and turned pale. Passed blood eight hours later. Vomited several times. Under chloroform a tumour felt on the right side of the abdomen. Warm olive oil was injected, and the mass disappeared. In a few hours the child passed three motions; no vomiting. Child was discharged the next day, as it was not weaned, but it was kept under observation for several days, and remained quite well.

CASE II.—K. R., aged 5 months, admitted to the Children's Hospital October 6th, 1895. Child ill seventy-two hours. Attack began with screaming. Then vomiting, and soon after passage of blood. The child has been constantly vomiting and straining since, and passing blood and slime. On admission mass could be felt in rectum. The abdomen was opened in the middle line, and as it was impossible without undue violence to reduce the intussusception (which was caecum into colon) completely, the mass was resected and the ileum was sutured to ascending colon. The child died soon after the operation (which only lasted thirty minutes) was completed.

CASE III.—O. W., aged 1½ year, admitted to the Children's Hospital October 17th, 1895. Ill five days; constant vomiting; no motion for five days. On the second day of illness some slime and blood were passed once, but not again till to-day. On opening the abdomen intussusception was found to be a double one, ileum into ileum, and then this into the caecum. In reducing these intussusceptions the ileum was torn. The damaged part, about 6 inches of ileum, was therefore resected, and the cut ends of the bowel united with a double row of catgut sutures. The child only lived one hour after the operation was completed.

CASE IV.—E. C., aged 5 months, admitted to the Children's Hospital on July 30th, 1896. Began to scream two days ago, and soon vomited and passed a motion consisting chiefly of blood and slime. Under chloroform cylindrical tumour, extending boomerang-like, in region of hepatic