outcome of the lifelong attention I have given to the subject upon which I have addressed you. The views I have set forth have sprung entirely from experimental physiological work on the one hand, and from practical experience in connection with diabetes on the other. With this foundation, they harmonise with the assemblage of facts to be dealt with belonging alike to physiology and pathology; and, at one and the same time, they bring the whole matter into a state of great simplicity, and supply a working basis upon which the treatment of diabetes may be rationally and reliably conducted. With the glyco-genic doctrine the teachings of diabetes are absolutely irreconcilable, and it may, moreover, be said that modern research has removed the support upon which it was originally based. And yet the doctrine still lingers in the mind, influencing its conceptions and shaping its views, for persons seem unable to look at the points belonging to the subject otherwise than through the prepossessions that have been engendered thereby. The chapter, however, will have, sooner or later, not simply to be revised, but completely rewritten; and the sooner this is done the better, I feel I am justified in saying, will it be for the right comprehension of diabetes by the medical profession.

CASE OF PYOPNEUMOTHORAX OF SEVERAL MONTHS' DURATION CURED BY FREE INCISION: WITH SOME REMARKS ON THE SURGICAL TREATMENT OF PNEUMOTHORAX.*

By SAMUEL WEST, M.D., F.R.C.P.,

Assistant Physician to St. Bartholomew's Hospital, and Senior Physician to the Royal Free Hospital.

I WILL in the first place relate the history of the case to which I desire to call attention, and afterwards make some remarks on the debatable points which are raised by it.

W. T., aged 34, a blacksmith, was admitted to the hospital under my care on March 8th, 1897, complaining of cough and difficulty of breathing. He proved to have a pyopneumothorax of the left side.

His previous history was this: On November and, 1896, he was taken suddenly ill while at work, with sharp pain in his left side, which was unbearable on coughing. He went home to bed, and stayed there for eleven weeks. He was told that he had pneumonia and pleurisy. Shortly after Christmas time, soon after getting up, he had an attack of violent dyspnoea, during which he became unconscious for a short time. He had had two slighter attacks since, and had been very short of breath on any exertion. For the last six weeks the patient had noticed a peculiar splashing sound in his chest as he walked, and he had shooting pains in different parts of the left side of the chest. His cough during this time had been frequent and paroxysmal. There had been much dark, thick expectoration but no hæmoptysis.

On January 4th the patient went to the Convalescent Home at Sandgate, where he remained five weeks, and while he was there the doctor drew off a small quantity of clear fluid from his chest.

Returning from Sandgate, the patient went to St. Bartholomew's Hospital, where he was shortly after seen by me and recommended for admission. During his illness he had lost two stone in weight, but he lost three during the first part of his illness, and regained one subsequently at Sandgate. There was nothing in the patient's previous history or in the family history which had any bearing upon the case. The patient looked pale and ill, and had evidently lost a good deal of flesh. He was disturbed frequently by paroxysms of coughing, which were occasionally very severe. He presented the signs of left-sided pneumothorax, the cardiac dulness reaching as far as two inches to the left of the sternum.

* Read before the Medical Society of London, November 22nd, 1897

The diagram shows the limit of hyper-resonance as the patient lay upon the back; in front the voice and breath sounds were amphoric in character, and the bell sound audible over the whole hyper-resonant area; behind, the breath sounds were absent at the base and altered in character at the apex, and the bell sound stopped at the limits shown upon the diagram; above and below between the two lines the bell sound and amphoric breathing and voice sounds were present, as in front. Succussion was easily elicited. The lung, it was thought, was adherent at the apex behind. The hole in the lung, in spite of the amphoric breathing, was evidently closed. As a cyrtometer-tracing showed, the affected side was rather smaller than the opposite one.

March 16th. A needle was inserted into the seventh intercostal space behind in the mid-axillary line and sero-purulent fluid obtained.

The patient had a pneumothorax with sero-purulent effusion. He was very ill, and quite incapacitated for work. He had been invalided for some months, and was getting steadily worse.

The question was: How was he to be treated? The fluid might be removed by paracentesis, but experience of similar cases was not encouraging. He could be left alone, but it was pretty evident what this would end in, and that to do nothing was tantamount to permitting the man slowly to die. Free incision seemed a better alternative than this, for it would give the patient, at the worst, a chance, even though so much time had been already lost. My own opinion was in favour of the treatment of pyopneumothorax on surgical principles by incision, and I did not take the unfavourable view of the prognosis after incision which was usually taken. Finally, if the side was to be opened at all, the sooner it was opened the better.

Accordingly, after giving the patient the advantage of a few days' rest and feeding in the hospital, I decided to have the side freely opened.

On March 30th a simple incision was made into the side in the sixth space in the posterior axillary line; a large amount of sero-purulent fluid escaped, probably about 4 pints. An examination with the probe immediately after the operation was completed showed that the lung was not more than an inch from the chest walls everywhere, so that it had expanded directly after the operation. Directly the side was opened the patient began to cough, and coughed violently. This might have been due to the irritation of the tube which was inserted, but was more probably owing to expansion of the lung. The patient took chloroform well, and bore the operation satisfactorily. A few minims of laudanum were given on the tongue to check the cough.

March 31st. The patient had been comfortable after the operation, except that once or twice during the night he seemed a little faint, and required some brandy. The next morning, when the wound was dressed, air was found to enter the whole of the left lung, and friction sounds were audible not far from the seat of incision. The patient complained of a good deal of pain in the left side, which was probably connected with this friction.

The discharge from the side was very free, so that for the first few days the patient had to be dressed twice daily. After this the discharge decreased greatly, and it was sufficient to dress the side once daily.

On April 7th the discharge became very offensive, some small pieces of sloughing tissue were removed from just inside the wound, and it was thought that a spot of bare bone was felt by the probe. The side was washed out with a weak solution of iodine and with boracic acid; the discharge rapidly lost its offensive smell, and the spot where it was thought the bone was bare healed over.

From this time the discharge rapidly diminished in quantity, and on April 25th it was about 2 or 3 ounces daily, and it continued to this amount for a good long time. Examination with the probe showed that the lung was in contact with the chest walls all round the incision, but that the track in which the tube lay extended for a distance of about 5 or 6 inches straight into the chest in the direction of the spine.

On May 15th the patient began to complain of a good deal of pain in the chest, sometimes round the wound and sometimes a little deeper, but no cause for it could be ascertained. The sinus had, up to the present time, been regularly washed out with carbolic acid solution. This was now stopped, and the pain was a little relieved, but as the pain continued various experiments were tried with the tube, as it was thought to be connected with some irritation the tube produced. The tube was shortened, but the pain was not much relieved.

On June 16th, on probing, a tender spot was found far back, and it seemed to me that at this spot the tube pressed on an intercostal nerve, and thus caused the pain. The tube was removed, and a longer one of smaller diameter inserted, and from this time the patient suffered no more pain. On June 25th it was stated that the patient felt very much better; he had been up some little time, and he managed to go up and down stairs moderately well; the amount of discharge was still about the same, but not more than could be accounted for by the irritation of the track of the tube. On July 8th he had some pain in the leg, which proved to be a little periostitis. This was rapidly cured with belladonna and glycerine, and gave no further trouble. The discharge was now not more than about an ounce in the twenty-four hours; the patient was up daily, was gaining strength and weight.

On October 12th the patient's weight was 10 st. 11 lbs.; the discharge was very slight, but, as it did not completely cease, the sinus was injected with creosote dissolved in olive oil (25 per cent. solution). Although this caused a good deal of smarting, it seemed to do a great deal of good. By the end of the month the tube had been considerably shortened, and by November 1st the tube was out, and the wound completely healed. On November 1st I made the following note: "The patient is in excellent health, looks fat and well, and says he never felt better in his life, and feels quite able to go back to his work, that of a blacksmith." The side was a good deal contracted, but no very large amount of deformity had

been produced. The vocal vibrations and vocal resonance were felt right down to the base behind. The percussion, at the lower parts especially, was still somewhat impaired; the breathing was audible over the whole side, though of course not as loud as on the opposite. The temperature was normal. The patient had gained 2½ lbs. during the last week.

As regards the temperature, all that need be said is this: that at the time of admission and up to the date of operation the temperature was normal, or rather somewhat subnormal, and remained subnormal for the first few days after the operation; it then began slowly to rise, and remained moderately hectic, varying daily from about 102° to 99.5° for about a week or ten days, after which it became subnormal again and continued so, with few occasional rises for a day or two, throughout. The patient may be regarded now as completely cured.

REMARKS.

Here is a case of pneumothorax of several months' duration with a purulent effusion. The patient had been under the care, shortly before I saw him, of an eminent authority, who had dismissed him as incurable; yet, as the result has shown, the case was not incurable when treated on general principles.

We may ask why are such cases as these regarded as incurable? Why are they left alone, that is to say, left to die, without the chance of recovering being given them which surgical interference might provide? The answer is—because of certain theories which are commonly accepted and frequently acted on, and which I believe to be erroneous.

1. That where the lung has been collapsed and compressed for some time, whether by air or fluid, it will form adhesions, and be bound down and be thus incapable of expanding again, so that if the side be opened a large cavity will be left, which cannot contract, and can only heal by the falling in of the ribs on one side, and by granulating up on the other. If this theory were true, of course operation would be undesirable, for there would be a chronic cavity left which could not close, and the patient would suffer from the results of prolonged suppuration, and die probably of exhaustion or of amyloid disease, and this is what is stated commonly to occur.

Now, is it correct to assume that when the lung has been compressed in the way I have described it must necessarily be bound down and be incapable of expansion? Can we make this a general statement or general rule of practice and act upon it? Certainly not. Of course it may be admitted at once that in many cases where patients have died after long-standing empyema, or pyopneumothorax, or sometimes even with serous effusion, the lung has been found tightly bound down by dense adhesions, and therefore would have been incapable of expanding and healing; but it must be remembered that these cases have been left alone with nothing done, and it does not follow that if some operation had been performed in early times the lung would have been bound down in that way.

We must also admit that there are cases in which very shortly after an inflammation of the pleura has taken place, as for instance with a common serous pleurisy, the lung has contracted adhesions enough to considerably hamper its expansion. These cases however, again, are quite exceptional, and we have daily evidence of the fact that the lung may be collapsed for a considerable period of time by serous effusion, and yet not contract these adhesions. Though the plea that these adhesions were likely to form was used as an argument some years ago in favour of the early performance of paracentesis in serous effusion, experience has shown that this fear is not well grounded.

That the lung is not bound down, even after a very considerable lapse of time, both in serous and purulent effusions, I will quote three or four cases to prove.

A young man, aged about 24, had been discharged from the army in July because he had a discharging empyema, which had made its exit through the side and also through the lung, so that he was expectorating about a pint of pus daily, and losing about a similar amount daily through the opening in the side. He came under my observation six months later, and as he was in good general health I decided to perform an operation upon him before his health began to fail. Accordingly a free counter-opening was made low down in the posterior part of the axilla so as to provide free drainage.

Within about a week from this time all expectoration through the lung had ceased, and within five weeks from the time of operation the wounds were closed and healed. The lung had expanded well, and breathing was audible over the

whole side. The patient came to see me for some time after that, and I know that he remained in excellent health for some months as long as he was under my observation, and I have no reason to believe that his health suffered afterwards.

Three years ago I read to this Society an account of the case of a lady who was suffering from a large right-sided serous effusion which had been deliberately, and after consultation, left alone for a period of eighteen months without even once being tapped. I tried paracentesis several times, and when this failed I ultimately decided to have the side laid freely open and the case treated as if it had been one of empyema. Paracentesis had shown me that the lung was not bound down as I had at first had reason to believe, for after each paracentesis it came out into contact with the ribs. I had hoped, however, after the repeated paracenteses, that the lung would have formed adhesions with the costal pleura, but to my surprise, when the side was opened, no adhesions were present, and the lung collapsed completely. However, this made no difference in the ultimate result, for the lung rapidly re-expanded and came into contact with the chest walls, and formed adhesions there. Soon nothing was left but the track in which the tube lay, and ultimately this also healed.

This case shows, first, that the lung in a case of serous effusion may be collapsed for at least eighteen months, and still be capable of ready re-expansion; and, secondly, that even after repeated paracentesis no adhesion need necessarily take place between the two layers of the pleura.

I have recently had another case of the same kind in a man who had been allowed to have a serous effusion not operated upon for about twelve months before he came under my observation. With him I tried also the effect of repeated paracenteses, but, encouraged by the experience of the previous case, after a few paracenteses I decided to have the side opened. Accordingly the side was opened, the fluid let out, and a drainage tube inserted. Within a week from the time of operation the lung was in contact with the chest walls over the whole side, and nothing was left except the long sinus in which the tube lay.

As I propose shortly to bring this case before one of the Societies, I need not describe it further here.

In the case which is the subject of this paper, again, although the lung had been compressed for some months, within a week after the incision had been made in the side the lung was in contact with the chest wall over the whole side, and the pleural cavity closed except for the sinus in which the tube lay.

We have, therefore, a series of cases: one of empyema, two of serous effusion, and one of pneumothorax, in which, in spite of the lung having been compressed by effusion for a considerable period of time, it was not bound down, but was capable of ready re-expansion when placed in a position to expand.

These are extreme cases, and although, on the other hand, it must be admitted that the lung may form adhesions which may bind it down quite early in a case of pleurisy, still there is no evidence that this always occurs, or even that it is the general rule; on the contrary, the evidence all points the other way. If this theory be unsound, the practice based upon it will be wrong.

2. A second theory, which is commonly stated in textbooks and is very widely accepted, and which has been often used as an argument against surgical interference, is that the compression of the lung by effusion or air, as the case may be, checks the progress of tubercle in it; so that though the presence of effusion or air in the pleura is an evil, it is outweighed by the advantage of checking the development of tubercle in the lung, and therefore no operation should be performed.

This again is, I believe, a completely erroneous theory though the statement is commonly made and has many distinguished names to support it; still I do not know that it rests upon any conclusive evidence, but remains a matter of opinion only; indeed, what evidence there is leads to the opposite conclusion. It is not the case that the compression and collapse of the lung does necessarily check the development of tubercle in that lung, for tubercles of recent formation may be found in the lung which is collapsed, and of date, as far as can be judged, subsequent to the time when the lung became collapsed. Certainly the collapse of one

lung is not infrequently followed by the rapid development of tubercle in the other.

The same theory was used at one time as an argument against the performance of paracentesis for serous effusion, and no doubt it was under the influence of this theory that the two cases I have recorded were deliberately not interfered with; but as a matter of fact at the present day this theory does not really affect our practice at all in respect of pleural effusions. They are tapped, and I believe rightly tapped, without any consideration as to the effect the removal of the fluid may have upon tubercle in either the one lung or the other.

In the case of empyema again we have almost forgotten that such a theory ever had any influence at all. No one nowadays thinks of leaving an empyema alone because of some possible effect it might have upon tubercle in the lung.

I maintain that these theories should have no more weight in relation to pneumothorax and its treatment than they have in relation to the treatment of empyema or of serous effusion.

3. Another argument that is used as against the treatment of pneumothorax by free incision, is that experience shows that the results are unsatisfactory. "Leave pyopneumothorax alone because the patient will die any way, and operation will only accelerate that result," used to be the teaching of not many years ago; and indeed on this point Fagge, who can hardly even yet be regarded as antiquated, was very precise.

"Operate only when driven to it," was another form the advice sometimes took. What would be the result upon our statistics if the same were applied to empyema, and if we never opened a side until the empyema was pointing, and never touched it even then if the empyema by good luck would discharge through the lungs?

A question of this kind must be settled by experience, and we cannot take the results of some years ago as a guide to practice at the present day in pneumothorax any more than in empyema.

In this respect, namely, the operative treatment of pneumothorax, we have to make our own experience over again; we cannot be guided by the experience of many years ago; especially when we remember that the statistics are based upon the cases of pyopneumothorax which were never operated upon unless operation became absolutely imperative.

I think I have shown, therefore, that there is no valid theoretical reasons why sero-purulent and purulent effusions in pneumothorax should be treated in any different way from sero-purulent or purulent pleuritis, and that the presence of air should not make any material difference.

Of course it is quite true that in the majority of cases of spontaneous pneumothorax the cause is a tuberculous cavity near the surface of the lung; but these ruptures frequently heal, and there is no reason why, when the rupture has firmly healed, the lung should not come into contact with the chest again, and the condition of the patient return to that which existed before the pneumothorax occurred.

Physicians have been in the past, it is true, too much inclined to leave things alone, though I do not think we can say the same of the present day, and it will be something in pneumothorax if we can brush aside the erroneous theories which have hitherto given an excuse for avoiding surgical interference.

If pneumothorax may be treated by incision when necessary, it next remains to consider when it becomes necessary.

1. *During the Early Stage.*—It is in the early stages of pneumothorax that the risk to life is greatest, and it becomes less and less as time goes on. This is due to the sudden embarrassment of the respiration and circulation, caused by the sudden collapse of one lung and the congestion of the other. If life is prolonged and time is given, the lungs and heart adjust themselves to the altered conditions and the urgent symptoms pass off.

To relieve these urgent symptoms in the early stage, paracentesis is often necessary, and may have to be repeated. If, after paracentesis, the conditions again became urgent in a very short space of time, so that paracentesis has to be repeated at short intervals, it may be wise to consider well whether the side should not be incised. This would have the

effect of giving a permanent vent to the air, and relieving all pressure symptoms; but it would have two obvious disadvantages: first, the lung, being partly expanded, would be in a less satisfactory condition for the perforation to heal soundly; and, secondly, an incision would almost certainly end in this case, as in others, with the production of an empyema. It is wise for these reasons to entirely avoid, if possible, a free incision in the early stage of pneumothorax; or, at any rate, postpone it as long as possible. All the relief that is requisite can usually be given by means of repeated paracentesis. However, if paracentesis should fail, I should not myself hesitate, even in the early stage, to lay the side open; and I should be inclined to do this in preference to using any form of valve tube. The only tube of this kind which I should be inclined, under any circumstances, to employ would be a fine trocar connected with a small india-rubber tube, extending, say, to the floor from the bed and opening under water; in this way a fluid valve would be formed which would permit the escape of the air from the pleura, but not allow any air to pass back. The objection to any of these tubes is that, do what we will, in most cases the puncture suppurates and sometimes leads to an opening into the pleura which might perhaps otherwise have been avoided. I should prefer in these cases repeated paracentesis, and, if that failed, I should be inclined to proceed to incision.

2. *During the Later Stage.*—If the early hours and days of pneumothorax pass over without symptoms of such urgency as to call for incision, the question of incision becomes one which may be considered at leisure, without hurry, and deliberately decided upon.

(a) If no fluid form, and the case remain one of simple pneumothorax, the air in time will be absorbed and recovery be complete. This may take some days, or even two or three weeks, but in the end the air will all be absorbed. I have never seen a case of simple pneumothorax in which the air did not in the end spontaneously disappear, and that as a rule in not many days. It is well, as a rule, not to interfere with these cases in any way, even by paracentesis; and if, with the view of accelerating recovery, paracentesis is performed, the aspirator should not be used, but the air removed without any suction. If suction be employed it should be limited to a few inches of water, say ten or twelve at the most. These cases are, however, best left alone untouched.

(b) Where serous effusion develops, so that the case is one of hydropneumothorax, it should be treated on the general lines of serous effusion; paracentesis may be performed without suction, care being taken to remove the fluid rather than the air. Cases may be cured in this way by paracentesis, but sometimes they spontaneously cure, the air being absorbed and fluid taking its place. Under these circumstances the subsequent treatment becomes that of serous effusion, and it may be that where frequent paracentesis has failed to cause the disappearance of the fluid, incision might become necessary, just as it does sometimes in cases of simple serous pleurisy.

(c) Where the effusion is purulent, that is to say, where we have a case of pyopneumothorax to deal with, and careful paracentesis fails to give relief, incision will become necessary. The chance of pus being spontaneously absorbed in pyopneumothorax is very much less than it is in empyema, and there it is as a rule small. Sooner or later incision will become necessary, and there should not be too long delay in deciding upon this operation. The indications for operation in pyopneumothorax are, I think, simply the same as those for operation in empyema, and the presence of air in these cases may, I think, be disregarded. What makes the prognosis in these cases less favourable than in ordinary cases of empyema is that in so many cases of pneumothorax tuberculous lesions in the lung are present, while they may be absent in empyema.

The indications for incision in cases of empyema are so well known that it is not necessary to refer to them here. The object of this paper is to show that there are no special reasons why the same considerations should not apply to pneumothorax as to empyema; and I do not hesitate to say that if they are—that is, if in the future cases of pneumothorax are treated on the same general principles as govern our practice in empyema, the results will be much more favourable than they have been hitherto.