The application failed to give relief, and when he presented himself at the hospital I advised him to have the ulnar artery tied.

On August 27th he was chloroformed, and the left ulnar ligatured above the wrist. Immediately after this the "beating" in the sac ceased. The hand and forearm were bandaged on a straight splint, and in the afternoon the young man was allowed to go home.

September 10th. The patient was seen to-day by me, when I found the wound above the wrist had healed. There was felt under the skin a hardish nodule, presumably the clotted contents of the aneurysm. The swelling had gone down, and all discomfort in the part had ceased.

I am quite satisfied that extirpation of the aneurysm as performed by Griffiths is the better operation. Yet, with care, I believe W. G.'s hand will give him no further trouble. The great advantage of dissecting out the sac lies in the fact that the future of such cases need give one no anxiety.

The wonder is that palmar aneurysms are so uncommon, because every engine driver and fireman uses the flat of his hand to start some stiff handle or lever of his locomotive. The blow often does produce a "dirling" sensation in the fingers supplied by the ulnar nerve, but, luckily, rupture of the intima coat of the artery does not often occur.

In conclusion, I may say I had it in my mind to tie the superficial volar along with the ulnar, but when I considered that the superficial arch was alone involved, and that the sac would be filled with firm clot ere the volar had had time to widen, I trusted to the above operation to effect a cure. I think it has succeeded, for W. G.'s hand will be fit for work very soon.

A CASE OF CERVICAL RIBS.

(With a Special Skiagraphic Illustration.)

By HERBERT ALDERSON, M.B., M.R.C.S.

CERVICAL ribs being rarely met with in the anatomical subject, and less often in the living body, a brief account of the following case with the radiograph here reproduced will, I believe, be of considerable interest.

A young lady, aged 21, consulted me this summer concerning a swelling in the right side of her neck. She complained of a little pain over the swelling, and was anxious to know if the protuberance would grow larger and thus cause a disfigurement; beyond this slight pain, which was not constant, and anxiety as to an increase of size, the swelling caused no inconvenience whatever.

The protuberance was believed to have been present some time, but it was only the last few weeks or so that the patient had taken any notice of it. The patient is a well built girl, accustomed to golfing and other athletic exercises, which are in no way interfered with by her extra rib.

On examination I perceived a slight swelling in the right anterior triangle of the neck, $2\frac{1}{2}$ inches above the clavicle, contiguous to the anterior border of the sterno-mastoid muscle. The protuberance was felt to be very hard, obviously bony, immovable, quite superficial, and over it the skin moved freely; there was no tenderness, a small artery was noticed coursing over the swelling. It appeared fusiform in shape, about the size of a brazil nut minus the shell, traced upwards or downwards it was lost in the deeper structures of the neck.

The left side of the neck appeared in every way normal and nothing could be determined by examination of the neck posteriorly.

The right subclavian artery appears to differ in no way from its fellow on the left; the pulsation of both radials is of equal volume.

The slight pain was easily accounted for by an irritation of the skin over the bony protuberance, caused by friction from a starched linen collar pretty constantly worn.

Supernumerary cervical ribs are frequently classified according to whether they possess a shaft, or merely a head, neck, and tubercle without a shaft. The radiograph reproduced on another page was taken for me by Dr. F. H. Low, and demonstrates the existence of a rib of each class, that on the left side being of the rudimentary class, having only head, neck, and

tubercle, articulating with the transverse process of the seventh cervical, but no shaft. Articulating with the right transverse process of the seventh cervical vertebra, a particularly good example of the full-grown supernumerary rib can be seen; the shaft will be noticed to pass under the clavicle to the first intercostal space, and probably joins the second rib by a cartilaginous union, an arrangement which I have not previously seen recorded. The radiograph also demonstrates a break in the continuity of the shaft of the cervical rib, with apparently the formation of a false joint, and it is this point which corresponds with the protuberance in the neck.

The patient gives a history of sundry mishaps when quite young, including a broken thigh and collar bone, which might also include a fractured supernumerary rib, or the rib and the collar bone might be synonymous.

The natural obliquity of the ribs is also well shown, the sterno-clavicular articulation and the first rib anteriorly being on a level with the fourth rib posteriorly.

The literature of supernumerary ribs, although not extensive, spreads over a considerable period of time. I find that in the *Edinburgh Medical and Surgical Journal*, vol. 40, Dr. Dymock records a case, with a drawing of supernumerary cervical ribs, met with accidentally in an anatomical subject during the year 1830. Dr. Dymock mentions that Galen, Fallopius, Winslow and others have seen such cases; but have not recorded details, an example not followed by M. Hunauld, who published an excellent paper on the subject in the *Mémoires de l'Académie Royale des Sciences* in 1740.

Professor Gruber, who collected all known cases from 1740 to the year 1809, found 76 examples of cervical ribs occurring in 45 individuals, since which time a few cases at considerable intervals have been recorded. In the *Edinburgh Medical and Surgical Journal* for 1892 a case is recorded where removal was necessary to relieve pain in the neck and arm on movement, and formication and numbness in the hand. The rib was successfully excised by Dr. George Fischer (Hanover) with complete relief to symptoms.

In the *Boston Medical and Surgical Journal* for 1896, Dr. J. Collins Warren records a case which gave rise to symptoms, pain about the middle of the clavicle passing down the arm, completely relieved by disarticulation, and removal of the rib piecemeal nearly up to its point of origin from the seventh cervical vertebra.

Supernumerary cervical ribs are probably more numerous than the literature of the subject would lead one to suppose. Many cases without doubt pass unnoticed, giving rise to no physical sign nor to any symptom, and insufficient dissection may in some cases cause them to remain unnoticed in the anatomical subject.

THE PATHOLOGY OF GENU VALGUM.

(With a Special Skiagraphic Illustration.)

By J. B. FOOTNER, F.R.C.S.,

Honorary Surgeon, Tunbridge Wells General Hospital.

IN the *BRITISH MEDICAL JOURNAL* of May 29th, 1897, there is a short but interesting communication from Mr. Charles A. Morton on the pathology of genu valgum. He there states that a skiagraph of a case of this deformity failed to show elongation of the internal condyle of the femur, but demonstrated, on the other hand, the presence of a very marked curvature in the shafts of the tibia and fibula.

The skiagraphs reproduced on another page of a severe case of genu valgum in a child aged 3 years tend to confirm his statement. It will be seen that there is some curving of the tibia and fibula, but that there is also considerable elongation of the inner tuberosity of the former bone.

One skiagraph (Fig. 1) shows the condition before operation. The other (Fig. 2) shows that the elongation of the inner tuberosity of the tibia has almost completely disappeared after operation.

I preferred division of the femur rather than the tibia, because the former is in my opinion a much more simple operation, and no important blood vessels or nerves are likely to be interfered with. In dividing the tibia, possibly some injury might be done to the anterior or posterior tibial vessels or nerves. It might also be more difficult to maintain the short

Skiagram showing cervical ribs

TO ILLUSTRATE MR. ALDERSON'S PAPER (page 1638).