PNEUMONIA: A MULTIPLE INFECTION.*

By J. W. MOORE, M.D., M.CH., B.A. UNIV. DUBL., F.R.C.P.I.,

Diplomate in State Medicine and Ex-Scholar Trin. Coll. Dubl.; Senior Physician to the Meath Hospital and Co. Dublin Infirmary; Professor of Practice of Medicine, Royal College of Surgeons in Ireland.

In a suggestive paper on Varying Infection in Pneumonia, which was published in the *New York Medical Journal*, October 9th, 1897, the author, Dr. W. H. Thompson, of the Bellevue Hospital, writes thus: "Of late years lobar pneumonia often fails to follow the definite course commonly ascribed to it. While its onset remains much the same in its suddenness, and in the rapid development of its acute symptoms, yet for some time I have declined to fix the probable date of the crisis or the duration of its subsequent stages. It may be that the advent in 1890-91 of the severest and most prolonged visitation of epidemic influenza recorded in history may have something to do with this change by contributing the influence of a mixed infection, but whatever be the cause, there is little doubt that acute lobar pneumonia now often departs more widely from its former characteristic course. In the histories of a series of 11 cases occurring consecutively in my winter service in Bellevue Hospital, in only 3 of them could it be said that they conformed to the old-fashioned type, with a definite crisis and a progressive change for the better afterwards, while in 3 a partial crisis only occurred, and in 5 none at all. In 8 out of the 11 the convalescence was very tedious, and marked by a variety of constitutional symptoms in which often the essentially toxic nature of the disease was strikingly indicated. Nothing could better illustrate than they did what a gain it was to modern pathology when lobar pneumonia was finally recognised as more an infection than an inflammation, and that its danger is due rather to systemic poisoning than to pulmonary damage. That infections by micro-organisms, however, should vary in their developments from time to time is what we should expect."

It will be observed from a close reading of the foregoing paragraph that Dr. Thompson inclines to the view that the varying phenomena presented by pneumonia in different cases probably depend upon a varying virulence of its supposed specific micro-organisms, the affection being regarded as an essential disease analogous to diphtheria, enteric fever, small-pox, or any other infective malady.

The micro-organisms in question are two: namely, (1) the micrococcus of sputum septicæmia (Fraenkel), micrococcus pneumoniæ crouposæ (Sternberg) or diplococcus pneumoniæ (Weichselbaum), and (2) the pneumococcus (Friedlaender) or bacillus pneumoniæ (Flügge). The former is now generally recognised as the usual agent in the production of acute croupous pneumonia, or, as I much prefer to call it, pneumonic fever. It was discovered by Dr. George M. Sternberg, now Surgeon-General of the United States Army, in September, 1880, in the blood of rabbits inoculated subcutaneously with his own saliva. Talamon, in 1883, demonstrated the presence of this micrococcus in pneumonic sputum, while Sternberg himself, in 1885, identified it with the micrococcus in the rusty sputum of pneumonia by comparative inoculation and culture experiments.¹ In 1886 Weichselbaum published the results of his extended researches relating to the presence of this micrococcus pretty constantly in the fibrinous exudation of croupous pneumonia. He obtained it in 94 out of 129 cases examined (54 times in cultures). To this observer we owe the name of diplococcus pneumoniæ, which serves to remind us that, as observed in the blood of inoculated animals, it is usually in pairs, consisting of oval or lance-oval elements (cocci), which are surrounded by a transparent capsule.

The second pneumonic micro-organism was obtained in 1883 by Friedlaender and Frobenius in pure cultures from the exudate into the pulmonary alveoli in cases of croupous pneumonia. Subsequent researches show that this microbe, in shape a short rod with rounded ends—hence called bacillus pneumoniæ by Flügge—is present only in a small proportion of the cases—9 times in 129 cases examined by

Weichselbaum, 3 times in 70 examined by Wolf, who pursued his studies in Weichselbaum's laboratory at Vienna.

Emmerich has demonstrated the presence of Friedlaender's pneumococcus in the soil of a room in which there were many pneumonia patients. The diagnosis was rendered certain by inhalation experiments with cultivations on 18 mice, of which 8 died of pneumonia. "Hence," says Flügge, "the soil seems to be one of the places where the pneumonia bacilli can be preserved, and whence, in suitable cases, they may pass into human beings."²

Flügge also says: "Friedlaender's bacilli are, without doubt, not the only cause of the pneumonic process. We are already acquainted with pneumonias which are caused by aspergillus and actinomyces; it is *a priori* not improbable that also among bacteria there are several other species which can set up pneumonia." Now this is the very point I wish to enlarge upon in the present communication.

The subject may be considered from both an etiological—or, we might say, a bacteriological—and a clinical standpoint. Naturally, it is chiefly from the latter point of view that I have had opportunities of regarding the question. It is impossible, however, to ignore the bacteriological aspects of the case.

In support of Flügge's statement just quoted I propose to adduce evidence that the micro-organisms peculiar to erysipelas, to influenza, to tuberculosis, and to enteric fever may one and all give rise to a specific pneumonia or pneumonic fever. So also may Loeffler's diphtheria bacillus and the bacillus of malignant anthrax, as well as other pathogenic bacteria.

I.—ERYSIPELAS.

In the form of this disease, which has been called "erratic" or "vagrant erysipelas" (erysipelas migrans)—the *erysipelas ambulans* of French writers—the attack may be protracted for one or two months. In such cases, not only every part of the surface of the body, but the whole tract of mucous membranes, and even the lungs and pleura, may in turn become affected. Dr. Peter,³ of Paris, has drawn attention to the spread of erysipelatos inflammation from the pharynx to the respiratory passages, causing in sequence bronchitis, bronchiolitis (capillary bronchitis), and pneumonia.

In a case observed by me at Cork Street Fever Hospital many years ago, the converse of this happened. A man was admitted suffering from pneumonia migrans. After some days a blush of erysipelas showed over one shoulder, and spread thence down the back, with the interesting result that simultaneously the pneumonic symptoms subsided. So great was the impression made upon me by this case that ever since I have recognised the propriety of looking upon erysipelas pulmonum as a distinct species of the great genus pneumonia.

The teaching of Levy,⁴ of Strassburg, that streptococcus pyogenes is an exciter at once of suppuration and of erysipelas is now generally accepted. This pyogenic bacterium was obtained by Fehleisen from the skin involved in cases of erysipelas in 1883, and by Rosenbach and Passet from the pus of acute abscesses within a year or two afterwards. Sternberg gives the following synonyms for streptococcus pyogenes: Micrococcus of erysipelas (Fehleisen), streptococcus erysipelatosus; streptococcus of pus, streptococcus longus (von Lingelsheim).

If, then, we admit the identity of the pus-producing streptococcus with that of erysipelas, we at once obtain a key to the occurrence of an acute pneumonia in erysipelas; for this very bacterium—the streptococcus pyogenes—plays a part that is second to none in the production of influenza pneumonia, to which I will now direct your attention.

II.—INFLUENZA.

In the great epidemic of 1889-90 in Dublin, it was my lot to see fatal cases of influenzal bronchitis, pneumonia, pleuritis, and heart failure. In a paper on the epidemic, read before this Section on February 28th, 1890, I wrote as follows: "The pneumonia (of influenza), while producing the ordinary physical signs of acute croupous pneumonia, is often latent in its course, or accompanied by a profuse muco-purulent expectoration, with scarcely any rusty sputa. The ebbing of the strength in some of these cases in elderly people is something awful—it is often absolutely beyond control." The fact

* Read before the Medical Section of the Royal Academy of Medicine Ireland, on Friday, November 19th, 1897.

is that influenza, infrequently directly fatal, causes an indirect loss of life which is appalling, chiefly through the complications affecting the respiratory organs and the heart, which have just been mentioned.

It will be remembered that, after the great pandemic of influenza in 1889-90, German medical literature in particular was flooded with writings upon the clinical, pathological, and bacteriological aspects of the malady. In the *Dublin Journal of Medical Science* for May, 1890,⁵ and August, 1890,⁶ will be found reports on the bacteriology and pathological relations of influenza, which I prepared from current German medical literature. From these reports I cull the following facts:

Leyden, in a communication to the Medical Society of Berlin,⁷ states that the pneumonias observed by him showed a peculiar course; severe pain in the side and dyspnoea were rarely noticed; the local process was not altogether typical; frequently it was necessary to watch for three or four days before any evidence of a localisation of the disease was forthcoming. Then a crepitating *râle* was heard over a wide area, and this perhaps the very next day would have disappeared to show itself in some other situation. Not very often a fine hepatisation occurred with clearly mapped-out dulness. Again, the typical sputum of pneumonia was often wanting. Bacteriological investigations revealed the presence of three kinds of microbes:

1. Diplococci, which represented the well-known pneumonia diplococci of Fraenkel.
2. Streptococci.
3. Staphylococci.

Leyden adopted the view that the forms of pneumonia are different: typical genuine pneumonias with deviating course; mixed forms, especially in those combined with pleural effusion; lastly, simple streptococci pneumonias.

Ribbert,⁸ discussing the possibility of a causal significance of the streptococcus pyogenes in relation to the phenomena of influenza, alludes especially to the inflammations of the lungs, whose peculiar erysipelas-like spread, on which Finkler lays so much stress,⁹ and whose anatomical relations admit of being referred back to the influence of the streptococcus. Ribbert points out¹⁰ that, in contrast to ordinary croupous pneumonia, the cut surface of the hepatised lower lobe in 3 cases presented an almost smooth appearance, the exudation was soft, very rich in cells, and poor in fibrin (hypinosis). Cultivation experiments with the tracheal mucus, the lung tissue, the spleen, and the kidneys, furnished in 5 out of 8 cases the streptococcus pyogenes, or else the streptococcus erysipelatosus (which has been shown to be identical with the former), the presence of which microbe could be demonstrated in the sputum also of the influenza patients. His investigations on the whole yield the result that in all cases in which micro-organisms were at all capable of demonstration the streptococcus pyogenes was found. Only once was there in addition a coccus, which had a great resemblance to the diplococcus pneumoniae, and probably represented a modification of the same.

Finkler¹¹ observed 45 cases of influenzal pneumonia, of which only 2 came under the description of typical lobar pneumonia, while the other 43 were regarded as cases of the disease which he has often described as "streptococcus pneumonia." Seven of his patients died, *post-mortem* examinations being made in 3 instances. He regarded the pathological condition as a preponderating cellular inflammation with participation of the interstitial tissue. The cellular nature of the inflammation, together with the pronounced tendency it exhibits to develop by spreading indefinitely, in Finkler's opinion justified him in describing this disease as an erysipelas of the lung. He points out that the resemblance of this form of pneumonia to erysipelas consists not alone in the anatomical characters of the inflammatory process, but also in the fact that both diseases depend on the presence of streptococci. Finkler looks upon this streptococcus pneumonia as a localisation of the exciting cause of influenza in the lungs. As to this last point, I should be more inclined to agree with Leyden¹² and Levy,¹³ that the question is much more one of a secondary infection, for which the influenza merely laid the foundation. Certainly the discovery by Pfeiffer, in 1892, of the bacillus influenzae in the purulent bronchial secretion, and by Canon in the blood of patients suffering from epidemic influenza, must be regarded as con-

clusive proof of the existence of a specific primary infection to which all other infections are accidental and secondary.

III.—TUBERCULOSIS.

It is not my intention here to allude to acute tuberculous fever (in which the lungs may escape unscathed), on the one hand, or on the other to the local peripneumonic processes which accompany sporadic depositions of tubercle in the lungs in ordinary catarrhal phthisis. Nor will I refer to those cases in which in the wake of an acute primary croupous pneumonia the wounded lung falls a ready prey to a secondary infection by the bacillus tuberculosis, when this micro-organism finds a fertile soil in the caseating exudation of an unresolved pneumonia. These several conditions are all beside the present question.

My concern is with acute phthisis, or scrofulous pneumonia, and the so-called acute tuberculo-pneumonic phthisis. Dr. C. Theodore Williams thus describes acute phthisis¹⁴: "The patient, generally young, who may have had cough previously, is attacked with sharp pain in one side of the chest, quick pulse, high temperature, the skin being quite burning to the ear of the auscultator, alternating with night chills and sweats. The general appearance betokens pneumonia, but the crepitation commences at the apices, extending to the whole lungs, and is not so fine and even as in pneumonia. The cough increases; the expectoration becomes opaque and purulent, containing quantities of lung tissue and swarms of tubercle bacilli; and the temperature assumes the intermittent type. The physical signs show at first gradual consolidation of both lungs, but later on indicate that excavation has taken place; and continues, the patient rapidly wasting and dying in a few weeks." In this disease the inflammatory nature of the lesions in the lung or lungs, and the rarity of miliary tubercle, are among its characteristics.

Acute tuberculo-pneumonic phthisis likewise presents consolidations in the lungs of a pneumonic origin, but tuberculation, as well as pneumonia, exists.

In both these varieties of "consumption" we have examples of true pneumonia resulting from an infection by the bacillus tuberculosis of Koch.

IV.—ENTERIC FEVER.

It is well known that pneumonia is more commonly observed as a complication in enteric fever than in typhus. Murchison noted it in 13 out of 100 cases, and Austin Flint (according to Bartlett¹⁵) in 12 out of 73 cases. It commonly occurs in the third or fourth week, but may usher in the disease. In this latter case its presence is probably an indication that the enteric fever poison has entered the system through the lungs. It is most commonly a lobular pneumonia, but occasionally it occurs under the form of ordinary croupous pneumonia.

It is indeed true that Eberth points out¹⁶ that anatomical investigations had up to 1881 afforded no evidence of the admission of the bacillus typhosus through the lungs. With this Gaffky¹⁷ does not agree, for he considers it highly probable (or, at least, the possibility cannot be contested) that the lungs may occasionally represent the seat of invasion. Eberth himself quotes a case observed by W. Meyer,¹⁸ of Berlin, in which death ensued on the second day of illness. In this case there were found at the necropsy hyperæmia of the lungs, spleen, and kidneys; in the lower portion of the ileum marked swelling of the solitary follicles and Peyer's patches. Microscopical examination revealed a very exceptionally large deposit of Eberth's bacilli in the cells of the submucosa and in the intermediate muscular layers of the intestine. Apparently they were not found in the lungs, notwithstanding their hyperæmic condition.

That, in infective diseases in general, infection may occur through the mucous membranes of the respiratory tract has been demonstrated, according to Sternberg,¹⁹ by several bacteriologists—especially by Buchner, who caused mice and guinea-pigs to breathe an atmosphere containing in suspension a powder consisting of dried anthrax spores mixed with lycopodium powder, or pulverised charcoal. In a series of 66 experiments, 50 animals died of anthrax, 9 of pneumonia, and 7 survived. Microscopical examination of sections and culture experiments showed that the lungs were extensively invaded. It may be objected that these results do not bear

on infection by the bacillus typhosus, which is believed not to assume the spore form. Positive results were, however, also obtained by Buchner with cultures of the anthrax bacillus not containing spores, which the animals were made to inhale in the form of spray. But in this case a considerable quantity was required, and a sero-fibrinous pneumonia was usually produced, as well as a general infection. "That man may be infected with anthrax by way of the respiratory organs," writes Sternberg,²⁰ "seems to be well established. In England the disease known as 'wool sorters' disease' results from infection in this way among workmen engaged in sorting wool, which is liable to contain the spores of the anthrax bacillus when obtained from the skin of an animal which has fallen a victim to this disease. That infection occurs through the lungs is shown by the fact that these organs are first involved, the disease being, in fact, a pulmonic anthrax."

Even if we take it as not yet proved that infection in enteric fever may occur by way of the lungs, there is no doubt that a close correlation exists between this disease and that variety of acute pneumonia, or pneumonic fever, to which the term "pythogenic pneumonia" has been commonly applied since 1875, when Dr. Grimshaw and I read a paper on the subject before the Medical Society of the College of Physicians of Ireland.²¹ Towards the end of October, 1882, the following remarkable outbreak of disease came under my notice.

On the 12th of that month a lad, aged 13, was admitted into Cork Street Fever Hospital from 6, Malpas Street, Dublin, suffering from croupous pneumonia. Malpas Street is very unhealthy; the houses are old and dirty, ill-drained and dilapidated. The street runs down to the bottom of a valley, through which a small tributary of the Poddle river flows sluggishly. The district is a prolific hotbed of disease. On October 31st the boy's father (J. C.), a boatman, aged 36, came in with the same disease. On the 20th of the same month two girls, both aged 14, were admitted to the Meath Hospital in enteric fever—one from 11, Malpas Street, and the other from No. 13. On November 27th a girl, aged 20, was admitted to Cork Street Hospital in enteric fever from 7, Malpas Street, next door to the house from which the two cases of pneumonia had come a few weeks previously. On December 12th, J. C., aged 36, was again admitted to the Meath Hospital from 6, Malpas Street, with "renal dropsy." It was he who suffered from pythogenic (?) pneumonia in the previous October as narrated above. Another coincidence occurred in March, 1883. On the 18th of that month W. N., aged 19, came into Cork Street Hospital from 6, Malpas Street, in an attack of "febricula," and the following day A. L., aged 27, was admitted from the same house with left basic croupous pneumonia.

A very similar instance of the correlation existing between enteric fever and pneumonia came under my observation in the autumn of 1881. Four cases of illness occurred in a training college in Dublin within a few weeks. Two of the four patients suffered from true enteric fever; a third, from an attack of acute gastro-intestinal catarrh, or, as some may think, from an abortive enteric fever; and a fourth from acute pneumonia, reminding one of Laennec's "epidemic pneumonia," which in recent times has received the names of "sewer-gas pneumonia" and "pythogenic pneumonia." The drinking water was proved by Sir Charles A. Cameron to be the source of the sickness in all four cases.²²

In November, 1891, there came under my care in the Meath Hospital a young woman with characteristic typhoid stools, and whose urine gave a striking reaction with Ehrlich's diazo-test. Her illness, however, had commenced with right apex pneumonia, with rapid breathing, cough, glutinous expectoration (not indeed deeply coloured when the patient was first seen by me), dulness on percussion, and, finally, the most typical crepitus redux.

One of the reasons which weighed with me in undertaking to make the present communication to the Royal Academy of Medicine was the remarkable tendency to a pneumonic element in enteric fever which has shown itself in Dublin during the present season. It will be remembered that the end of August and beginning of September proved both wet and cold, while the air temperature continued below the average through the greater part of September.

These atmospheric conditions were doubtless the prime reason why the epidemic enteric fever assumed the so-called thoracic form. I venture to submit brief notes of two cases in illustration of this statement.

CASE I.—R. A., aged 40, a bootmaker, was admitted to the epidemic wing of the Meath Hospital, from New Street, on September 17th, 1897, on the eighth day of his illness. His evening temperature was 104.3°, pulse 100, respirations 32-36. When I saw him next morning I found him suffering from acute catarrhal laryngitis; the base of the left lung was quite dull, with the other signs of consolidation. He was expectorating a viscid sputum from which the rusty colour was already quickly disappearing. There was active diarrhoea, the daily number of motions varying from three to seven. Forty-eight hours after admission, profuse intestinal hemorrhage occurred, after which the attack ran the usual course of a typhoid fever, with laryngeal and pulmonary trouble ("laryngotyphus" and "pneumotyphus"). Laryngeal examination by Dr. Richard Lane Joynt revealed a severe catarrh. The chart is a characteristic one of enteric fever, but the rapidity of the respirations sufficiently indicates

the severity of the pulmonary lesion. In convalescence the patient suffered from boils as a secondary infection.

CASE II.—J. B., aged 22, a domestic servant, was admitted to the epidemic wing of the Meath Hospital, from Ranelagh, on September 15th, 1897. She was then a week ill. She was sent in as suffering from acute pneumonia, and the physical signs of left basic lobar pneumonia were present. She breathed from 40 to 44 times a minute, while her pulse at first did not exceed 100, although some days later it rose to 120. The temperature was only 101° on the evening of her admission, but gradually rose until the evening of the twelfth day of her illness, when it reached 104.3°. For several days the patient's condition was extremely unsatisfactory, and in appearance and physical signs the case strongly resembled acute phthisis, or "scrofulous pneumonia." However, she gradually emerged from the fever, the evening spiking of the temperature subsiding after the twenty-fourth day. She left hospital for the Convalescent Home at Bray, on October 22nd, the forty-fifth day from the commencement of her attack. At that time the state of her left lung remained far from satisfactory, though it was fast improving. There was still dulness on percussion over the base, the breath and voice sounds were feeble, and some effusion still existed in the pleura.

These cases are types of the form which enteric fever assumed in the unseasonably cold after-summer of the present year. That the lung attacks were the outcome—direct or indirect—of a specific poisoning by Eberth's bacillus can scarcely be doubted. This poisoning may, it is true, have been secondary to a localisation in the intestine, although the fact that in each case the lung trouble occurred at the outset of the fever points with much force to a primary localisation in the lungs. However that may be, the presence of an acute pneumonia in each case must be conceded.

Drs. Muir and Ritchie, writing in 1897,²³ say that most observers will agree with Gaffky in attributing any failure to find typhoid bacilli in the organs of a typhoid patient to the difficulties of the search.

These writers further state that in the lungs there may be patches of congestion and of acute bronchopneumonia. In these, typhoid bacilli may sometimes be observed, but evidence of a toxic action depressing the powers of resistance of the lung tissue is found in the fact that pneumococcus is frequently found in such complications of enteric fever.²⁴ As to this, I repeat that the very early appearance of pulmonary trouble in a certain proportion of cases which ultimately prove to be undoubtedly enteric fever is altogether in favour of a primary infection with the bacillus typhosus by way of the lungs. A pneumonia brought about through lessened resistance to the specific micro-organisms of this disease caused by the toxic action of the typhoid bacillus on the system in general, including the organs of respiration, would be much more likely to develop in a more advanced stage of the fever.

In conclusion, I venture to submit that there is clinical evidence to show that a true pneumonitis may occur in any one of the four diseases with which this communication deals—that is to say, erysipelas, influenza, tuberculosis, and enteric fever. Further, it is reasonable to suppose that in each case the pneumonitis is directly due to a localisation of the specific poison of the disease in the lung, whether that poison be a micro-organism itself, or a toxin derived therefrom. Indeed, in respect of three out of the four diseases named, the evidence, from a bacteriological standpoint, in favour of such a view is incontrovertible.

As regards enteric fever, the influence of season and weather in determining pneumonic trouble is, no doubt, considerable; but it cannot be accepted as paramount or exclusive. And, if it is objected that the bacilli typhi abdominalis of Eberth have not as yet been often found in the lungs or sputum of enteric fever patients, I am justified in attributing this to the comparative infrequency with which, so far, search has been made for these bacteria in the pulmonary organs. Here, in any event, an almost untrodden path of investigation lies open to the adventurous footsteps and the keen perception of our Irish pathologists and students of bacteriology.

REFERENCES.

- ¹ *American Journal of the Medical Sciences*, July 1st, 1885.
- ² *Micro-organisms*, by Dr. C. Flüge. Translated by W. Watson Cheyne. London: The New Sydenham Society, 1890, p. 259.
- ³ *Dictionnaire Encyclopédique des Sciences Médicales*, Tome IV, p. 720, Art. "Angines."
- ⁴ Ueber die Mikroorganismen der Eiterung, *Archiv. für experiment. Path. und Pharm.* xxix, p. 135.
- ⁵ Vol. lxxxix, No. 221, third series.
- ⁶ Vol. xc, No. 224, third series.
- ⁷ *Berl. klin. Woch.* 1890, No. 10.
- ⁸ *Deut. med. Woch.* 1890, No. 15.
- ⁹ *Ibid.*, 1890, No. 5.
- ¹⁰ *Ibid.*, 1890, No. 4.
- ¹¹ *Ibid.*, 1890, No. 5.
- ¹² *Loc. cit.*
- ¹³ *Berl. klin. Woch.*, 1890, No. 7.
- ¹⁴ Quain's Dictionary of Medicine, vol. ii, Art. "Phthisis," new edition, 1894.
- ¹⁵ *The Fevers of the United States*, fourth edition, Philadelphia, 1856.
- ¹⁶ *Virchow's Archiv.* lxxxi, 486, 1881.
- ¹⁷ On

the *Ætiology of Enteric Fever*, *Mittheilungen aus dem Gesundheitsamte*, B. ii, Berlin, 1884. ¹⁸ *Untersuchungen über den Bacillus des Abdominaltyphus*, *Inaugural Dissertation*, Berlin, 1881. ¹⁹ *Loc. cit.*, p. 230. ²⁰ *Loc. cit.*, p. 231. ²¹ *Dublin Journal of Medical Science*, vol. ix, No. 41, third series, May, 1875, p. 399. ²² *Ibid.*, vol. lxxiii, No. 122, third series, February, 1882, p. 131. ²³ *Manual of Bacteriology*, Edinburgh and London, Young J. Pentland, 1897, pp. 298. ²⁴ *Loc. cit.*, p. 312.

A FATAL CASE OF THYROIDECTOMY.

By F. T. PAUL, F.R.C.S.,

Surgeon to the Royal Infirmary, Liverpool.

A CONSIDERABLE interest attaches to a certain remarkable train of symptoms liable to supervene upon operations on the thyroid gland. With me these have occurred in cases in which the operation had been completed in the most satisfactory manner, and in which the wounds have been perfectly healthy, so there can be no doubt that the symptoms were due to a disturbance of the gland, and were not incidental to the wound. When the first case occurred, which was the one ending fatally, I had not seen anything of the kind published, and formed my own conclusions independently. Since then I have had other cases, and some have been published by Mr. Victor Horsley and Dr. Rodocanachi, from all of which it appears clear that the symptoms are due to the absorption of thyroid secretion through the wound.

The case is shortly as follows: A.M., aged 15, a slim, tall girl, had a rather large parenchymatous goitre. She was a native of Liverpool, but her father was a south-countryman, and an aunt has a goitre. The tumour was soft, but not cystic, and did not pulsate. It involved both lobes about equally. The heart was excitable, the impulse diffused, and there was a soft systolic *bruit* at the apex. The pulse was 104. There was dyspnoea on exertion. She was said to be a good useful girl at home, and, though of the thin nervous type, appeared to be quite strong enough to bear a serious operation. On December 3rd I removed the left lobe and isthmus of the gland. The operation was perfectly straightforward, and the patient left the theatre in excellent condition. All details as to dressings, drainage, etc., were carried out in the usual manner. During the night she became restless and uneasy, an irritable cough set in with a mucous rattle in the throat. The temperature was 100°, the pulse 130, and the respirations 26 and laboured. On the morning of December 4th the same condition was present; the wound was dressed and appeared perfectly healthy. In the afternoon the temperature rose to 103°, and fearing that the ether inhalation had set up bronchitis, steam and eucalyptus vapour were used, with stimulants and extra nourishment. Subsequently the restlessness, anxiety, distressed breathing, and frequency of the pulse increased, and the temperature was generally between 101° and 102° F. There was no further evidence of bronchitis. She took nourishment and stimulants well, but notwithstanding the most devoted care on the part of my house-surgeon, Dr. Badger, and the sister of the ward, the Graves's disease-like symptoms increased in severity, and the pulse became uncountable, and she died on December 6th, just two and a-half days after the operation. Small doses of morphine were given a few times, and on each occasion were followed by a little sleep. Probably, as observed by Dr. Rodocanachi, it would have been better to have used this drug more freely. A necropsy was made. In regard to the wound, there was a want of healing action, and the fluid contained in it was of a very watery character, but there was nothing suggestive of septic changes. The divided surface of the thyroid looked quite fresh, as though repair had not yet commenced. The bases of the lungs were congested, and there was a little tenacious mucus in some of the tubes, but no general bronchitis. The heart and all the other internal organs were healthy, except the liver, which was very fatty and perfectly nodular throughout from idiopathic cirrhosis. At first I thought this remarkable liver might be sufficient to account for the death of the patient, but soon dismissed the idea, for on reviewing the whole circumstances I could come to no other conclusion than that the symptoms were due to the absorption of an excessive amount of thyroid secretion, but how this came about, and what I could do to prevent it, did not at once occur to me.

A fortnight later we had another object lesson. A pale,

sallow, nervous woman, aged 43, with rather chronic Graves's disease, was sent to me for operation by Dr. Abram, as she had undergone no improvement after a course of medical treatment. The thyroid was only moderately enlarged, and exophthalmos, though present, was not marked; otherwise the symptoms were usual.

On January 18th, the patient being in fairly good condition, I removed the right lobe of the gland without difficulty, and with practically no loss of blood; in fact, the operation could not apparently have been more satisfactory. For several hours the patient was very comfortable; but during the night she became restless with quickened respiration. The temperature was 101°, the pulse 128. Next morning she was a little better, and on changing the dressings the wound looked well. Later in the day the restlessness, anxiety, and irritable cough increased, and the pulse and respiration became more frequent. The ward sister was already anticipating a repetition of the last case, and it seemed to me only too probable that her fears were well grounded. I again removed the dressings. There was now no blood stain, but the wound was discharging freely a clear watery fluid, and on cutting the stitches it was seen to be filled with the same. I imagined that this fluid, or at any rate a considerable part of it, was derived from the thyroid, and wanted Dr. Abram to test it for me; but it appears that there is no chemical means of recognising thyroid secretion. However, acting on the belief that the patient was suffering from a sort of traumatic thyroidism, I left the wound open, but packed it to the bottom with dry salicylic wool. The opening up of the wound was followed by a marked improvement, but only for a time. During the night—the second night after the operation—she became worse than ever; the temperature was 104.8°, the pulse almost uncountable, the respirations 36. I removed the plug of wool, and found it saturated with watery discharge, replaced it with dry wool, and left instructions that it was to be changed as often as it became at all moist, which proved to be about every two hours. The following day, January 21st, she was better in every way. On January 22nd the temperature was only just above normal, and continued so until convalescence was established, but the pulse and respiration went down more gradually. We had no further anxiety about her.

These two cases were the twenty-fifth and twenty-sixth operated on for thyroid enlargements, and I have done about ten since. None of the others were at any time dangerously ill, but I can see now that a few of them suffered from the same condition in a minor degree, and at about the same time, I therefore set myself to try and discover what could be the reason that an operation which at first appeared so safe should all at once have become so uncertain. Upon looking back, this explanation occurred to me. In the earlier operations I handled the gland very little, tying the vessels on each side before dividing them, and ligaturing the isthmus before separating the lobe. The operations then were longer, and more blood was lost. Indeed, in Graves's disease the hæmorrhage was sometimes alarming, for in these cases the blood wells out of the gland as if it were a saturated sponge. Gradually I came to grasp the gland firmly in the left hand until the thyroid arteries were secured, and in this way found that all serious bleeding was readily avoided. About the same time I gave up ligaturing the isthmus, as it appeared unnecessary. Now in these two changes it was probable that the danger lay. From barely handling the gland it had come to be firmly squeezed during most of the operation, and from securely tying the isthmus it had come to be cut straight across—and it must be remembered that in enlarged thyroids the isthmus is frequently upwards of an inch in diameter. I take it that squeezing the gland may help to liberate secretion contained in the follicles, and that the same may escape into the wound from the lymphatics in the divided capsule round the severed isthmus, the lymphatics being the normal channel for absorption of the secretion. If the condition from which these patients suffered is to be regarded as thyroidism, and not, as Mr. Horsley has said, athyroidism, then every possible source of contamination of the wound with thyroid secretion should be avoided.

I cannot recommend that the safe grasp of the gland should be altogether given up; but I believe it may be rendered harm-