doubtedly some examples occur (particularly of dorsal dislocation of the index first phalanx) in which great difficulty is met with in reduction. M. Polaillon¹⁰ showed three cases, two concerning the index and one the ring finger, of dorsal dislocation which resisted all attempts at reduction under anaesthesia and remained out of place. In all such cases it can hardly be doubted that the real difficulty is in the glenoid ligament, and its division may be recommended with confidence should the ordinary manipulation fail.

Finally, it may be said that with the correct understanding of the part the sesamoid bones and the glenoid ligament play in backward dislocation of the thumb, and with a resort to the simple method of operation already described in cases otherwise irreducible, all the difficulties of this much discussed subject practically disappear.

NOTES AND REFERENCES.

¹ A mistaken attempt has been made in one of the most recent works on anatomy to do away with the glenoid ligament of this articulation, and to describe merely "sesamoid bones in place of a glenoid ligament." It is hardly necessary here to combat this view. ² *Bull. de la Soc. de Chir.*, Tome iv, 1878, p. 743. ³ *Edinburgh Medical Journal*, 1815, p. 188. ⁴ *Treves's System*, vol. i, p. 592. ⁵ *BRIT. MED. JOUR.*, August 24th, 1895. ⁶ Mr. W. H. Battle, *Lancet*, December 22nd, 1888. ⁷ *Affections Chirurgicales des Membres*, 1895, p. 451. ⁸ Mr. T. Bryant, *Manual of Surgery*, vol. ii, p. 372. ⁹ *Transactions*, vol. iv, p. 250. ¹⁰ *Loc. cit.*, p. 452.

CLINICAL REMARKS

ON

CATHETERISATION OF THE MALE URETERS UNDER ELECTRIC LIGHT.

By E. HURRY FENWICK, F.R.C.S.,

Surgeon to the London Hospital, Surgeon to St. Peter's Hospital for Urinary Disease.

THE MODE OF EXAMINATION.

SINCE the introduction of the electric cystoscope it has been foreseen that the next important step in the progress of precise urinary surgery would consist in an easy method of catheterising the male ureters. It was felt that once this manœuvre was rendered practical, much of the obscurity in surgical renal disease would disappear, many of the false impressions we still retain about the symptomatology would be effaced, and the mortality of nephrectomy would be assuredly lessened. It is generally allowed that the female ureter is easily entered and traversed by a suitable catheter guided either by the finger (Simon, Pawlik), or by the eye (Newman, Hirst, Hamill, Kelly, Goldschmidt); but the disappointment caused by the failure of Brenner's instrument in 1889 has rendered us sceptical as to the possibility of guiding an instrument into the male ureter under electric light.

This manœuvre is, however, possible, and has been now in active use for two years. Not only is it practical in most cases, but to the dexterous it is easy. It will probably be easiest to those who have studied the varying positions and aspects of the ureteral mouths under electric light, and to those who have become adepts in guiding fine bougies under electric light over the mouths of false passages in the strictured male urethra.

Fingers and eyes thus skilled can engage most ureters in a couple of minutes, provided a suitable cystoscopic instrument is employed. I wish, however, before drawing attention to the instrument, to dissent somewhat strongly from the opinion of one surgeon, who asserts it can always be performed in the consulting room under cocaine. Of course it may be done, but not, I submit, with safety. No harm can follow when there is little chance of inflammatory reaction (cystitis, pyelitis, suppression) resulting. Thus it will probably do no harm to catheterise ureters in one's room, provided the patient has not reached 45, the urine is quite clear, the kidneys are healthy, and that no tuberculosis of the urinary tract is present. But under such healthy conditions the procedure is not called for.

It will probably be found that those patients who really need ureteral exploration for obscure disease possess kidneys too sensitive to ureteral or urethral shock to permit any risk of a subsequent chill being run. The immediate and remote death-rate due to washing out the bladder in cases in which pyelitis or irritable explosive kidneys coexist is sufficiently

high to discourage anyone from washing out that viscous in one's consulting room for the first time—a requisite in catheterising ureters; in fact, to do so at all, except under the most favourable circumstances of rest and warmth.

If the ureter has to be explored, let it be done with care and caution, let the patient be in bed, and remain there quietly for a few hours after the examination.

THE INSTRUMENTS.

Several forms of catheterising male cystoscopes are on the market, but as yet they fall mainly under two patterns—the Nitze and the Casper. As will be seen from the accompanying Figs. 1 and 2, both consist in a cystoscope

Fig. 1.—The Nitze Male Catheterising Cystoscope.

Fig. 2.—The Casper Male Catheterising Cystoscope.

with its usual electric lamp L, its prism P, and its telescope. Attached to one side is a channel for the catheter c to run in. This is so placed as to allow the point of the catheter to emerge into the bladder under the eye of the operator. He is thus able to guide it into the mouth of the ureter.

It is difficult to say which instrument will find the greatest favour. Both are valuable, both need, in my opinion, improvement. For my own part, I work the Casper without difficulty, and prefer it because the point of the catheter can be so easily manipulated through a large angle by pressing on the ring D.

This instrument is introduced into the bladder, the catheter lying hidden in its channel. The mouth of the ureter selected is brought into view, and the catheter is pushed along its groove until its point is seen to emerge. A little movement of the cystoscope itself, or manipulation of the ring D, to alter the angle, or a slight rotation of the catheter, is sufficient to guide the latter into the mouth of the ureter (Figs. 3 and 4) and it can now be watched swelling up the wall of the bladder as it passes up the ureter. In a few seconds urine will drop from the end of the catheter, and may be collected in a test

tube. It is best to give the patient plenty of fluid some time before the examination in order to excite the renal function. I have for long relied upon half a bottle of Contrexéville water

Fig. 3.—The Nitze Catheter entering left Ureter.

Fig. 4.—Cystoscopic picture of a Casper catheter entering mouth of right ureter.

taken an hour before exploration. Cocaine is ample in most instances when the urethra is tolerant and capacious. If an anæsthetic is needed chloroform is perhaps more suitable, as it interferes less with the rapid secretion of the kidney than ether (?).

RAPID DETECTION OF THE SITE OF THE URETER MOUTH AFFORDED BY STAINING THE EFFLUX WITH ANILINE DRUGS.

When the bladder base is much swollen or the ureteral orifices are displaced, or the kidney is sluggish, or herniæ coexist, difficulties may arise. In detecting the mouths of the ureters and distinguishing a thin-edged patulous orifice from a minute hernia the value of a stain for the urine will be obvious. I have for some years made use of fuchsin ($\frac{1}{2}$ gr. in pill). This, if taken an hour or so before the examination, will colour the urine a delicate pink, so that after the bladder has been washed out streams of pink urine will be seen to issue from the ureters. I may be permitted to remark that fuchsin does not affect some kidneys at all, and others which are undoubtedly weak pass a good deal of the colour through. Some years ago I hoped to be able to test the efficiency of the secreting cells by means of fuchsin, but extended experience showed me the test was unreliable.

SOUNDING THE KIDNEY PELVIS AND URETER FOR STONE UNDER ELECTRIC LIGHT.

Before the male ureter could be catheterised under the light, I used to have the female ureteral catheter tipped with metal for sounding the female ureter, for the fingers can detect by these means the presence of any small calculus in the channel of the ureter. Sounding the ureter is difficult in the male because of the many bends the catheter has to make and the many corners it has to pass; the gripping and friction of these tend to obscure the sense of touch. But there is no doubt in my mind that this is a step in the future.

The microphone perhaps may render it of more value. I believe that ureteral obstructions from kinks and stones are more frequently the cause of renal pain and colic than is at present allowed.

ON THE CAUSE OF DEATH BY ELECTRIC SHOCK.

BY
THOMAS OLIVER, and ROBERT A. BOLAM,
M.A., M.D., F.R.C.P., M.D.,
Professor of Physiology, Joint Lecturer in Physiological
Chemistry and Physics,
University of Durham, College of Medicine, Newcastle-on-Tyne.

THE increasing employment of electricity within the last few years has demonstrated, from the accidents to workmen engaged in its generation and distribution, not only what a dangerous force it is, but the necessity for using all the precautions suggested by experts. In this country alone within the last few years, sixteen deaths have been caused by electricity. It has seemed to us, therefore, that the present time is opportune for reopening the subject with the view of ascertaining the cause of death in such cases.

A powerful electric current kills immediately. Two opinions are before the profession (1) that death is due to failure of the respiratory centre (d'Arsonval) (2) that it is due to sudden arrest of the heart's action. From the appearance presented by the internal organs after death some physiologists have maintained that death is due to asphyxia. We have it on the testimony of workmen employed in electric generating stations that when one of their mates has been fatally injured by coming into contact with highly charged metal, he has generally breathed a few times although to all appearance dead. This fact, therefore, would suggest that death was not due to failure of the respiratory centre. To what other cause may it be attributed? It is desirable that a definite opinion, if possible, should be arrived at as to the cause of death in electric shock, so that we may have at our command means for the resuscitation of those who have been apparently killed by such discharge. As this subject formed part of an inquiry by a Home Office Committee upon which one of us served, and for which we instituted a series of experiments upon animals, most of them under anæsthesia, with the view of determining these points, we feel that the data thus obtained have a scientific not less than a practical value, and as such we offer them for consideration.

The animals employed in this research have been dogs and rabbits. In almost every case ether was used as the sole anæsthetic. The blood pressure was recorded in the ordinary way, the left femoral artery being usually opened in the dog, the left carotid in the rabbit. Respiratory movements were registered by means of a Verdin's double tambour applied to the chest of the animal, and communicating with a recording tambour. When artificial respiration was used in the attempts at resuscitation after shock it was generally performed by the Sylvester method; occasionally by inflation through the usual tracheal cannula inserted at the outset of the experiment. An alternating current supply in the laboratory running at 100 volts and 100 cycles per second was transformed as required. Wet sponge electrodes mounted on 6 inch vulcanite handles for efficient insulation were applied to the fore and hind limbs on portions of the skin previously shaved and thoroughly moistened. The period of stimulation was shown by an ordinary time marker on a shunt circuit in which were placed two incandescent lamps in series. We would take this opportunity of cordially acknowledging our indebtedness to Mr. R. J. Patterson, Lecturer in Physics, Durham College of Science, for much help in the physical portion of the work as well as for personal aid in many of the experiments.

RESULTS.

When an animal is exposed to electric currents of sufficiently high potential death, practically speaking, is instantaneous. The animal is thrown into a condition of opisthotonos, during which breathing is suspended; the heart's beat momentarily quickened and then arrested, while with currents of somewhat higher potential still it is immediately arrested without any preliminary quickening. On shutting