THE POWERS OF THE COUNCIL.

The powers of the Council are, I consider, specifically laid down in Clause 20 of the Act of 1858, which gives to it control over professional study and examination, whilst Clauses 28, 29, and 40 lay the foundation for the exercise of those penal powers which the Council has so frequently put in force for the protection of the public and for the maintenance of professional ethics. So that not alone does the Act confer the power of interference with the internal discipline of the profession, but the procedure of the Council shows how freely that power has been exercised.

In connection with the ambiguous wording of some of the clauses of the Act of 1856, I may briefly allude to a point raised by Mr. Horsley in one of those addresses which he has delivered on the construction of the Medical Acts, addresses so creditable to him not alone for the cogency of their reasoning, but for the disinterested zeal which they display. Mr. Horsley points out that the Act specifically refers to members of the profession who are registered as "qualified practitioners," and declares that they should be distinguished from unqualified practitioners. He maintains, therefore, that the Act protects the individual, not the title he possesses. It confers upon the individual the right to practise, as well as certain other rights and privileges laid down in Clauses 31, 32, 35, 36, and 37 relating to recovery of fees, etc., immunity from serving on juries or in the militia, holding public medical appointments, and signing certificates, etc. That Mr. Horsley is correct in his reading of the meaning of the Act is made manifest by the fact that removal of the name from the *Register* does not remove the title, which the person still continues to hold until the body from which he obtained it strikes the name off its list of licentiates or graduates as case may be. Lord Justice Lopes, in his judgment given in the appeal case of Allbutt against the General Medical Council, lays down that by Section xxxiv of the Act of 1858 a "legally qualified practitioner" is to be construed to mean a person registered under the Act, which to any ordinary mind endowed with normal, not forensic, intelligence would indicate a person alone privileged by law to practise.

But whilst fully recognising that those only who are registered are entitled to practise, it must, I think, be admitted that the wording of the Act is too vague to permit of prosecution in all cases where persons practise without being registered. The Act states that "any person who shall wilfully or falsely pretend to be or take or use the title of Physician, Doctor of Medicine, Licentiate of Medicine or Surgery, Bachelor of Medicine, Surgeon, General Practitioner, or Apothecary, or any name, title, addition, or description implying that he is registered under the Act.....shall, upon a summary conviction for any such offence, pay a sum not exceeding twenty pounds." Take, however, the case of a person who was a *bona-fide* "Doctor" or "Surgeon," and who practised as such without being registered, could it be held that such a person "wilfully and falsely" assumed the title which he had legitimately obtained? Furthermore, it would be contended, in the absence of any direction making registration obligatory, that he was sufficiently penalised by deprivation of the rights and privileges laid down in the clauses already referred to. The extraordinary contention of Lord Justice Lopes in the case cited, that the General Medical Council should publish the grounds upon which a name was erased from the *Register* to enable former patients to decide whether or not they might employ the practitioner who had ceased to be legally qualified, seems not alone to be at variance with the spirit and meaning of the Act, but contradictions of the construction put upon Section xxxiv by the learned judge himself.

AMENDMENTS DESIRABLE IN THE MEDICAL ACTS.

If I might venture to enumerate the amendments which it might be desirable to make in the Medical Acts, I would tabulate them as follows: 1. No person should be allowed to practise medicine, surgery, or midwifery unless he was registered. Any breach of this regulation to be punishable by fine or imprisonment. 2. One licensing Board to be established in each division of the kingdom for the purpose of holding an examination of a uniform standard, and granting a registrable qualification in medicine, surgery, and midwifery. The Board should have no connection with the exist-

ing universities or licensing bodies. Its members to be appointed by the General Medical Council, which should have full powers given to it to regulate the scope and character of the examination. 3. Power to be given to the General Medical Council, in dealing with penal cases, to examine witnesses upon oath, and to inflict in certain cases a less grave form of punishment than that of erasure of the name from the *Register*. 4. That all criminal prosecutions under the Medical Acts should be undertaken by the police, and the fines obtained by convictions remitted to the Council.

THE SUGGESTED ONE-PORTAL EXAMINATION.

It may, doubtless, be considered by some that the suggested change is too sweeping and radical in its nature, and that due regard has not been paid to the position and interests of those bodies which at the present time grant the registrable qualification. To such persons a one-portal system, which would represent a combination of the universities and licensing bodies in the formation of a conjoint board of examination for each division of the kingdom, would be the best solution of the difficulty. Such a scheme would, it is admitted, be preferable to the existing state of things, under which, for example, we have here in Dublin four different bodies in competition for the purpose of granting a registrable qualification. But we must bear in mind that our claim for reform gains strength not in relation to professional interests, but in direct proportion to its public utility. I regret to have to admit that neither the public nor the State is very much concerned with the privileges or status of our profession, and as yet we are too weak, politically, to secure those rights to which we are justly entitled. It is only when a reform appeals to the selfish instincts of the community at large that it is likely to be carried; hence I hold that the principle of a State examination, securing as it would a minimum uniform standard free from the taint of anything that would suggest corporate competition, is the one which in the end would command most sympathy and support. I personally entertain no doubt as to its effects upon the profession and upon the future of the various colleges. To the profession it would bring higher status in proportion to the standard which was created, and less storm and stress in the battle of life. To the colleges it would mean the removal of the lowering if not degrading influence of a monetary competition, leaving them the source of those grades of professional honour which they could make attractive in proportion as they gave them in recognition of high attainments and distinguished merit. In most earnest loyalty to the College of which I am a humble member, and in whose hall we meet here to-night, I venture to express a hope that I will live to see the principle for which I have just contended realised.

A FURTHER NOTE ON THE TECHNIQUE OF
SERUM DIAGNOSIS.

By A. E. WRIGHT, M.D. DUBL.,
Professor of Pathology, Army Medical School, Netley.

THE method of employing capillary sedimentation tubes for the purposes of serum diagnosis may be claimed to possess certain advantages over all other methods. In the first place, it enables us to dispense with prolonged microscopic observations. In the second place, it is a method which may, where necessity arises, be employed even in the absence of a microscope. In the third place, it is a method which lends itself better than any other method to the quantitative determination of the intensity of the serum reaction. The following note deals with certain improvements that have been introduced into this method. These improvements relate first, to the method of collecting blood from the finger, and, secondly, to the method of diluting the serum.

THE METHOD OF COLLECTING BLOOD FROM THE FINGER.

The form of blood capsule which on the whole appears to present the greatest advantages is the form which is illustrated in Figs. 6 and 7.²

These capsules are readily made by heating a piece of glass tubing in two places in succession and drawing it out there into capillary tubes,

The lower of these capillary tubes is then bent up so as to form a somewhat acute angle with the long axis of the capsule. The only point that requires to be attended to here is that the capillary tube must not be allowed to kink at the point where it is bent over. This, however, may be effectually prevented either by blowing down the upper end of the capsule while the lower end is being bent up, or, more simply, perhaps, by giving the capillary tube time to cool down and then bending it over very slowly.

The blood capsule is filled in by capillarity and gravity in the manner which is indicated in Fig. 6. It is advisable only to half-fill the capsule. When a sufficient quantity of blood has flowed in, the flame of a spirit lamp or of a match is rapidly passed over the empty half of the capsule. Before the air, which has thus been rarefied, can contract again the tip of the capsule is to be sealed up as indicated in the figure. When the air now contracts it necessarily draws the blood up into the capsule in such a way as to leave the orifice of the bent tube quite free. The flame may now be applied to this end without any fear of overheating the blood.

METHOD OF DILUTING THE SERUM AND OF MIXING IT WITH THE BACTERIAL CULTURE.

When the serum has exuded from the clot, it is to be aspirated either into such sealed sedimentation tubes as are represented in Figs. 1 to 5, or into an unsealed sedimentation tube which has been fitted with a small rubber teat.³ The aspirating force which is required is obtained in the former pattern of tube by heating the air-chamber at the upper end of the sedimentation tube. In the latter pattern of tube it is obtained by compressing the rubber teat. In both cases this is to be done before introducing the tip of the tube into the blood capsule. Also before introducing the tip of the tube into the capsule a mark is to be placed upon the stem of the capillary pipette at any convenient point. One of the red wax pencils, which are employed for writing on glass, may be conveniently employed for this purpose.

As soon as the serum has run up as far as the point indicated by the mark the sedimentation tube is to be withdrawn from the blood capsule. The point of the tube is now to be introduced into a watchglass of normal saline solution. During the interval which necessarily elapses between the withdrawal of the sedimentation tube from the capsule and its introduction into the salt solution a bubble of air is to be allowed to pass up into the orifice of the capillary stem. This bubble of air is destined to serve as an index. Salt solution is now allowed to run up from the watchglass into the pipette until the distal end of the air-bubble index reaches the mark which has been placed upon the stem of the pipette. A volume of salt solution, which is exactly equal to the volume of the serum employed, will now have been introduced into the tube. A second air-bubble index is now required. This is readily obtained by merely raising the point of the tube out of the salt solution. As soon as it has been obtained the point of the tube is once more lowered into the fluid. In this manner the sedimentation tube can rapidly be filled in with any desired number of equal volumes of salt solution.⁴

For ordinary serum diagnosis, where a tenfold dilution is to be employed, the most convenient procedure is to introduce into the sedimentation tubes, first, one volume of serum, and then four equal volumes of salt solution. As soon as these have been taken in, the tip of the pipette is to be transferred to a watchglass containing the bacterial emulsion with which the serum is to be tested. Five equal volumes of this emulsion may now be allowed to pass into the pipette. The sedimentation tube will thus have been filled in, first, with one volume of serum, secondly, with four volumes of normal salt solution, and thirdly, with five volumes of bacterial emulsion. This will represent a tenfold dilution of the serum, for the serum will obviously stand in the relation of one to ten to the total volume of fluid which has been introduced into the tube. Mixture of the contents of the sedimentation tube is now readily effected by allowing the air in the air chamber to draw up the whole fluid into the mixing chamber. If sufficient aspiratory force has been obtained at the outset, this will effect itself in an automatic manner. If, on the other hand, sufficient aspiratory force was not then obtained (and this case can only occur where sealed tubes are employed), it will be necessary now to break off the upper capillary end of the air chamber. The air in the air chamber may then be reheated without in any

way displacing the contents of the tube. In either case, as each successive volume of fluid is drawn up into the air chamber, its air-bubble index will rise to the surface of the fluid, and will thus be got rid of. The escape of these air bubbles will be facilitated if (a) the mixing chamber has been made reasonably large, (b) if the tube is held vertically, and (c) if the tube is gently rotated between finger and thumb while the air bubbles are passing up through the fluid.

When the contents have been effectually mixed, the rubber teat is again compressed, or gentle heat is again applied to the air chamber, and the mixture of diluted serum and bacterial culture is driven down into the stem of the capillary pipette.⁵ The orifice of the capillary tube may then be immediately sealed up in the flame. The sedimentation tube is now placed upright in a test tube which has been labelled with the date and with the patient's name. The amount of the serum dilution may be conveniently inscribed upon the sedimentation tube. In all cases it is advisable to put up a control tube. This control tube is to be filled in with a mixture of equal volumes of bacterial culture and normal saline solution.

The procedure which has just been described is, as has been already said, the procedure which is most applicable to ordinary diagnostic purposes. Where, however, it is desired to make a quantitative estimation of the intensity of the serum reaction, it is preferable, first, to take into the tube one volume of serum, as is represented in Fig. 1; secondly, to take in four or more equal volumes of normal salt solution (as is represented in Fig. 2); thirdly, to expel the diluted serum into a clean watchglass; and, fourthly, to mix one volume of this diluted serum, or of any further dilution that may have been made from it, with one volume of the bacterial culture.

The most convenient form of culture to employ in these sero-sedimentation tubes is a dead bacterial culture. These dead bacterial cultures are easily prepared for use in serum diagnosis by the following method:—

First, young agar cultures of the desired micro-organism are emulsified with a measured quantity of sterile normal salt solution. Four to five c.c. of salt solution is an appropriate amount to add to each agar tube. The bacterial emulsions are then heated to a temperature of 60° C. for 10 minutes. Lastly, an addition of 0.5 per cent. carbolic acid is made to the dead cultures in order to prevent the possibility of accidental coagulation. The dead and carbolised cultures are now filled in from a syringe into capsules, such as those which are illustrated in Fig. 9 and Fig. 10. Or, where larger quantities of bacterial cultures are required—as, for instance, for class purposes, or for the daily routine of blood examinations—the dead cultures are simply introduced into such drop bottles as are represented in Fig. 8.

METHOD OF APPRAISING THE RESULTS OBTAINED BY THIS METHOD.

The positive or negative result of a blood examination can in an ordinary case be judged of in the course of twelve hours by a mere naked eye inspection of the sedimentation tubes. Figs. 4 and 5 represent respectively such a positive and negative result. In Fig. 5, which represents a negative result, the fluid has remained turbid throughout. In Fig. 4, which represents a positive result, the supernatant fluid has become completely clarified, and a floccular precipitate, which consists of masses of agglomerated bacteria, has settled down to the bottom of the tube. A less complete positive reaction would have manifested itself in the formation of a precisely similar floccular precipitate in an incompletely clarified fluid. Such a floccular and unevenly disposed precipitate can generally be distinguished at a glance from the evenly stratified and non-floccular precipitate which gradually accumulates at the bottom of the control tubes. A difficulty in discriminating these two varieties of deposit can only occur in cases where the characters of the bacterial deposit are masked by either a sediment of red blood corpuscles or by a sediment of flocks of serum globulin. The former source of confusion can be completely avoided by employing for serum diagnosis only serums that are free from blood corpuscles. The latter source of confusion can be avoided by employing normal salt solution, and not distilled water, as a diluting agent for the serum.

Where these fallacies have been admitted into an experiment, it will, however, still be possible to discriminate between a sediment that contains agglomerated bacteria and a sediment that contains only non-agglomerated bacteria. For this purpose recourse must be had to the microscope. A sample of the sediment may be obtained for examination by

breaking off—first, in the case of a sealed tube, the tip of the air chamber, and then afterwards the tip of the capillary stem. The sediment may also be examined in this way for agglutinated bacteria in cases where it is desirable to learn the result of a blood examination at the earliest possible moment. In such cases it will generally be found that, if the serum possesses the characteristic agglutinating

Fig. 1.—Sero-sedimentation tube which has been filled in with one volume of serum. Fig. 2.—Sero-sedimentation tube which has been filled in with one volume of serum and four volumes of normal salt solution. Fig. 3.—Sero-sedimentation tube which has been filled in with equal volumes of a fivefold diluted serum and of bacterial emulsion. Fig. 4.—Sero-sedimentation tube showing positive result of a serum diagnosis, that is, clarification of supernatant fluid and formation of a pellet of agglomerated bacteria at the bottom of tube. Fig. 5.—Sero-sedimentation tube showing a negative result; no clarification of supernatant fluid and no pellet of agglomerated bacteria.

Fig. 6.—Method of filling in blood capsule from finger. Fig. 7.—Showing method of sealing upper end of blood capsule. Fig. 8.—Drop bottle filled in with dead bacterial culture. Fig. 9.—Glass capsule filled in with dead bacterial culture. Fig. 10.—Method of emptying dead bacterial cultivation out of capsule.

power, evidence of agglomeration can be obtained in a very few minutes by examining a drop of fluid from the bottom of a sedimentation tube.

In conclusion, it is perhaps worth noting that the capillary sedimentation tubes may be used over and over again. They may be blown out after breaking off the ends. This done, they may be cleaned in distilled water. They may then be dried by passing them through a flame.

NOTES.

¹ It was recently pointed out by Surgeon-Major Semple and myself (BRITISH MEDICAL JOURNAL, May 15th, 1897), that the requisites for undertaking a serum diagnosis may be narrowed down to (a) a supply of dead bacterial culture, and (b) a supply of sero-sedimentation tubes—or, in lieu of the ready-made sedimentation tubes, a supply of material for making these, to wit, a few lengths of glass tubing and an alcohol spray, which can by ignition be converted into a blowpipe flame. ² These blood capsules, and all the other appliances for serum diagnosis which are described in this paper, may be obtained from Mr. A. E. Dean, jun., 73, Hatton Garden, E.C. ³ The beginner will find the mechanical appliance which is here figured very convenient; he can obtain the fine movements which are required for filling in the pipette by moving the controlling screw (B) and he can obtain the coarser movements which are re-

quired for mixing the contents of the tube by pressing directly with the finger upon the spring (A). My attention was called to this mechanical appliance by Mr. A. E. Dean. ⁴ This method of dilution is obviously applicable to a great number of other purposes. It is applicable, among others, to haemocytometrical and haemalkimetical purposes, and to the dilution of antitoxins. ⁵ If the total amount of fluid which is being dealt with more than suffices to fill the stem of the capillary pipette, the excess may at this stage be conveniently expelled from the tube.

ON THE DIAGNOSTIC VALUE OF TUBERCULIN.

BY

C. F. MARTIN, B.A., M.D., and G. D. ROBINS, B.A., M.D.,
Lecturer on Medicine, McGill University; Assistant Physician, Royal Victoria Hospital, Montreal.
Senior Resident Physician, Royal Victoria Hospital, Montreal.

It is now more than six years since tuberculin was placed before the medical world as a possible reagent for the diagnosis of tuberculosis. Its employment in the veterinary profession is now so universal and so productive of accurate results that not to use it is to stigmatise oneself as being of low rank in that profession. For human tuberculosis, too,