the clusters of swarm spores; that is, I could find no pigment in the latter; though it would have been difficult to do so even if it had been present, owing to the numbers of struggling flagellulæ. Since then I have examined no more grey mosquitos fed on tertian blood; and, indeed, have not been in a position to continue researches on malaria; but I have dissected thirty horse flies (*tabanus*), parasitic on tame pigeons, whose blood contained halteridium (Labbé). Though halteridium was found in the blood in the insects' stomach cavity, the stomach walls never contained pigmented cells; yet the histology of *tabanus* is closely like that of *culex*. It is interesting to note that in the four instances in which the cells have been found, their size is in proportion to the length of time between the feeding and death of the insects—namely, 7μ after two days, 17μ after four days, 19μ after five days, and 25μ after about a week.

I must correct an error due to the unfortunate use of formalin as a preservative in the two specimens sent to England. Pigment appears to have been found, not only in the pigmented cells, but also in the ordinary stomach cells, and elsewhere; and Dr. Thin draws the natural inference from this that the former are an altered form of the latter, induced by the absorption of malarial pigment from the stomach cavity. The effect is due to formalin, which not only blackens the green and yellow granules and globules always found in the tissues of *culex*, but converts traces of hæmoglobin in the stomach cavity into a multitude of minute acicular crystals very like pigment, and even has the same effect on the pigment of the pigmented cells. I know this to be the case from specimens in my possession, one of which now appears to contain pigment everywhere. I can say with confidence, and after special and repeated search for it, that when the specimens were fresh pigment was to be found nowhere else in them except in the pigmented cells; nor, except in the pigmented cells, have I ever found pigment in the stomach walls of mosquitos fed on malarial blood.

So long as the cells had been found only in two individuals (the only two properly examined) of a rare species of mosquito, it was quite possible to suppose them some kind of physiological cell, either pigment-producing or pigment-absorbing; the new facts show them to be too exceptional for this. If they be normal pigment-producing cells they should be found in most individuals of a species in one of which they have been observed; but Cases III and IV show that they may occur in a single individual out of many scores of the same species examined, while, except in these four instances, I have never observed them in allied diptera (*pulex*, *tabanus*, *musca*), nor even in allied species of *culex*. If, as Dr. Thin believes, they be ordinary epithelial stomach cells which have taken up the pigment of dead malarial parasites from the stomach cavity, they should, if found in one individual of a species fed on malarial blood, be found also in others of the same species similarly fed; but Case IV, in which they were observed in an insect fed on tertian blood, but not at all in many insects of the same species fed on crescent blood, is in direct opposition to this postulate—unless we can venture to assume that the stomach cells of this species are capable of taking up tertian pigment to the exclusion of crescent pigment. But there are now many reasons against Dr. Thin's theory. Thus, in Case III, the cells were much too small to be epithelial cells. Again, the pigmented cells are not like phagocytes in one important particular—they all contain roughly the same amount of pigment and an arrangement of it, which, as Dr. Manson says, forcibly recalls the arrangement of pigment in the malaria parasite. Phagocytes contain varying amounts of pigment, from one granule to several large clumps of it, according to the quantity they have been able to seize (compare phagocytes with parasites in malarial blood for example). Still further, if the pigment in the cells is due merely to phagocytosis, how comes it that they do not contain other granules from the stomach cavity besides, such as the yellow detritus of ingested blood corpuscles? Lastly, it remains to be proved that the stomach cells of *culex* are capable of taking up solid particles from the cavity. I have never seen any evidence of such, either in *culex* or in other families of diptera. Yet the stomach of the larva and imago contains multitudes of coloured vegetable and hæmoglobinous particles which could

easily be detected if contained in the interior of the cells of the stomach wall. I have never observed such a phenomenon. Why, then, should these cells absorb matter so exceptional in the ingesta as malarial pigment, and reject the numerous other granules which are much more commonly presented to them? The cells of the stomach are not scavengers like the phagocytes of the blood; what object can they possibly have in taking up the pigment of dead parasites, which otherwise will be naturally evacuated by the bowel? Or, are we to suppose that this dead pigment has the power of forcibly entering the cells? In short, I doubt whether this hypothesis will bear careful scrutiny.

Hence, the facts already collected, especially the new ones, make it difficult, if not impossible, to believe that the pigmented cells can be any kind of normal physiological cells, whether pigment-producing or pigment-absorbing cells. It would seem then that they must be some kind of pathological cells. What kind—whether new growths, or parasites peculiar to the mosquito, or a developmental form of the parasites of malaria in the mosquito—I cannot say. If not the last of these, it will always remain a curious coincidence that they should contain pigment so closely like malaria pigment, and that they should as yet have been found only in insects fed on malarial blood.

REFERENCE.

¹ No parasites found. Vide *Proceedings of South Indian Branch, British Medical Association, 1897.*

EUCHININ IN MALARIA.

By ST. GEO. GRAY, M.B. B.Ch. UNIV. DUBL.,
St. Lucia, West Indies.

SINCE last August I have been observing the effect of euchinin on malarial fevers, and, as far as my experience goes, I find it highly satisfactory in suitable cases.

Up to now I have only employed it in undoubted cases of malaria, where I find it superior to quinine in being tasteless and requiring a smaller dose to reduce the temperature. However, I must differ *in toto* from the statement reproduced in the BRITISH MEDICAL JOURNAL of December 11th, 1897, that euchinin does not cause cinchonism, for I have seen it cause tinnitus aurium, deafness, and derangement of vision and sensation in a more marked degree than the same dose of quinine.

Contrary to the statement of Professor von Noorden, that 15 grains of quinine are equal to 25 or 30 grains of euchinin, I find that euchinin is a more powerful antipyretic than quinine, and that, in malarial fevers at least, 10 to 15 grains of euchinin are as efficacious as 20 to 25 or 30 grains of quinine sulphate, and that it nearly always, in doses of 12 to 15 grains, causes buzzing in the ears if not other symptoms of cinchonism.

The largest dose I have given has been 15 grains once or twice a day, always commencing with a good purge, which I consider essential in the treatment of all malarial fevers. This is sufficient in most cases, following the treatment with tonics and change of air if possible after the temperature has remained normal for a few days. I append notes of a few cases:

CASE I.—August 11th, 1897. A. J., a young negro woman, had had quotidian ague for eight days; temperature 104° ; severe headache, constipation. She was ordered a purge and euchinin gr. xv twice a day. August 12th, temperature 98.4° . Patient states that the fever left her one hour after the first dose of euchinin. August 15th, temperature has been normal since evening of August 11th; fever has not returned, but patient feels weak. Ordered tonics. She had buzzing in her ears while taking euchinin.

CASE II.—August 11th, 1897. J. J., husband of above, says that he also suffered from fever every day. He took one of his wife's powders, and says that the fever left him after it, and has not returned. No buzzing in ears. I did not see him during the attack of fever.

CASE III.—August 14th, 1897. F. E., European, male, aged 34. Quotidian ague for five days. Temperature 102.8° . He was ordered a purge and quinine sulph. gr. xx, acetaminilid gr. v, once a day. On August 15th he was no better. Temperature 102.8° ; weak from purge. No tinnitus. He was ordered euchinin, gr. xij, at bed time. He could not sleep from buzzing in the ears and other unpleasant sensations, but the fever was gone next morning. He felt well but weak, and the temperature was 98.2° . He was ordered tonics. He has had one attack of ague since then, but was afraid to take euchinin on account of the cinchonism, which was not caused by the large doses of quinine with which he finally got rid of the fever.

CASE IV.—October 22nd, 1897. W. B., a young negro from the goldfields of French Guiana. Quotidian ague for several weeks. He was ordered

euchinin gr. xv once a day. On October 30th he returned to me to report that he had had no fever after the second dose. He said that he was taking a supply of euchinin back to Cayenne with him. He had some tinnitus, but not much.

CASE V.—November 8th, 1897. Rev. —, European, had tertian ague. He was ordered euchinin gr. xv once a day. The fever left after first dose, and did not return. He found it "stronger" than quinine, and said that half the dose was sufficient to keep away ague, while even that caused noises in the ears.

I have used euchinin in many other cases, but they are all similar to the above. To sum up:

1. Euchinin is as effective as quinine in malarial fever.
2. It causes cinchonism.
3. It is tasteless, therefore easily administered. This is its great advantage over quinine.

The readiest form of administering euchinin is the simple powder placed dry on the tongue and washed down by a little water. As it is very bulky, some patients prefer it in cachets; but all solutions of euchinin that I have seen are decidedly bitter, presenting no advantage whatever over quinine.

I do not consider cinchonism such a fatal objection as the intensely bitter taste of quinine. My experience of malarial fevers is that quinine and the malarial poison being antidotes to one another, cinchonism is the sign that a sufficient quantity of quinine has been taken to overcome its antagonist the malarial poison, as mercurialism is the sign that the syphilitic poison is under control.

If euchinin can be proved to be even nearly as effectual as quinine, its tastelessness alone should recommend it to many as a substitute, notwithstanding that it is not altogether free from some of the other objections to quinine.

REMARKS ON A CASE OF HORSESHOE KIDNEY SIMULATING MALIGNANT DISEASE OF THE ABDOMEN.

By THOMAS OLIVER, M.A., M.D., F.R.C.P.,
Physician to the Royal Infirmary, Newcastle-upon-Tyne.

THE following case is reported with the view of directing attention to the simulation of malignant disease by an abnormal kidney, and of the need for physicians and surgeons giving consideration to the possibility of the existence of such an abnormality before undertaking an exploration of the abdominal cavity. In most cases it must be that the existence of a horseshoe kidney is diagnosed only after the abdomen is opened. It was so in my own patient in whom there were no physical signs that strongly suggested such an abnormality.

Wm. K., aged 51 years, a coal miner, was admitted into the Royal Infirmary, Newcastle-upon-Tyne, complaining of headache and loss of appetite of seven years' duration, and of pain over the region of the stomach and down the spine of two months' duration. There is nothing in the family history that calls for comment. He has been married thirty-one years, is the father of eleven children, one of whom died in infancy. His wife has had two miscarriages. He has had a good home and plenty of food: he has been a teetotaler for the last eighteen years. As he has suffered much from indigestion he has been extremely careful in regard to his diet, particularly within the last twelve months, taking principally rice, fish, milk, and fresh eggs, etc. At the age of 24 years he had pneumonia; seven years ago he had influenza, and for the last ten years he has been more or less a dyspeptic.

Two months ago, while hewing coal in the pit and working in a bent position, he was seized with pains in the region of the umbilicus. It was so severe that he felt faint and was obliged to lie down for a little. He could work no more that day on account of pain, which by degrees gradually subsided. Afterwards he only felt the pain after eating, and although it often made him feel sick it only caused him to vomit once about a month ago.

He is on the whole a well-nourished man, but he states that within the last few weeks he has lost flesh considerably. Myxœdema is well marked. He has the jaded look and mental depression so frequently observed in old dyspeptic subjects. His tongue is dirty brown and coated. Chest: Heart and lungs are healthy. Abdomen: On inspection very distinct pulsation can be observed over a circumscribed area of 2 ins in diameter just above the umbilicus. A hard rounded mass with smooth borders can be felt, through which apparently the pulsation of the aorta is being transmitted. The tumour lies rather to the right than to the left of the aorta, and is felt to move up and down with respiration. It is painful on deep pressure, but is not attended by any sickening sensation when handled nor by a feeling of faintness. As the patient lies on his back a soft blowing systolic murmur can be heard through the stethoscope placed a little above rather than either over the tumour or below it. On placing him in a sitting posture the pulsation can hardly be detected and no *bruit* is thus heard. Turned on his face there is no longer pulsation felt when the hand is placed in the umbilical region until it is plunged deeply in, and even then it is not very distinct. On auscultating the spine no *bruit* is heard nor is pain elicited on percussing the spinal column. The legs are well developed; the knee-joints are normal. The bowels are

regular and there is no trouble with the bladder. The urine is acid Specific gravity 1028, no albumen, no sugar.

A diagnosis was never really made. Taking into consideration the history of the case, the long continued dyspepsia, recent pain, the presence of a circumscribed solid tumour in the abdomen almost in the middle line and movable during respiration, the age of the patient and his emaciation, and the fact that the physical examination excluded aneurysm, it is only right to say that the probability of malignant disease suggested itself, although there was nothing that helped us to locate the tumour in any particular organ in the abdomen. It was a case in which there was a large element of doubt in the diagnosis, and as it was evident that medicine could do nothing but palliate, and as the patient on account of pain was unable to follow his employment, he himself pressed for an abdominal exploration so that his mind might be placed at rest. My surgical colleague agreeing with me that the operation would at any rate enable us to make a complete diagnosis, Mr. Williamson on July 27th, 1897, opened the abdomen by an incision 6 ins. in length slightly to the right of the middle line. After examining the contents of the cavity the tumour was found to be a horseshoe-shaped kidney, the central part of the arch of which was lying upon the aorta. The abdominal incision was at once stitched up and in ten days thereafter the patient was practically well again and has remained so since, having lost all his pain and other unpleasant symptoms.

The question naturally arises, Was the operation justifiable? The end justifies the means, and as the end obtained was the cure of the patient, mental and physical, quite apart from the making of a diagnosis, I have no hesitation in answering the question in the affirmative. It is more than likely that strain at work loosened the attachments of the kidney, and by pulling upon delicate nerves caused the severe pain and feeling of faintness. Probably it was in consequence of this strain that the arch of the fused kidneys became movable. The range of movement was extremely limited, up and down during respiration only, and in extent nothing like what is observed in ordinary floating kidney, which has frequently been mistaken for a tumour of the omentum or mesentery. The possibility of a horseshoe kidney misleading the physician in his diagnosis is thus alluded to by Mr. Henry Morris in *Surgical Diseases of the Kidney*, p. 104: "It might easily happen that a horseshoe kidney or two kidneys more or less completely fused and situated in the median line of the abdomen or in some part or other of the pelvis might give rise to the opinion that a morbid growth existed." It was rather a morbid growth than an abnormal organ which the physical signs and symptoms presented in my patient, and as one horseshoe kidney is met with in every 1,600 *post-mortem* examinations, the value of my remarks will be enhanced if they direct attention to the necessity of giving some consideration to the possibility of fused kidneys in the diagnosis of obscure tumours of the abdomen.

SYMMETRICAL PARTIAL DETACHMENT OF FINGER NAILS FROM THEIR MATRICES.

By J. STANLEY KELLETT SMITH, F.R.C.S.ENG.

THE accompanying photograph (p. 553) illustrates a curious condition of symmetrical partial detachment of the finger nails from their matrices.

The patient, a young married woman, aged 30, states that about a year ago, without any apparent cause, the skin beneath the free edges of the affected nails began to peel away in whitish shreds. This continued for three to four months, at the end of which period the excavations beneath the nails had reached their present limits.

The fingers attacked are the index, middle, and little fingers in each hand. The nails themselves are lustrous and healthy looking, growing with an even free edge, and presenting no scaly, fibrillar, or opaque appearance. Over the area of separation, into which a probe can easily be passed, the matrix looks glazed, and is cicatricial to the touch. So marked is the symmetry of the condition that not only are the corresponding nails in the two hands affected, but the