rabbit, guinea-pig, white rat and mouse, on the other hand, are more or less depressed. Even in the same species of animals there is a considerable variation in the action of morphine, and men who are most susceptible to its depressant effects are sometimes markedly excited by it. Comparatively recently Amblard and Grasset have returned to the question of the convulsant action of morphine, and from researches on animals with cut spinal cords have concluded that morphine and thebaine are not diametrically opposite in action and that the effect of morphine on frogs and mammals is not dissimilar. This was known before, but these authors lay stress on the fact that the excito-motor action of opium is not exclusively dependent on the amount of so-called convulsant substances it contains, but it is also dependent on the morphine present. An amount of opium containing 5 cg. morphine is more convulsant than the amount of thebaine (about 1 mg.) present.

But whatever the effects in other animals, to man and the dog morphine is decidedly depressant, and differs markedly in its action from thebaine. And according as the other alkaloids of opium approach nearer to one or other of these in action, they may be classified. Claude Bernard, commencing with the most convulsant, obtained the following series: narceine, morphine, codeine, narcotine, papaverine, thebaine; the order of toxicity he found was narcotine, morphine, narceine, papaverine, codeine, thebaine, the last named being the most powerful.

Schröder divides the alkaloids of opium into two groups: the morphine group containing morphine and oxydimorphine, or, as it is now sometimes called, pseudo-morphine; the codeine group containing papaverine, codeine, narcotine, and thebaine. Hydrocotarnine, laudanose, and cryptopine also belong to this group, but our knowledge of their actions, Schröder says, is not sufficient to place them. Meyer, in connection with a research on other papaveraceous alkaloids, has modified this classification. He divides them thus: 1. Morphine group, containing morphine, chelidone, and α -homochelidone. 2. Codeine group, containing papaverine, codeine, laudanose, narcotine, hydrocotarnine, sanguinarine, thebaine, laudanine, hydrastine. 3. Protopine group, containing protopine, β -homochelidone, and cryptopine. Not one of these classifications seems to me satisfactory. Claude Bernard's has been criticised by many observers, mainly in connection with the relative positions of morphine and narceine. The latter in man has not been found by the majority of observers to be more narcotic than the former, and in practice it has proved of little value. The position of papaverine in the two classifications also seems to me erroneous. In both it is given as a convulsant, and in Bernard's as a marked convulsant. In my experiments the prominent symptom has been depression.

In frogs spasmodic inco-ordinate movements may sometimes be obtained on stimulation for a short time after the administration of a lethal dose, but at no time is any tetanus or marked increased activity present. In rats 0.3 gram per kg. produced muscular weakness, slowing of the pulse and respiration, but no tetanus or distinct narcosis; the symptoms lasted four hours, and recovery occurred. Still larger and lethal doses failed to produce increased reflex excitability. In a guinea-pig 0.2 gram (0.47 gram per kg.) produced prostration and running movements of the limbs, which lasted 30 seconds, but these often occur after the administration of alcohol, and they cannot be regarded as certainly indicating a convulsant action. This animal died from paralysis of the respiration in ten minutes. A rabbit to which a similar dose (0.48 gram per kg.) was given also died in ten minutes, and in this case the heart and respiration ceased together. No sign of narcosis, rigidity, or tetanus occurred.

It seems to me that these experiments show that papaverine is not possessed of any convulsant action in any sense of the word. As far as I can see its action is mainly depressant, and as the nerves still retain their irritability after death, the effect is apparently due to an action on the central nervous system.

My experiments with the other alkaloids corroborate, on the whole, those of previous observers. Thebaine is almost a pure convulsant, and the same may be said of laudanine and laudanose. Hydrocotarnine, protopine, and codeine also possess convulsant properties, and, in frogs and many mammals, so does morphine. Protopine produces more clonic convulsions than the other alkaloids, but this does not seem sufficient to justify its being placed in a separate division. Narcotine and cryptopine I have not found markedly convulsant, and narceine seems to possess no action whatever in this direction.

V. Schröder found it without effect when given *per os* to a rabbit; and in a kitten to which 0.1 gram per kg. was given by the mouth I obtained no

action. Injected intraperitoneally into a rat, 0.5 gram per kg. (1.25 c.cm. of 2.5 per cent. solution), only produced slight and transient depression, and in a frog 0.02 gram produced after some time depression, followed in a few days by death. Previous to death the animal became slightly tetanic when touched.

The order of toxicity, according to my experiments is — narceine, narcotine, morphine, codeine, papaverine, hydrocotarnine, laudanose, laudanine, protopine, and thebaine. As regards therapeutic utility, none of these substances except those already in use (morphine and codeine) are likely to prove of value in practice.

[The lecturer at this point discussed the chemical constitution of the opium alkaloids, and then proceeded as follows:]

The general action of morphine is well known to all of you, and I have nothing new to add as to its effects. It cannot be employed habitually as a hypnotic, mainly on account of the rapid tolerance which is produced, and also on account of its tendency to induce a habit and produce untoward effects on the system. But whenever pain is present morphine is the hypnotic *par excellence*.

With morphine we have no longer to deal so much with a depressant effect on the circulation, as in the case of chloral and its allies, as with an effect on the respiration. This comes out in nearly every tracing taken with this drug. At first there is often increase in the frequency and depth of the respirations, and sometimes this effect may last a considerable time; but during the sleep produced by the drug the respirations are almost invariably slower, and often shallower. Later they may become irregular and rapid, or may take on a periodic rhythm.

The effect of morphine on the circulation varies somewhat, Guinard says, according as the animal is narcotised or not by the drug. In animals which are narcotised, such as the dog, the blood pressure at first rises, then falls below the normal to the extent of about 20 mmHg. In animals not narcotised there may be a preliminary fall, but the pressure afterwards rises and remains above the normal level until near death. The heart itself, Guinard believes, is stimulated directly, more especially in those animals which are not narcotised by morphine. After large doses in the dog the heart-beat sooner or later becomes irregular. The irregularity he is inclined to attribute to a central action, as section of the vagus caused it to disappear. This is of importance in connection with the use of morphine as a hypnotic in advanced heart disease, for although in this condition it is one of the best hypnotics we possess, it is by no means free from danger. Attempts have been made to find a less dangerous and equally effective substitute. To two such compounds I intend to draw your attention in the next lecture.

REFERENCES.

- ¹ Arch. d. Pharmacodynamie, iii, 191. ² Münch. med. Woch., 1898, xxxvii.
- ³ Rev. Méd. de la Suisse Rom., 1893, pp. 549, 628. ⁴ Rif. Med., 1888, No. 81.
- ⁵ Ther. Monats., 1889, p. 515. ⁶ Zeit. f. Phys. Chem., xiv, p. 52. ⁷ Deut. med. Woch., 1896, 185. ⁸ Ther. Monats., 1896, 421. ⁹ Diss., Halle, 1896. ¹⁰ Münch. med. Woch., 1897, 355-391. ¹¹ La Morphine et l'Apomorphine, 1898.

CASES OF SEPTICÆMIC INFECTION TREATED WITH ANTISTREPTOCOCCUS SERUM : RAPID RECOVERY.

By J. MITCHELL BRUCE, M.D., F.R.C.P.,
Physician to Charing Cross Hospital.

By the kindness of two of my professional friends I am able to submit the histories of two cases of an infectious kind which I had an opportunity of seeing with them, and in which the use of antistreptococcus serum was entirely successful.

SEPTIC ENDOCARDITIS.

(Notes by ALEXANDER T. SCOTT, M.R.C.S.)

Mr. X., 39, married, a merchant, had suffered from protracted chorea in childhood. About the age of 14 he had a neurotic attack following a blow on the abdomen at cricket. At 17 he had diphtheria. The present illness commenced about the middle of December, 1897, with "smoky" urine and a feeling of "seediness." For about twelve months previously I had noticed that his complexion, which was usually sallow, had assumed gradually a cachectic tint; indeed, I had frequently asked him how he was, but only to be told "Quite well."

On February 13th, 1898, at 10.30 P.M., he had a shiver and vomiting. This was repeated several times during the night. Next day I visited him. I found the temperature 101°, and the pulse 100. There was hæmat-

CHART I.—Mr. A. T. Scott's case.

CHART II.—Dr. Abercrombie's case.

uria, and he had pain in the loins, right knee, and left forearm. He continued in this condition for about ten days, when the temperature became normal, and the pains disappeared, and he proposed to return to business. During this time he had no rash. He was now apparently approaching convalescence. The urine improved considerably, and on two occasions was found free from albumen or smokiness. However, on March 3rd, all the symptoms returned, and an eruption resembling petechiae appeared on the hands and forearms; but in two or three instances, especially one on the back of the right wrist, looked like small furuncles with sloughy centres of minute size. At about 2 P.M. that day the temperature, which had been 100.5° in the morning, rose to 103.4° with profuse sweating. There were no pulmonary or cardiac complications, and the pulse remained steady and not proportionately accelerated. The case was treated with 15-grain doses of sodium salicylate every four hours and cold sponging. The daily rise of temperature went on till March 6th, when an apparent defervescence set in again until March 13th, when a further rise took place, this time to 103.4°, on March 15th to 104°, and on March 16th to 105°, with increase of rash. A peculiarity in the case was a sudden rise of temperature in the afternoon with lower ranges at night. The high temperature was always preceded by an intense feeling of dry heat followed by profuse sweating.

On March 17th Dr. Mitchell Bruce saw the case in consultation. The temperature was then 104.8°; there was an eruption over the greater part of the body, partly erythematous partly resembling stigmata; hæmaturia still existed. A soft systolic murmur was audible over an area of about a

square inch in the mitral region. The diagnosis was that the case was septic, with recent endocarditis.

An injection of antistreptococcus serum was given at 6.30 P.M. At 6.30 A.M. on March 18th the temperature had fallen to 98.7°, and a second injection was given. By 6.30 P.M. the rash had faded, and the murmur was only just audible; a third injection was given, the temperature being 100°. On the morning of the 19th the temperature was normal and the rash gone, leaving only minute dried-up sloughs.

The patient rapidly recovered, having no subsequent drawbacks. The temperature chart is appended. (CHART I.)

The case just related cannot be considered complete without reference to another case with which it was associated in a remarkable way, although the exact bearing of the two on each other can be only a matter of speculation in the present state of pathological knowledge. While this patient's illness was in progress, his two children (boys, aged 12 and 9) had measles; and after recovery I operated on the elder boy for post-nasal adenoids on March 6th, and he made a good recovery. The younger boy had his right tonsil removed on March 14th, and that afternoon, being fretful, was allowed to lie on his father's bed for a time. This was three days before Dr. Bruce saw Mr. X. On March 19th it was discovered that the wound on the boy's tonsil looked sloughy, and that his temperature was 103°. The case being highly suggestive of infection, an antistreptococcus injection was given on the evening of March 23rd, with no benefit either locally or constitutionally.

An injection of antidiphtherial serum was administered on the even-

ing of March 23rd, as the case seemed obviously diphtherial. (A specimen of membrane sent to the British Institute of Preventive Medicine was said to contain no diphtheria bacilli.) Next day tracheotomy was performed, followed by another injection of antidiphtherial serum. The patient made an uninterrupted recovery. A portion of membrane sent on the day of the operation to Messrs. Burroughs and Wellcome's laboratory was reported: "Much diphtheria." Until the injections were used the throat had been painted with zinc chloride and carbolic acid, and insufflations of precipitated sulphur also employed.

The lessons to be learned from the case, as far as one can reach a conclusion safely from a single case, are: (1) That antistreptococcus serum may succeed in septicæmia after ordinary measures have failed; (2) that it may act with great rapidity, all the phenomena disappearing in the course of forty-eight hours.

SEPTICÆMIA OF CEREBRO-SPINAL TYPE. (Under the care of Dr. ABERCROMBIE.)

J. R. B., aged 15, clerk, was admitted into Charing Cross Hospital on May 31st, 1898, for fever with delirium and drowsiness. The previous history was unimportant. He had been ill for twenty-six days. The first symptoms were a feeling of drowsiness which lasted for three days, then nocturnal restlessness, vomiting, and the development of the phenomena which continued prominent throughout. These were: Fever, ranging between normal or subnormal and 105°, with occasional rigor; pains in the head and down the back, and tenderness of the head, muscular spasms and a degree of opisthotonos, and bouts of restlessness, insomnia, and incoherence or active delirium.

A striking feature of the case, next to the cerebro-spinal type of the phenomena, were periods of remission, each extending over one or two days; the boy looking well and feeling well, calling for food and anxious to be permitted to read; and the sudden interruption of this condition by rigor, rise of temperature, possibly sickness, restlessness, and incoherence, all passing off again presently. The urine was found albuminous repeatedly. Epistaxis occurred on three occasions. On June 7th (thirty-third day of illness) a soft systolic murmur was heard over the apex of the heart; next day it had disappeared; on the third day it was caught again, and again it disappeared; but later on, June 18th (forty-fourth day) it was well developed and partly presystolic in time.

On June 13th (thirty-ninth day) 10 c.cm. antistreptococcus serum were injected under the integuments of the abdomen. The temperature fell steadily, but this proved to be only one of the ordinary periods of remission, for although a second injection was given next day, the temperature rose again on the third day, and two days later still the patient was as ill as ever, with a temperature of 105.2°, vomiting, delirium, severe headache, retraction of the head, and well-marked mitral murmur. Quinine was now tried (grs. v. bis die), and the temperature fell as usual, only to rise again to 104° nearly. (Chart II, p. 77.)

On June 22nd (forty-eighth day) a second course of antistreptococcus injections—the serum obtained from another source—was commenced, six in number. The temperature never again reached 100°, becoming practically normal, and therewith the patient improved steadily. The cardiac murmurs continued audible for a few days, then was heard only at intervals, and finally disappeared. All the phenomena connected with the head and spine subsided at the same time; and the boy was discharged on July 20th (seventy-sixth day) perfectly well, having gained 6 lbs. in the last sixteen days in hospital.

The conclusions to be drawn from this case are obvious: (1) As before, that in some cases of a septic kind antistreptococcus serum is successful; (2) as before, that the effect may be

extremely rapid; (3) that success may be obtained after other measures failed; (4) that success may be attained with one serum after another serum has failed. This is very important. There was no suggestion of the first serum being bad. It was derived from an unimpeachable source. I regard it simply as different in some occult respect; and after all the test of difference is the result. One practical lesson to be learned from this case is not to be content with the trial of one serum if it prove unsuccessful.

TREATMENT OF A CASE OF PUERPERAL FEVER BY ANTISTREPTOCOCCUS SERUM.

By DAVID ARTHUR, M.D., C.M., D.P.H.

Mrs. P., aged 25, was confined of her first baby on August 5th, 1898. Everything was apparently normal. I first saw her about three hours after delivery, examined the placenta and found it entire, and the patient felt very comfortable. The nurse was not certificated; she was stated to be "very clever" and thought it unnecessary to send for me. All went well till the third day when the patient felt ill, complained of thirst, headache, and a sense of fullness in the lower part of the abdomen. There was no rigor; her temperature registered 103° F.

I douched the uterus with 1 in 4,000 mercury perchloride, and gave a mixture of iron perchloride and quinine. The douching of the uterus was continued for the following seven days thrice daily; between times the nurse douched out the vagina with the same lotion. On the third day of the fever intractable diarrhoea commenced, which lasted during the whole illness and was only partially controlled by the use of lead and opium pills. The patient gradually became weaker, all attempts to reduce the fever ended in failure, and the end seemed near.

Dr. Robinson was called in consultation on the eighth day of the illness. We found the patient in a very weak condition, the uterus extended above the umbilicus, and felt soft and boggy; there was only slight tenderness on pressure. The whole of the vaginal wall, the labia minora and majora were covered with a yellowish membrane, diphtherial in appearance, and the sense of touch gave the impression that the same membrane extended right up into the uterus. We both thought the patient was dying. Dr. Robinson, judging from the nature of the membrane, thought it due to a pure streptococcus infection; this I afterwards confirmed by a cover-glass preparation.

As the uterine douching and medicinal treatment had failed, we decided to stop all treatment, with the exception

Chart of Dr. Arthur's case.