REMARKS ON THE SURGICAL TREATMENT OF PYLORIC OBSTRUCTION, WITH AN ACCOUNT OF TWENTY CASES, AND OF A NEW WAY OF PERFORMING GASTRO- ENTEROSTOMY.*

By F. T. PAUL, F.R.C.S.,
Surgeon to the Royal Infirmary, Liverpool.

STENOSIS of the pyloric orifice of the stomach is brought about in several different ways. It is due to—

1. Congenital insufficiency.
2. Obstruction from the external pressure of tumours, bands, etc.
3. Reflex spasm.
4. Fibrous contraction.
5. Cancerous and other growths.

In dealing with these conditions I propose to restrict my remarks to personal experience.

CONGENITAL INSUFFICIENCY

I have no positive knowledge of in the adult, unless it was the primary cause of stenosis in Case IV.

OBSTRUCTION FROM WITHOUT

in such a degree and under such circumstances as to suggest surgical interference is very rare, owing to the mobility of the stomach. I have known omental bands connected with a hernial orifice and acting through the transverse colon to so drag on the stomach as to give rise to symptoms of obstruction. In such a case the treatment is obvious and simple. Scirrhus cancer of the gall bladder also drags on the pylorus, but in the opposite direction. It is somewhat rare. I have seen two cases in which obstruction was caused in this way.

REFLEX SPASM

occurs both in association with ulceration and independently of it. In 1896 I operated upon a case in which symptoms of pyloric obstruction apparently depended upon muscular spasm only.

CASE I. The patient was a man, aged 21. He was admitted into the Royal Infirmary under the care of Dr. Caton in November, 1895, for pain in the epigastric region and a lump. The lump proved to be a largely dilated stomach, in which splashing sounds could be easily produced. He was treated by careful dieting and washing out the stomach, but not having materially improved in January, 1896, was transferred to me. The patient was anæmic, and decidedly of neurotic type. On Jan 6th the abdomen was opened for exploratory purposes. The stomach was found dilated, the pylorus thick but not fibrous, and upon invaginating the anterior wall upon the forefinger into the pyloric orifice abnormal resistance was met with. This was gradually overcome by continued steady pressure, and ultimately the opening was over-dilated without splitting by passing three fingers through it. No cut was made in the organ. The patient healed well, but was subsequently attacked with acute mania, and had to be sent to the County Asylum, where he remained five months before he got his discharge. He had no recurrence of the gastric symptoms.

FIBROUS STRICTURE

is a condition presenting the greatest possible interest for surgical treatment. It is secondary to ulceration or other inflammatory change in the walls of the stomach. Out of nine cases, all but two have occurred in men, although gastric ulceration is so much more common in women. None were traumatic. Perhaps fibrous stricture with or without preceding ulceration may sometimes be of syphilitic origin, a point to which I drew attention at this institution in 1881.¹

Whatever the cause of these strictures, they sometimes lead to almost complete occlusion, with extreme gastric dilatation and great emaciation of the patient. Several operations have been suggested and employed for the cure of fibrous stricture, including pylorotomy, divulsion, pyloroplasty, and gastro-enterostomy. Pylorotomy ought not to be practised for a stricture known to be of simple nature, unless under some exceptional circumstances, and the same may be said for divulsion or Loreta's operation. The choice therefore lies between pyloroplasty and gastro-enterostomy. I have never employed pylorotomy for simple stricture, but

have used divulsion in two cases in addition to the one just described.

LORETA'S OPERATION.

CASE II.—P. K., aged 36. Fibrous stricture of pylorus. Two years ago vomited blood in small quantities on four occasions. Eleven months ago first began to suffer from the collecting of "burning wind" in his stomach, with nausea but no vomiting. He gradually gave up various articles of diet, living mostly on bread and milk. He lost 3 st. in weight. Was admitted under Dr. Caton in November, 1890, and was for some time under medical treatment for obstructed pylorus with dilated stomach. He was transferred to me in January, 1891. On January 7th the abdomen was opened, a fibrous contraction of the pylorus found, which was forcibly dilated with the forefinger. The viscus was not opened, its anterior wall being pushed before the finger. The stricture gave way suddenly, the coats rupturing nearly to the peritoneum. The wound did well for a few days, and then discharged rather a lot of pus from the upper end, and subsequently healed. He gradually got on to common diet, and left the infirmary cured on March 30th. During the succeeding two years he was twice readmitted with complaints regarding the "burning wind," but as he ate common diet heartily without vomiting, and was a man of very weak intellect, nothing more was done. I thought his symptoms greatly relieved, but not nearly to the extent they would have been by the more modern pyloroplasty, for which the case was eminently suitable.

CASE III.—G. S., aged 41. Fibrous stricture of the pylorus. For four years this patient had suffered from marked symptoms of pyloric obstruction, during which time his weight had dropped from 12 st. to 8 st. He had been under the care of Dr. Glynn, and was transferred to me in April, 1892. On April 5th the abdomen was opened, and a thick tough fibrous stricture discovered, through which I was unable to pass a finger, even after incising the anterior wall of the stomach. Ultimately the stricture was overcome by the use of Hegar's uterine dilators, but again it was not a genuine dilatation, as it was felt to be ruptured on the under side. Progress was pretty good at first. The patient recovered sufficiently to get up and commence going out, but he was never thoroughly comfortable, and on May 24th, exactly seven weeks after the operation, he had a fatal hæmatemesis. At the necropsy it was seen that the gastro-duodenal artery had been exposed on the floor of the ruptured stricture, and was now opened into by the ulceration which had supervened.

Of the three cases of Loreta's operation, in two only was it employed for the treatment of genuine fibrous stricture. The first was certainly benefited, but the other died from an accident for which the operation was directly responsible. Even the first probably ran considerable risk, for the suppuration is quite as likely to have originated with the injury to the pylorus as in the superficial wound, and, though the patient was relieved, he was not thoroughly cured. In slight cases, where the stricture is chiefly spasmodic, dilatation without cutting into the stomach may be beneficial, and is practically devoid of danger; but for tough fibrous strictures the day is past for using divulsion, which must give place to the more effective, and I believe less dangerous, operations of pyloroplasty and gastro-enterostomy. In choosing between these, the choice will now always fall upon the former whenever it is possible to do it thoroughly and efficiently, the latter being reserved for those cases in which, owing to great thickening and fixation of the parts, or the presence of still active ulceration, it is undesirable or impossible to adopt pyloroplasty. I have performed pyloroplasty for fibrous stricture five times, with one death, the death being due to peritonitis pure and simple, the stomach wound having closed perfectly.

PYLOROPLASTY.

CASE IV.—G. D., aged 31. Fibrous stenosis of pylorus; pyloroplasty; recovery. Ever since he was a little child of about 6 has had to be very careful in regard to food, living chiefly on milk. For several years has often vomited a large quantity at night before he could sleep, and for the last eighteen months has regularly emptied his stomach with a tube and washed it out. Has been steadily losing flesh, and is now very emaciated, as he is above average height and weighs only 8 stone. Was sent to me by Dr. Masson, of St. Helens. October 8th, 1896, pyloroplasty was done. The pylorus was very much constricted, but no great thickening and hardness, so that a good band of healthy mucous membrane could be brought into the pyloric orifice. He made excellent progress. On October 24th had chicken and potatoes and rice pudding for dinner. On the 31st was discharged weighing 9 st. 5 lbs. When seen a few months later was in excellent health weighing 11 st. In 1897 he was accepted as a first-class life in the Lancashire Alliance Insurance Company, and has since remained quite well. Altogether this proved one of the most satisfactory surgical cases I have ever met with.

CASE V.—W. A., aged 26. Fibrous stricture; pyloroplasty; recovery. Sent to me by Dr. Pinegan. First began with indigestion five or six years ago. For the last two or three years has suffered from periodic vomiting. Is thin but not emaciated. Has a considerably dilated stomach, in which splashing sounds are easily produced. On January 20th, 1896 explored and found a slight fibrous stricture. Pyloroplasty was easily done, and the recovery was uneventful. On February 15th he was discharged apparently quite well, and when last seen a few months later was still in good health.

CASE VI.—W. H., aged 38. Tough fibrous stricture; pyloroplasty; recovery. Fourteen years ago was said to have had an attack of biliary colic. The attacks occurred subsequently about every six months. Has also suffered much from indigestion. Was admitted under Dr. Caton, February 1st, 1897, with all the symptoms of pyloric obstruction and

* Read at the Liverpool Medical Institution.

dilated stomach, possibly complicated with gall stone. He was not much wasted. Patient improved considerably under medical treatment, but the symptoms of obstruction remaining he was transferred to me on April 1st, 1897. On April 5th pyloroplasty was done. The pylorus was found thickened, hard, and adherent. The operation was satisfactory, but not quite so much as in Case IV. He made an uninterrupted recovery. Commenced common diet April 27th, and was discharged cured, May 6th. He was seen several months later by Dr. Gullan, and was then in excellent health. A report just to hand states that he has been perfectly well ever since.

CASE VII.—Mr. J., aged 46. Old fibrous stricture; pyloroplasty; recovery. Subject to indigestion with pain since about 20. Had attacks of vomiting about every two months, always after food that had disagreed with him. Often used to make himself vomit to obtain relief. For the last five or six years stomach symptoms have been steadily increasing in severity, and in March, 1897, they became decidedly worse. Vomits now three times a week, and is much wasted. In September Dr. Wilson Paton asked me to see patient with him with a view to operation. There was no doubt he had pyloric obstruction with considerable gastric dilatation, but no tumour could be felt. On September 14th the abdomen was explored, and old fibrous contraction of the pylorus being found, pyloroplasty was done, though the parts were too hard and adherent to permit of a perfect result. He recovered well, and was greatly benefited for a time; but has lately had a certain amount of recurrence of symptoms of obstruction. Whether this is due to some subsequent contraction, or to the development of malignant growth not detected at the time, or to careless habits in diet which are undoubted, I cannot say; but should the symptoms continue I am afraid nothing less than a gastro-enterostomy would afford him any permanent relief. A later report states that he is much better again, and gaining weight.

CASE VIII.—Miss H., aged 42. Fibrous stricture; pyloroplasty; death from peritonitis. Had suffered for many years from what she regarded as bilious attacks, which of late had been accompanied with a good deal of pain in the region of the gall bladder, but not so acute as the pain of biliary colic. Vomiting was severe, and during the attacks almost continuous. On October 13th, 1897, I explored the abdomen with the assistance of Dr. McAfee. There were no gall stones, but the pylorus was scarred and contracted. Pyloroplasty was easily done, and I felt no anxiety as regards the result. The latter, however, was very disappointing. The operation was followed by septic peritonitis, from which she died on the 15th. At the post-mortem the pyloroplasty wound was found to have healed perfectly, the peritonitis evidently having originated directly from the operation.

GASTRO-ENTEROSTOMY FOR FIBROUS STRICTURE.

CASE IX.—E. F. aged 24. Active ulceration; gastro-enterostomy; recovery. Patient was a very healthy man until eleven months ago, when he began to vomit after food. He dieted himself carefully, and worked until eight months ago, when pain in the epigastric region set in, and the vomiting continued. The pain became very bad both night and day. He was for three months under the care of Dr. Davidson, in the Royal Infirmary, and later in the Northern Hospital, under Dr. Dickinson, where clots of blood were washed out of the stomach on two or three occasions. He went from bad to worse, and was sent to me by Dr. Finegan in a very wasted and broken-down condition. In the dresser's report he is described as "an emaciated, miserable looking, dejected, neurotic man." He was in almost constant pain, and frequently vomiting. The stomach was considerably dilated. On May 10th, 1895, the abdomen was opened. The pylorus, duodenum, and colon were all found matted together in a large, hard mass, the duodenum being apparently constricted in two places, and what was believed to be a large ulcer was felt in the pyloric region. Pyloroplasty and pylorotomy were out of the question, therefore gastro-enterostomy with Murphy's button was done. The patient did as well as such a neurotic type of individual could be expected to. He had plenty of pains and discomforts, but the temperature never exceeded 99.5°, and he began to get up on July 8th. On July 14th he commenced common diet. As the button was not passed he was given a very full meal on the 18th, and made to vomit afterwards, but without effect. So far as we know, the button has never been passed. I have had him examined with the x-rays on two occasions, but without a definite result. He is now well-nourished, florid, and healthy-looking; but came to see me the other day nearly three years after the operation with a slight return of the old pain. A few days' rest in bed seemed to put him all right, but I feared a slight recurrence of the ulceration. The obstruction seemed to have been quite relieved.

CASE X.—Mr. A., aged 44. Old duodenal ulcer; stenosis; gastro-duodenostomy; death from cardiac failure on seventh day. Two brothers had similar but less severe gastric symptoms. At the age of 17 had first gastric attack, which lasted three or four months. Has been subject to such attacks since at intervals. They are characterised by flatulence, pain, constipation, and vomiting of quantities of fluid. Between the attacks he is quite well, and is able to enjoy food. The symptoms have tended to gradually increase in severity. He has now flatulence, pain, frequent nausea, and occasional vomiting of large quantities of frothy fluid containing sarcinae. The abdomen is distended, there is tenderness over the epigastric region, and considerable dilatation of the stomach. On May 19th, 1895, the abdomen was opened. The pylorus was thickened, contracted, and tucked up under the liver. The second part of the duodenum lay near the pyloric end of the stomach, and an anastomosis was established between them with Murphy's button. The subsequent course of the case was unsatisfactory throughout. He began with nausea and frequent cough with mucous expectoration, which we attributed to the ether. The pulse became rapid and bounding. Temperature ranged from 99.5° to 102°. He suffered from flatulence, and frequently vomited small quantities of fluid. After a slight temporary improvement he began to fail, and died on May 26th, seven days after the operation. The patient was markedly of the neurotic type, but sensible and patient throughout. Dr. Carter of Preston, whose patient he was, made a post-mortem examination of the parts. He found the pylorus constricted to the size of a cedar pencil, and an annular ulcer in the duodenum. The button was still in position, there had been no leakage, and there was no septic peritonitis. Apparently the patient was not in a condition to bear

the operation, and had died from cardiac failure without any fault which could be directly attributed to the operation.

CASE XI.—A. C., aged 35. Gastric ulcer; pyloric stenosis; gastro-enterostomy; recovery. Patient began to suffer from symptoms of gastric ulcer six or seven years ago. Since then she has been three times in the infirmary for this complaint. The fourth attack occurred a year ago with hæmatemesis. On January 30th, 1897, she was admitted under Dr. Caton with a largely dilated stomach, and was moderately emaciated. She improved under medical treatment, and was transferred to me in March for operation. On March 11th the abdomen was opened. The parts in the pyloric region were hopelessly matted together and adherent to the underside of the liver. Pyloroplasty had to be abandoned, and gastro-enterostomy with Murphy's button was done in its place. A finger was introduced through the stomach opening to examine the pylorus; it was found thick, hard, and contracted, with a very large and thick-walled ulcer spreading into the pyloric end of the stomach. Patient made a good recovery, and was discharged well on April 24th. The button was not passed. She was seen on June 11th in very good health, eating all kinds of food with enjoyment. Her weight then was 108 lb. Before the operation it was just 8 st.

Of the three operations used in the treatment of fibrous stricture of the pylorus, division, pyloroplasty, and gastro-enterostomy, the former, as has already been said, is unsatisfactory, and must now give place to the other two. Pyloroplasty is the best operation, but unfortunately it is not applicable to all cases. When the parts are sufficiently yielding to permit a broad band of healthy mucous membrane to be drawn into the contracted pyloric orifice, the result is perfect; better I think than is ever obtained by any other stomach operation. But when the hardness and fixation of the surrounding tissues is such as only to allow the approximation of a little mucous membrane from stomach to duodenum, the result is not superior to that obtained by gastro-enterostomy, if as good, and when this can only be effected with much tension on the sutures, gastro-enterostomy is distinctly the better operation. Moreover, when active ulceration is present, it is generally undesirable to operate in its immediate neighbourhood. Pyloroplasty is probably a less severe operation than gastro-enterostomy, but one need not attach so much importance to this in vigorous subjects, since in such cases the dangers of the latter are greatly lessened, and patients with fibrous stricture are as a rule much stronger, and have much more reparative power than those suffering from malignant disease. When the patient is exhausted pyloroplasty should be done if possible.

In regard to the two deaths, Case VIII died from septic peritonitis, the result of the operation, though I was not aware of any source of contamination of the peritoneal cavity. The pyloroplasty wound could not have looked more firm and healthy than it did at the post-mortem examination, the sutures being entirely covered with firm plastic exudation. Some people are specially susceptible to septic inflammation of the peritoneum. Case VIII was one of these, for the operation was easy and there was no accident of any kind. The death following Murphy's operation (Case X) occurred in a similar neurotic type of subject, though not exactly in the same way. Vomiting commenced soon after the operation, and was accompanied later by rapid pulse and elevation of temperature, with increased flatulence, but he did not die until the seventh day, and at the post-mortem examination no evidence of septic peritonitis was found. I think there must have been some septic element in the case, perhaps connected with the slough produced by the Murphy's button. The result would probably have been different had it occurred later, when the advantages of pyloroplasty were known and appreciated.

For malignant stricture of the pylorus I should have said there were only two operations—pylorotomy and gastro-enterostomy—were it not that my colleague, Mr. Rushton Parker, has recently been trying pyloroplasty with some success in cases in which the pylorus was not entirely surrounded by growth. The value of the operation in malignant cases must remain *sub judice* pending further experience. There seem to be some *a priori* objections to it in these cases; but such objections are readily met by practical experience, and it cannot be said that the latter under similar circumstances presents gastro-enterostomy in a very favourable light. I have done pylorotomy in two cases only. The first died, the patient being almost *in extremis* at the time of the operation. The second recovered, dying sixteen months later from recurrence, after having been very strong and well for about a year.

PYLORECTOMY.

CASE XII.—A. A., aged 45. Pyloric cancer; pylorotomy; death from

shock. Unfortunately the report of this case is missing, and I have only the essential details. Patient had been suffering from increasing symptoms of pyloric obstruction for nearly a year. She was admitted into the infirmary under the care of Dr. Davidson, with frequent copious vomiting and great dilatation of the stomach. She was wasted and considerably exhausted for want of food, but, as she vomited everything, had to be kept entirely on nutrient enemata. After ten days' rectal alimentation, as the patient was rapidly losing ground, she was transferred to me for operation. On April 22nd, 1897, the abdomen was opened, and a small hard carcinoma was found in the pylorus almost completely obstructing it. The growth was freely movable, and the stomach was greatly dilated and distended with gas and dark brown fluid. I rapidly excised the pylorus, closing the wound with sutures. It was a very easy case to operate upon, the entire proceeding occupying under an hour, and the state of exhaustion did not seem to be very much increased. However, the patient showed no signs of rallying, and she quietly faded out next day. At the *post-mortem* examination the condition of the parts operated upon was found perfectly satisfactory. This would have been a very good case for pylorotomy had the operation been undertaken at an earlier stage of the complaint. It was, of course, an error in judgment to attempt it in her then condition.

CASE XIII.—A. S., aged 56. Carcinoma of pyloric end of stomach; pylorotomy (gastrostomy); recovery; death sixteen months later from recurrence. Patient was a widow with children; worked hard and supported her family. For the last six months she had been very ill with indigestion and pain, but no vomiting. Gave up all solid food and was easier. Seven weeks ago first noticed a lump in the epigastric region the size of a hen's egg, which grew steadily larger. She was sent to me by Drs. McDonald and Huskie. The patient was wasted, but healthy looking. The tumour was perfectly visible through the abdominal wall, and felt as large as a good sized orange. She had no vomiting. On May 25th, 1895, I opened the abdomen. The pyloric end of the stomach for 4 inches was the seat of a dense, thick, hard mass, more like sarcoma than the ordinary carcinoma, and a few adjacent glands were white and hard, and about the size of peas. On May 27th the affected structures were excised with scissors, and the viscera reconnected with a Murphy's button and sutures. The patient suffered very little from shock, but was not comfortable after food until June 12th, when perhaps the button became dislodged. It was not passed until June 25th, after which the patient felt quite well. She soon returned to her work, and continued in excellent health for a year, when evidence of recurrence began to manifest itself, not in the stomach, but in the glands and peritoneum. She lost flesh rather rapidly, and died at the end of six months from the date of the operation.

Pylorotomy should be reserved for favourable cases. I would not again attempt it in an exhausted patient like Case XII. A gastro-enterostomy is the most that a person in such a condition can be expected to survive. In Case XIII the limit from another aspect, namely the extent of the growth, was about reached. It is true we have before us the recently published case of complete excision of the stomach by Dr. Carl Schlatter, but one may be allowed to hope that such an example will not be taken as a practical guide in the surgical treatment of cancer of the stomach. Strongly as I would urge, when reasonable, a curative in place of a palliative operation, I think pylorotomy should only be attempted when the growth is fairly small, and above all things freely movable. In regard to the manner of doing the operation the only differences of opinion relate to the method of closing the wounds. Here the choice lies between suture alone, suture and a mechanical appliance, and complete closure of the cut ends combined with gastro-enterostomy. In some cases the latter is imperative, because the cut ends will not come together, otherwise I should not think this plan desirable. As regards the choice between simple suture and suture combined with a button or bobbin I have some doubt, but incline towards the former when approximation is fairly easy. When there is any difficulty in applying the sutures the posterior ones are in danger of proving insecure, and under such circumstances the use of a button would probably be advantageous.

GASTRO-ENTEROSTOMY FOR MALIGNANT PYLORIC OBSTRUCTION.

I have done this operation on seven occasions, and am sorry to say that the mortality has been heavy. Two were done with Senn's bone plates, 1 died and 1 recovered; 1 with bone rings, which was fatal; and the remaining four with buttons, of which 2 died and 2 recovered. Of the 4 deaths the Senn's case and 1 of the Murphy's were due to shock and previous exhaustion, 1 was due to the button, and 1 to peritonitis.

CASE XIV.—R. C., aged 60, was admitted under Dr. Caton for scirrhus pylorus. On July 13th, 1889, I did gastro-enterostomy by Senn's method. The patient did very well for a few days, but in these early cases we were too much afraid of feeding, and this patient got nothing but small sips of water by the mouth until the end of the seventh day, when it was too late. The exhaustion was too profound, and he died the next day.

CASE XV.—Mrs. B., aged 48, a patient under the care of Dr. Barnes, suffering from cancer of the pylorus. I operated by Senn's method on November 17th, 1891. She recovered and was up by the end of the month. After a few weeks of comparative comfort the symptoms of obstruction recurred, and she died little more than two months subsequent to the operation. At the necropsy we found the opening had nearly closed again.

CASE XVI.—J. G., aged 45; symptoms of pyloric obstruction for upwards of a year. On June 2nd, 1892, I did gastro-enterostomy on the posterior aspect of the stomach by a method I recommended for producing a circular punched-out opening by strangulating a portion of the viscera with two bone rings ligatured together. The effect was exactly that caused by Murphy's button, but the rings could not be used with the same ease and speed as the button. Unfortunately the patient died from peritonitis on the second day, so the operation was never fully tested in the human subject, for I of course gave it up as soon as the better appliance suggested by Dr. Murphy was introduced.

CASE XVII.—John N., aged 35. Patient was admitted under Dr. Caton in August, 1896, having suffered from indigestion and vomiting for five months. Had gastric dilatation, and was anæmic and wasted. Improved under medical treatment. On November 5th I opened the abdomen and found a carcinoma of the pylorus, which was unsuitable for excision, and therefore did Murphy's gastro-enterostomy. He recovered rather slowly, and did not pass the button, which was beautifully shown by Professor Oliver Lodge, about three inches to the right of the umbilicus. He was discharged on January 1st, 1897, much relieved. In May he had hæmatemesis, and soon after a nodule of growth was found in the abdominal scar. In June the growth had much increased, and on July 19th he died from exhaustion. The necropsy was made by Professor Boyce. The gastro-enterostomy orifice was 6 inches from the pylorus. It was completely surrounded by growth, but still admitted a finger easily. It was 3 inches distant from the pyloric growth, which was extensive, and had ulcerated into the gall bladder and liver. The abdominal scar was also infected, though not connected with the primary growth. Death was considered to be due to the liver complication, and hæmorrhage into the stomach. Professor Boyce pointed out that this was a marked case of secondary infection of scars.

CASE XVIII.—Harry C., aged 34. Symptoms of pyloric obstruction for nine months. Wasting, frequent vomiting, and a nodular lump in the pyloric region. The abdomen was opened on April 6th, 1897, and the growth proving too extensive for pylorotomy, Murphy's gastro-enterostomy was done. Patient made a good recovery, and left on May 8th, not having passed the button, but eating fish and minced meat with perfect comfort. The vomiting stopped completely. After his return home the growth increased rapidly, and he lived rather less than six months after the operation.

CASE XIX.—Mr. C., aged 50. Seen with Drs. Barron and Chisholm. This patient had an old history of indigestion, with signs of pyloric obstruction coming on for the last twelve months. When first seen in April, 1897, he was in good condition, and no tumour could be felt. Exploration was advised, but not acceded to. Subsequently he lost flesh rapidly, and a tumour in the pyloric region could be easily felt. On June 25th I did Murphy's gastro-enterostomy. The operation produced no shock, and for some days he progressed splendidly, having no pain and no elevation of temperature. On July 1st he became very weak, and later in the day something evidently gave way, as he was attacked with abdominal pain, became collapsed, and died the following day.

CASE XX.—C. D., aged 65. Had suffered severely from indigestion, vomiting, and hæmatemesis for the last seven months. When seen had a pyloric tumour and a dilated stomach. Was emaciated, anæmic, and prostrated. It was really too late for operation, but he was very anxious for it, so I consented to try gastro-enterostomy. The operation was easily and quickly accomplished on December 5th, 1897, a Murphy's button being used, but he died two and a half days later from exhaustion. The operation caused very little apparent shock.

Gastro-enterostomy when done for malignant disease is a much less satisfactory operation than when done for fibrous stricture. It has all the disadvantages, associated with a continuance of the disease, which accompany colotomy for cancer of the rectum, without the safety characteristic of the latter. The two patients that recovered from the Murphy's operation were no doubt much relieved, and if the mortality could be substantially lowered, one would, I believe, look much more favourably on this method of treatment. Gastro-enterostomy, however, must not stand or fall on the results obtained in cases of advanced malignant disease. In non-malignant cases, and in young and robust subjects of all sorts its success is already very good.

The conclusions at which I arrive regarding the four operations of which the above 20 cases constitute my personal experience, are the following: Divulsion should be reserved for very exceptional cases, other and better methods having almost completely excluded it. In the complete form of Loreta's operation with an incision into the stomach, it is not likely to be employed in the future, but one can imagine cases in which one might explore with the expectation of finding a fibrous stricture, and discovering only a dilated stomach with a small and perhaps spasmodically contracted but not otherwise strictured pylorus, one might make use of such a simple and safe method of dilating the orifice as to pass one's fingers through it by invaginating the anterior wall of the stomach. Of the other three operations, pylorotomy under present conditions is only likely to be the best method of giving relief in a small percentage of cases, and though nothing can compete with pyloroplasty in the complete success which follows its employment when circumstances are favourable, there must always remain a considerable number of cases in which gastro-enterostomy alone is available.

After all, this operation with its many failings must re-

main a most important one, because it is always practicable when the others prove impossible. We may well, therefore, turn our attention towards endeavouring to improve it. Its greatest failing has always been a high mortality, primarily due no doubt to the exhausted condition of the patient at the time of the operation, or rather to the unsuitability of the operation for the class of cases for which it is required. One surgeon will blame the mechanical appliance for causing death, whilst another, who eschews buttons and bobbins, admits the danger of prolonged suturing operations with their risk of peritoneal contamination and their want of perfect water-tight security. As a matter of fact both methods are very well in their way, and succeed very well in the right sort of patients, but they are not suitable for exhausted subjects, and, unfortunately, the sort of subjects requiring this kind of relief usually are exhausted. When the patient is robust he reaps all the advantages of the button, plate, or hobbin, and is able, owing to vigorous and sound union, to withstand their dangers; but when, as is too often the case, the patient is in an exhausted condition, the nutrition low, and the union soft and delicate, the mechanical support becomes an active source of danger. As a foreign body it lies in contact with the wound, and in the case of the button about the sixth or seventh day perforation is liable to occur, with extravasation of gastric or intestinal contents into the peritoneal cavity, and rapid collapse and death. Plain suture in gastro-enterostomy has difficulties and dangers quite as great. The time occupied in effecting a water-tight union, the greater danger of contaminating the peritoneum during the operation, and the subsequent tendency which the opening shows to close are substantial objections to this mode of operating. At present one has to choose between the rapidity and primary security due to the ease and perfect coaptation of button operations, and the slowness and less perfect primary security of suture operations without a mechanical appliance.

Neither, unfortunately, are favourable methods to employ upon exhausted subjects. In fact, gastro-enterostomy still presents an open field for surgical ingenuity, as the objections referred to apply equally to the latest operations, such as those suggested by M. Faure and M. Desfosses. I make, therefore, no apology for proposing yet another plan. If it does not altogether hit the mark, it may be that a useful application for the principle will be found elsewhere, or that it will stimulate other and better proposals.

Anterior gastro-enterostomy by a new method. 1 The eschar produced by chloride of zinc. 2 Wound in the stomach made by excising a portion of the outer coats. 3 Wound in the intestine made by incising and rolling back the outer coats. The commencement of the continuous suture is shown.

The plan is as follows: The bowel and stomach having been exposed, a longitudinal incision is made in the form—
—for about 2 inches in the human subject—through the

peritoneal and muscular coats only. These are reflected with a small, curved, blunt dissector over an oval area, having a diameter of about 1 inch in the centre. The exposed submucosa is then rubbed with a stick of zinc chloride, say half a dozen times, being well dried between each application, when it assumes a grey dead look. The bowel is now wrapped in a piece of moist absorbent wool and laid on one side whilst the stomach is dealt with in an almost similar way. In preparing the stomach for the anastomosis, instead of making a linear incision, an oval patch of musculo-peritoneal coat is excised; the muscular coat being so much thicker in this organ, it is rather in the way if only reflected, and what is of more importance is that the loss of substance helps to prevent subsequent contraction of the opening. Both the wounds having been duly cauterised, they are brought together, and a continuous suture of the finest green gut or silk is run round the edges, as shown in the illustration—not, of course, penetrating the mucous membrane, but picking up as far as possible the tough submucous coat. Finally, a few Lembert or Halsted sutures are generally desirable, especially in anterior gastro-enterostomy.

The result of this operation in dogs—it has not been done in the human subject—is that the mucous and submucous coats slough and entirely disappear between twenty-four and forty-eight hours after the operation, and good firm union seems to take place with great rapidity. The complete removal of the slough might be a somewhat slower process in the average human patient; but probably an anastomosis would be effected within forty-eight hours—which is practically as soon as food could be digested, and quite early enough having regard to the proper adhesion of the external coats.

The features of this operation which it is hoped may be reckoned as advantages are:

1. The effect is one of pure traumatism. The viscera not being opened, all risk from such sources is avoided.
2. No foreign body is used.
3. The time occupied is less than for a suture operation, though more than for Murphy's.
4. The anastomosis resulting from loss of tissue should be more durable than in the majority of methods.

In performing the operation experimentally upon animals I have had the kind and valuable assistance of Professor Sherrington, who has personally helped me on each occasion, for which I feel very much indebted to him.

The chief points to which our attention has been directed were to ascertain:

1. The best chemical for causing necrosis of the mucous and submucous coats.
2. The interval between the operation and the breaking-down of the slough.
3. The character of the union.
4. The efficiency and durability of the anastomosis.

Eight animals were operated on, and the experiments have been continuously progressing from November 2nd, to the present time.

In regard to the general result the most noticeable feature has been the complete absence of shock in every case in which gastro-enterostomy alone was done. In estimating the severity of the operation, I am able to compare the effect of this method with that of several previous gastro-enterostomies in dogs in which the viscera were opened, and I attribute the absence of shock, and, indeed, of any appreciable pain or disturbance, to the fact that nothing of a dangerous nature was done. The viscera were not opened, and therefore the local peritoneal reaction was purely plastic and aseptic, and was not accompanied by that sensation of pain and insecurity which is present when the localised peritonitis is associated with septic as well as traumatic influences. Professor Sherrington has had a much larger experience of the effect of operations upon animals than I have, and he quite endorses this view.

Only three chemicals were tried, as in the first instance I was limited to five experiments, though the limit was subsequently withdrawn. Strong nitric acid well applied failed to produce an anastomosis. At the end of a fortnight firm adhesions were present over the area operated on, but no opening at all. Chromic acid was more successful, but I

thought not quite so efficient as zinc chloride, which has now been used seven times on six animals without once failing. Several other escharotics were thought of, but as the zinc chloride seemed to answer so well they were not tested.

The interval between the operation and the separation of the slough has not been quite definitely fixed. At 16 hours it was on the point of giving way. At 48 hours it was entirely removed, and strong union was present all round the opening. There were no other cases bearing upon this point.

The character of the union was very good in all the cases. The continuous suture was thoroughly embedded in a firm adhesion of plastic lymph which left nothing to be desired on the score of safety.

It has been difficult to decide as to the permanent efficiency of the anastomosis in dogs. In one the duodenum was divided, and gastro-enterostomy performed at the same sitting. The animal died at the end of two days from gangrene of the distal end of the duodenum, due presumably to division of the gastro-duodenal artery. Another in which the duodenum was greatly narrowed thrived well, but it seemed probable that some of the gastric contents continued to pass by the natural way, though the constriction of the bowel was sufficient to have produced dilatation above it. In a third, contraction of the pylorus was effected by destruction of the mucous membrane with zinc chloride, gastro-enterostomy being delayed until symptoms of pyloric obstruction were marked. After a few days' relief the severity of the symptoms increased, and the operation was a failure. At death a fortnight later the anastomosis was present, and would admit a little finger. It had either been made too late, or too small, or in a bad position, for it was high up near the lesser curvature towards the cardiac end. Probably all three conditions contributed to bring about failure. In this case the external coats of the stomach were incised only, and not cut out as I would now recommend, and the original incision itself was too small. Probably to get an orifice half an inch in diameter after complete recovery has taken place, one must operate on an area of twice that extent, and the loss of tissue should involve all the coats of the stomach if subsequent contraction is to be safely guarded against.

There was no evidence of regurgitation of intestinal contents into the stomach in any case. All the gastro-enterostomies were anterior, and in my limited experience in dogs and men this has not occurred in the anterior operation, though it did in one or two cases of the posterior operation in dogs which I performed in 1892.

In conclusion, I may say that I feel satisfied as to the safety of the operation I have proposed, and I believe there is a fair prospect that it will prove successful in effecting a satisfactory lateral anastomosis. The result of Murphy's operation in exhausted patients is sufficiently discouraging, in my judgment, to exclude it in this class of cases, though it is very successful under more favourable circumstances. I shall therefore feel justified in employing this method for my next gastro-enterostomy.

REFERENCE.

¹ *Liverpool Medico-Chirurgical Journal*, No. 1, p. 160, On Some of the Most Remote Effects of Syphilis.

THE OCCURRENCE OF MENSTRUAL SECRETION IN THE FALLOPIAN TUBES IN THE HUMAN SUBJECT AND ITS SIGNIFICANCE.

By C. J. BOND, F.R.C.S.,
Surgeon to the Leicester Infirmary.

I have published in the *Journal of Physiology* for February, 1898, a preliminary note on certain observations bearing on the normal and pathological secretions of the uterus and Fallopian tubes in the human female and in some of the mammalia. In the present communication I wish to draw attention to the subject as far only as it affects the human subject, and especially to the occurrence in some cases of a menstrual secretion in the Fallopian tubes.

Three years ago, on cutting across a Fallopian tube removed with the ovaries for uterine disease on the first day of the menstrual flow, I was struck by the fact that the tube contained a quantity of blood-stained mucus resembling

menstrual fluid. Further observations were then carried out on cases operated on during the menstrual periods. Immediately after removal the tube was cut across near the uterine end, and the contained fluid, which exuded on gentle pressure from the cut end, examined both in the fresh state, and by means of dried films stained with logwood and eosin.

Fig. 1.—Film preparation of normal menstrual fluid from the external os on the second day of menstruation. Stained logwood and eosin. $\frac{1}{2}$ obj. A. Leucocytes. B. Red blood corpuscles. C. Altered epithelial cells.

Portions of the tubes immediately after removal were also hardened in formol 5 or 10 per cent. solution, and sections cut and stained with the same reagents.

Fig. 2.—Film preparation of fluid from Fallopian tube on third day of menstruation. $\frac{1}{2}$ obj. Stained logwood and eosin. A. Leucocytes. B. Red corpuscles. C. Epithelial cells.

It is, of course, necessary to exclude all cases in which at the time of operation the tubes are evidently diseased, and in which there had been peritonitis with obliteration of the infundibular ostium and distension with either blood, serum, or pus. Consequently the observations have been limited to those cases in which the tubes appeared to be, and indeed were, healthy, and on which the operation was performed for other conditions, such as fibroids or cystic or diseased ovaries, etc.

In four such cases, in which the tubes were removed and