the right lumbar region were increased, and a definite rounded swelling could be made out. On the same day, in the evening, 10 c.cm. of serum were injected under the skin to the front of the swelling. The urine was examined on February 25th, and contained a quantity of pus and some albumen.

On February 26th in the morning her condition was much the same, except that the urine which had been passed during the night was much clearer, and contained only a small quantity of pus. In the evening the temperature had fallen to 102° and the pulse to 126. The urine passed during the day was quite clear, and contained a little albumen but no pus. Pain and tenderness in the right lumbar region had almost disappeared. Ten more c.cm. of serum were injected in the right lumbar region, but closer to the spine. This injection, for which some dried serum was used from the Pasteur Institute, was followed by some erythema and tenderness of the skin around the spot, which gradually disappeared.

On February 27th the improvement was most marked, the temperature was only 101.2°, and the pulse 126. The tongue was moist and red at the edges, with slight fur down the centre. Pain and tenderness had quite disappeared. The urine still remained clear, and the albumen had diminished.

The next day there was a still greater fall in temperature and pulse. In the evening the temperature was normal. The pulse was still rather rapid, but this was probably due to the stimulant, which was ordered to be left off.

The patient continued to improve, and on March 3rd the pulse had fallen to 80, and she was practically well. The child was living. The urine still remained clear, all tenderness and pain had disappeared.

The serum was supplied by Messrs. Burroughs and Wellcome.

A CASE OF HÆMORRHAGIC INFARCTION OF THE SMALL INTESTINE: SUCCESSFUL RESECTION.*

By T. E. GORDON, M.B., F.R.C.S.I.,
Surgeon to the Adelaide Hospital, Dublin.

THE history of the following case seems to me to possess sufficient clinical and pathological interest to justify my bringing it before you:

The patient is a woman about 45 years of age. She gave me a tolerably clear history of secondary syphilis contracted some eight or ten years ago, and she has now undoubted evidence of tertiary syphilitic ulceration of the hard and soft palate.

In November, 1896, she was operated on by Mr. Scott for strangulated femoral hernia on the right side, and she made a complete recovery. Her health continued good until the day before her readmission.

On the morning of that day—that is, August 17th—she was feeling as well as usual, and her bowels were moved. While she was having her lunch at 12 o'clock she got a sudden and most severe gripping pain in the lower part of the abdomen. She was carried to bed, and Dr. Woods, of Amiens Street, was sent for, who gave her morphine when he arrived. At about 5 o'clock on the morning of the 18th she commenced to vomit, and this symptom persisted until the time of admission. The bowels were not moved subsequent to the attack of pain. She was sent at once to the Adelaide Hospital, and placed under my care in Mr. Scott's absence.

When I saw her, she appeared to be very ill but not collapsed. The abdomen was not distended, and was sufficiently flaccid to enable me to easily detect a tumour in the right iliac region, apparently formed by a distended coil of intestine. Dr. Peacocke saw the patient with me, and both he and I thought the case one for immediate operation. We supposed that there existed some form of mechanical obstruction. I accordingly operated, being assisted by Dr. Smyly.

On opening the abdomen in the middle line below the umbilicus, a considerable amount of blood-stained fluid escaped and a distended coil of very dark bowel presented in the wound. The peritoneum had not lost its polish, and, indeed, there were none of the unequivocal signs of gangrene present. There seemed to be a constriction at the upper end of the loop, and this marked off the darkened region very abruptly; below the transition to normal intestine was more gradual. The wall felt thickened, and emitted a heavy but not offensive odour. In the mesentery I found a continuation over a limited area of the same condition. I decided to resect and establish end-to-end anastomosis. I used for the latter purpose Mr. Ball's bobbin, of the smallest size I could obtain, and fixed the ends of the intestine into the groove with purse-string sutures. Over this I applied a continuous subserous suture of fine silk. For reasons already indicated I removed a considerable area of mesentery. Finally, I sponged out any fluid in the neighbourhood and closed the abdominal wound.

The amount of bowel removed measured 2 feet. It was found to contain a mass of foul-smelling blood clot. The wall was much thickened, and the constriction at the upper end proved more apparent than real. There was no stricture and no ulceration of the mucous membrane.

* Read at the Surgical Section, Royal Academy of Medicine in Ireland.

I did not discover any definite cause for the condition found. It is worth noting that some vessels in the mesentery bled freely as they were divided, and one of considerable size was cut, which may have been, in part, responsible for a complication which I must now refer to. On the third day after the operation, at 6.30 A.M., the patient complained of the most intense pain in the abdomen. The house-surgeon gave one-sixth of a grain of morphine, which relieved her, and when I arrived she did not appear to be in any great degree of danger. With the attack of pain a slight fall of temperature occurred, and the pulse was 128. By the 22nd—the fifth day since operation—the temperature reached normal, and the course of the case, until the 27th, seemed to promise an uninterrupted recovery. On that day I found, on removing the dressings, that a considerable leakage of intestinal fluid had taken place. I opened up the wound to some extent, and packed with iodoform gauze. A couple of days later (the 29th) a slough came away which seemed to represent the entire line of union, and contained a considerable length of both sutures, superficial and deep. Up to this time the bowels had not been moved satisfactorily, but there was clearly no obstruction, nor was there evidence of other than a purely local peritonitis. On the 30th I ordered a castor oil enema, which acted well. The discharge of bile-stained fluid was now so great as to necessitate very frequent change of dressings, on two occasions nine times in the twenty-four hours. On September 5th the patient passed a natural motion, and the discharge from the wound was reduced to very little, and had ceased entirely on the following day. It recurred to a very slight extent a few days later, a mere soiling of the plugging, before it finally ceased, that is, ten days after the discharge was first observed, and about three weeks from the date of operation. The subsequent history has been uneventful, and the patient now appears to be in perfect health.

REMARKS.

It is scarcely possible to form any definite opinion concerning the cause of sloughing at the suture line. The possible influence of the division of the mesentery I have already alluded to. It may, however, have been due to an extension of the pre-existing local condition, and the accident might have been avoided by a more liberal resection. Or again, the flange of the bobbin may have exerted undue pressure on the wall of the gut as it passed over it. This would have been avoided by using a smaller tube.

With some hesitation I have arrived at the conclusion that this case was one of hæmorrhagic infarction of the intestine dependent upon a sudden obstruction of some branch of the superior mesenteric vessels. The condition might conceivably have been due to:

1. Inflammation. This we can certainly exclude, for it is impossible that such a violently acute inflammation could have remained so distinctly local. The onset, too, was more sudden than could be explained on this hypothesis; and, finally, the microscopic appearances were not those of inflammation.

2. Some of the usual causes of internal obstruction. None of these were to be seen at the operation, and the appearance of the involved segment of gut was not such as one would expect to find if the cause had been, let us say, a peritoneal band. Had such been present, though overlooked, surely the delimitation of the area of strangulation would have been as marked at the one extremity as the other, and this, as we have seen, was certainly not the case. There were no evidences of previous inflammation about the hernial orifices.

3. If it is allowed that none of these extrinsic causes were at work, it follows that the cause of obstructed circulation must be looked for within the vessels themselves; must, in fact, have been either embolism or thrombosis.

The possibility of the occurrence of an infarction as the result of occlusion of the superior mesenteric vessels seems at first sight very difficult to admit. One is accustomed to find such a lesion in areas supplied by terminal arteries only. Experiments, however, as well as numerous recorded cases of intestinal infarct, place the possibility beyond doubt. The subject is dealt with in *Ziegler's Pathology* as well as by Cohnheim in a most important and convincing essay, by Coats and others. I will quote a few passages of special interest from the last-named pathologist. He writes: "It has been said that the hæmorrhagic infarction forms in the case of end arteries, but there is an apparent exception to this in the case of the superior mesenteric artery. If the superior mesenteric artery or one of its larger branches be obstructed, hæmorrhage occurs in the intestine and mesentery, and the portion of gut suffers necrosis, but it is well known that the anastomosis of the mesenteric arteries is tolerably free. The explanation of this apparent anomaly has been worked out by Cohnheim and his pupils in a very interesting manner. As a rule, it may be said that main stems have, in proportion to their size, much less considerable anastomotic communications than the smaller branches. If the main stem be obstructed, therefore, the restoration of the circulation occurs

much more slowly than when a small stem is the seat of obstruction. In the case of the superior mesenteric and its larger branches, obstruction is slowly recovered from, and for a time the circulation is virtually at a standstill."

"But here a very important element comes into consideration. The vessels of all parts of the body are not equally resistant. Those of the skin and muscles are peculiarly resistant, and are able to stand a considerably prolonged deprivation of blood. It is different with those of the intestines." The author then proceeds to describe an experiment made to test the vulnerability of the intestinal vessels. The conclusion drawn from this is that "before the anastomosis can be established the backward flow from the veins has already begun, and even the diapedesis. The re-establishment of the circulation even increases the tendency to hæmorrhage, as it subjects the blood to a higher pressure than that of venous blood. The experiments referred to showed that when the superior mesenteric artery is obstructed the hæmorrhagic infarction forms with a backward flow from the veins just as in other cases."

I must allow that it is more difficult to explain an infarction limited to a single loop of intestine than the occurrence of a more extensive lesion, for it seems necessary to assume that the obstruction which produces the less effect is situated in a part of the superior mesenteric distribution where the collateral circulation is extremely free. It is possible, however, that this may be explained by supposing that when one vessel had become occluded a rapid extension of thrombotic occlusion followed along the alternative blood circuits. This would be less difficult to assume possible if it could be shown that the vessels of the area were atheromatous or otherwise abnormal. I shall have something to say with reference to this point presently.

To conclude this part of my subject, I may add that the appearances in my case, both macroscopic and microscopic, were such as one would expect to meet with in a case of sudden vascular occlusion. Thus there was considerable hæmorrhage into the peritoneal cavity, the coil itself was full of foul smelling blood clots, and microscopic examination showed that the wall of the intestine was everywhere occupied by extravasated blood.

Although cases of hæmorrhagic infarction of the intestine are uncommon, I have found records of a considerable number. Osler, for instance, has recorded 3 which have occurred in his own practice. Most of the cases have been, as one would expect, associated with heart disease, but different causes have operated in others. Ulceration of the duodenum appeared to be the primary lesion in one instance, and aneurysm of the superior mesenteric artery caused infarction in one of Osler's cases. In the present case two important questions naturally suggest themselves. Could the lesion be in any way traced to the syphilis, and did the strangulated hernia take any part in its causation? Syphilitic arteritis is a well-recognised pathological condition, and although most writers on the subject deal almost exclusively with it as it affects the cerebral vessels, there can, I think, be little doubt that any artery may show similar changes. This is rendered probable by *a priori* reasoning, and more than probable by the record of numerous cases of endarteritis in vessels other than cerebral in syphilitic subjects. Instances of its occurrence in those whose age is much below that at which arterial disease from other causes usually takes place are of special importance. There is then nothing improbable in the supposition that this case may have been one dependent on arterial thrombosis in syphilitic vessels. Moreover, cases have been reported of mesenteric thrombosis stated to be of a syphilitic nature.

As regards the second question, it is worth noting that the damaged coil lay in the right iliac region, and the previous strangulation had been on the same side. It is, too, of interest in this connection to recall to mind Mr. Hutchinson's statement that "in almost all" (tertiary affections) "we see good reason to believe that in addition to the syphilitic taint some localising influence takes an important share in evoking the local changes." Is it not just possible that the strangulation may have acted in this instance as such an influence, and determined the occurrence of the syphilitic arteritis in this particular region?

The extent of intestine involved may range from a few

inches to almost the entire length of the small intestine, as in a case recorded by Dr. Frederick Taylor; and I have seen a report of one case in which three distinct infarctions had taken place.

The disease is very generally fatal, but, it is supposed, not necessarily so in all cases. A case of Gull's is quoted in proof of this. In this patient copious hæmorrhage at the outset, with subsequent passage *per rectum* of long sloughs of mucous membrane, together with a preceding œdema of the arm of supposed syphilitic origin, were the signs relied on in making the diagnosis.

The symptoms are, as one would expect, very variable. Hæmorrhage from the bowels, followed by signs of intestinal obstruction, might in some cases lead one to suspect an infarction if there was some possible cause for embolism or thrombosis present. It will have been noted that there was no blood passed in this case of mine.

Little need be said with reference to treatment. Any case presenting such symptoms as these would be looked upon by most medical men as one urgently requiring operative interference, and it would then depend upon the extent of bowel implicated whether resection could be carried out or not. An example of successful resection for this condition is recorded in the *Annals of Surgery* for 1895.

NOTE ON A CASE OF PERFORATED GASTRIC ULCER.

By G. REINHARDT ANDERSON, F.R.C.S.,
Hon. Medical Officer, Southport Infirmary.

I WAS consulted on February 18th, 1897, by a young woman, apparently in good health, complaining of pain after food. The pain was not constant, nor accompanied by vomiting. She had vomited after every meal on February 16th, but this she attributed to a bilious attack. She had never vomited blood. The pain, she stated, had existed off and on for some months. I subsequently ascertained that twelve months previously, in the early part of 1896, she was a ward maid in the infirmary, but had to give up her situation because of anæmia and persistent dyspepsia. I found her somewhat anæmic, with pain and local tenderness in the epigastrium. I prescribed a milk diet and the usual remedies.

On February 21st, at 5 P.M., I received a call to see her as it was thought she was dying. I found the patient in a state of profound collapse, profuse perspiration, ashen face, and with a rapid, thready pulse. She complained of severe abdominal pain. There had been no vomiting.

It seems that after a very hearty meal at 1 P.M., consisting of calf's heart, peas, potatoes, and pastry, she had taken a walk of over three miles. When nearing home, at 4.45 P.M., she was suddenly seized with such severe pain in the abdomen that it became necessary to put her in a cab to get her there, though only three or four hundred yards away.

From this history I suspected perforation of a gastric ulcer, and on examining the abdomen I made out the following facts: The walls were rigid, hardly moving at all with respiration; there was marked tympanitic resonance with absence of the liver dulness to the upper border of the seventh rib, and there was shifting dulness in either flank to a small extent. Palpation elicited general acute tenderness. A small injection of morphine was given, and arrangements made for her immediate removal to the infirmary over a mile distant.

At 7.30 P.M., 2½ hours after the onset of pain, I operated. The abdomen was opened by a mesial incision 5½ inches in length from just below the xiphoid cartilage to within an inch of the umbilicus. On entering the peritoneal cavity there was an escape of gas and turbid serum with a marked gastric odour. Flakes of lymph were already visible on the surface of the liver and omentum as they presented in the wound. On drawing down the stomach, a small punched-out perforation readily came into view, from which the dark-brown fluid contents of the stomach escaped in plenty, unavoidably at first rushing into the peritoneal cavity. Traction on the stomach induced retching, increasing the flow. The perforation was now surrounded by sponges, and the stomach emptied as much as possible into a porringer. I then secured