The coexistence of diphtheria of the fauces, nasal or buccal mucous membrane, skin, vulva, or elsewhere, would suggest the nature of the eye mischief; but, even then, proof as to the identity of the two affections would be lacking in the absence of a bacteriological examination. The occurrence, within two to five weeks after inflammation had subsided, of loss of knee-jerks or of pareses in various parts of the body would be almost conclusive² with regard to the nature of the antecedent eye ailment. Such cases have been recently recorded in England by Mr. Silcock,³ Dr. Collins,⁴ and others. But these signs, valuable though they be, come too late to be of any service from the therapeutic point of view, which is all-important now that we possess antitoxin. In diphtheria of the fauces or larynx, as everybody knows, the urine is often found to contain albumen from an early period of the disease, and this sign would be helpful if it chanced to coincide with a membranous conjunctivitis, as in cases lately reported by Dr. W. J. Collins⁴ and by Mr. Percy Flemming.⁵ Albumen, however, has not been present in any of my patients. Indeed, in this particular form of disease, as in wound diphtheria, the existence of albumen is possibly the exception rather than the rule. Still, it is always advisable to test the urine of any patient who presents membrane upon an inflamed conjunctiva.

One must recollect that the mere presence upon the palpebral conjunctiva of a false membrane is not conclusive as to the nature of the inflammation. As a matter of fact, a fibrinous exudation may take its appearance in almost any form of conjunctivitis, supposing the inflammation to be severe enough to give rise to a coagulation necrosis. For example, it is not unknown in the purulent ophthalmia of the newly-born, as pointed out by Chaisaignac⁶ many years ago. It forms an almost constant sign of the inflammation set up by pneumococci; it is sometimes seen in that form of conjunctivitis which is associated with Weeks's bacilli; while I have observed it along with pus cocci, as in a case recorded by me in the *Lancet* of November 13th, 1897. Apart from these acute conditions, there is a curious chronic affection, where a thick greyish, fibrinous exudation is seen upon the palpebral conjunctiva for weeks or months together. The malady does not seem to be contagious, generally occurs in delicate children, and sometimes implicates the cornea. Instances have been reported in this country by Hulme,⁷ Mason,⁸ Hogg,⁹ Nettleship,¹⁰ Critchett,¹¹ Juler,¹² Bronner,¹³ and Stanford Morton.¹⁴ Then, there is the fact that false membrane may follow injuries to the conjunctiva, whether mechanical or chemical. Lastly, cases have been published by Sourdille,¹⁵ Aubineau,¹⁶ and by H. Coppez,¹⁷ where Klebs-Loeffler bacilli were present, but membrane was absent. Hence there is nothing in the least pathognomonic about the presence (or absence) of these fibrinous conjunctival deposits. Neither can any conclusion be safely drawn from the naked-eye characters of the film, which in diphtheria is sometimes but lightly adherent to the mucous membrane, whereas at other times it can be peeled away with difficulty or not at all. Any attempt to base a diagnosis upon the state of the underlying conjunctiva when the membrane has been stripped off appears to me fallacious. At all events it could only be in the most pronounced cases, where the subconjunctival tissues were densely infiltrated ("brawny"), that a conclusion might, with tolerable certainty, be reached as to the nature of the diseased process. The existence of small, dusky-red hæmorrhages, scattered over the surface of the diphtheritic plaques, has been a notable feature in some of my cases, although not observed in the present ones. For my part, however, I should hesitate to lay any great stress upon the point.

It is now well known that the preauricular glands (between which and the conjunctiva there is a direct connection) are liable to become involved in various forms of conjunctivitis. Some few years ago I¹⁸ described a case of unilateral conjunctival diphtheria, in which the corresponding preauricular gland was swollen and markedly tender. But further experience has convinced me that no dependence is to be placed upon this sign, since the glands in question sympathise more or less with many of the severer conjunctival inflammations, especially when due to such organisms as tubercle bacilli and gonococci. For that matter, it was not observed in the two cases that form the text of the present communication. A similar remark applies to implication of other glands, as those

about the angle of the jaw, which may or may not be enlarged.

In short, as it seems to me, the diagnosis of this ailment must be grounded upon one feature only—the presence of *diphtheria bacilli*. If these are found there can be no reasonable doubt as to the nature of the case, while their absence, after repeated and careful examination, tends to show that one is dealing with some other malady. In the latter event, a conclusion must be based on (a) the organisms demonstrated in membrane or discharges from the eye; (b) the general features of the inflammation; (c) the history of the case. In conjunctival diphtheria, then, as in diphtheria of the fauces and elsewhere, the discovery of Klebs-Loeffler bacilli must be regarded as the only certain, trustworthy, and scientific way of diagnosing the disorder, a fact that I have emphasised elsewhere.¹⁹

It is hardly necessary to make any remarks as to the method of identifying the bacillus in question, since that is so well understood at the present time. I will merely say that in conjunctival diphtheria the making of cultivations should not be employed to the exclusion of simpler means. Morax²⁰ has pointed out (and I am in a position to confirm his statement) that in a certain number of cases the discharge from the eye contains only that microbe which is the cause of the inflammation. Hence, in marked instances, examination of film preparations has brought to light such numerous and characteristic Klebs-Loeffler bacilli that he has been able at once to affirm the nature of the disease. There is one other point that should certainly be alluded to—namely, the confusion likely to arise between Klebs-Loeffler bacilli on the one hand and xerosis bacilli on the other. The latter are to be found in very many cases of conjunctivitis, and so marked is their resemblance to diphtheria bacilli that a mistake may be easily made if the investigation be not conducted with an eye to the fallacy.

Briefly, the main points of distinction are these:

1. Both stain by Gram's method, but in my experience the diphtheria organism loses its gentian violet, when in alcohol, much more quickly than the xerosis bacillus.
 2. Klebs-Loeffler bacilli, according to Dr. Eyre,²¹ give rise to an acid reaction when grown in neutral bouillon or milk, while xerosis bacilli never do.
 3. The xerosis organism when inoculated into guinea-pigs, causes nothing more than a swelling at the site of puncture.
- The last test (3) cannot, of course, be carried out except under licence from the Secretary of State.

NOTES AND REFERENCES.

- ¹ Fuchs, *Textbook of Ophthalmology*, 1894, p. 84. ² A similar form of paralysis may follow measles, typhus, and typhoid, so that it cannot be taken as wholly conclusive of a preceding diphtheria. At the same time, it may occur as a sequel to the mildest attacks of diphtheria, and is much commoner after that than after other zymotic disorders. ³ Silcock, *St. Mary's Hospital Gazette*, December, 1896. ⁴ Collins, *Lancet*, June 27th, 1896. ⁵ Flemming, *Lancet*, October 2nd, 1897. ⁶ Chaisaignac, *Ann. d'Oculist.*, 1847. ⁷ Hulme, *Med. Times and Gazette*, October 31st, 1863. ⁸ Mason, *Oph. Hospital Reports*, 1871. ⁹ Hogg, *Lancet*, March 1st, 1873. ¹⁰ Nettleship, *St. Thomas's Hospital Reports*, 1880. ¹¹ Critchett, *Trans. Ophth. Society*, vol. iii, p. 1. ¹² Juler, *ibid.* ¹³ Bronner, *Trans. Ophth. Society*, vol. xiii, p. 26. ¹⁴ Morton, *Trans. Ophth. Society*, p. 29. ¹⁵ Sourdille, *Rev. des Mal. de l'Enfance*, 1895, p. 68. ¹⁶ Aubineau, *Essai sur l'Applic. de la Sérothérapie dans le Trait de la Diphthérie Conjunctivale*, Paris, 1895. ¹⁷ Coppez, *Rev. Gén. d'Ophthalmologie*, 1896, p. 51. ¹⁸ Stephenson, *Medical Press and Circular*, March 23rd, 1892. ¹⁹ Stephenson, *Epidemic Ophthalmia*, 1895, p. 12 (Young J. Pentland, London and Edinburgh). ²⁰ Morax, *Ann. d'Oculist.*, April, 1895. ²¹ Eyre, *Journal of Pathology and Bacteriology*, July, 1896, p. 54.

SYMPATHETIC OPHTHALMIA.*

By CECIL E. SHAW, M.A., M.D., M.Ch.,

Assistant Surgeon to the Belfast Ophthalmic Hospital; Ophthalmic Surgeon to the Ulster Hospital for Children and Women, etc.
(From the Pathological Laboratory, Queen's College, Belfast.)

SYMPATHETIC ophthalmia is to be sharply defined from sympathetic irritation, the latter being recognised by functional changes only in the second or sympathising eye, while sympathetic ophthalmia is always characterised by visible organic changes of the nature of irido-cyclitis. Sympathetic ophthalmia never comes on in less than a fortnight from the time

* The expenses of this research have been in part defrayed by a grant from the Scientific Grants Committee of the British Medical Association.

of the injury to the exciting eye, and seldom so soon; and even if the injured eye is enucleated, sympathetic ophthalmia may appear in the other at any time within a month of enucleation, its severity and duration being in no way affected, apparently, by the fact that the exciting eye is removed.

Up till the last decade, the generally accepted theory was that sympathetic ophthalmia was due to nervous sympathy of one eye with the other, the irritation being conveyed by the ciliary nerves. In 1882, Deutschmann¹ published a research which has become a classic on the subject, and over which a very vigorous controversy has been carried on. He stated that after infecting one eye of a rabbit with a small quantity of a suspension of staphylococcus albus or aureus, he could detect the micro-organisms in both optic nerves and in the chiasma, and in some cases in the posterior half of the second eye; and he assumed that if the rabbits had lived long enough they would have developed sympathetic ophthalmia in the second eye, though, as a matter of fact, they all died of general infection at various periods, depending on the dose of the infecting material which they had received.

Deutschmann further said that in examining eyes removed to avoid sympathetic ophthalmia, he rarely failed to find micro-organisms.

Deutschmann's theory was at once widely accepted, for it explained (1) why sympathetic ophthalmia exceedingly seldom follows any but perforating wounds of the globe, an entry being thus afforded to micro-organisms. (2) Why sympathetic ophthalmia does not start at once, but only after several weeks, this time being needed for the germs to travel up the one optic nerve, through the chiasma, and down the other, to the second eye, there to give rise to the inflammation. (3) Why enucleation of the wounded eye does not always prevent sympathetic ophthalmia—the germs may already have started, and be on their way up the nerve or in the chiasma, out of reach of the surgeon.

Deutschmann's experiments have been repeated many times, on rabbits, dogs, and guinea-pigs, but never, I believe, with success. Gifford,² Randolph,³ Mazza,⁴ Limbourg and Levy,⁵ and Bach,⁶ have all experimented with negative results. Then, in the examination of injured eyes enucleated for fear of, or on account of, sympathetic ophthalmia, confirmatory results have been almost totally wanting. Randolph³ examined 20 eyes, and only found microbes once, in an eye enucleated a fortnight after injury. Ohlmann⁶ examined 30 eyes, and never found micro-organisms at all. Baquis⁷ examined 10 eyes with negative results. Bach⁶ examined 16 eyes, 4 of which had already caused sympathetic ophthalmia in the other, and his results, too, were negative. My own results I shall mention later. Deutschmann has so far modified his earlier views as to say that the micro-organisms are small and not easily stained.

In support of Deutschmann's results, Pincus⁸ records two cases in which he found numerous micro-organisms, one being specially interesting, as it was an old case of cysticercus, which only developed sympathetic ophthalmia after the affected eye had been operated on, and so, possibly, an entrance for micro-organisms afforded. Both in the scar and in the posterior part of the eye, and especially under Tenon's capsule and in the sheath of the optic nerve, he found numerous micro-organisms, chiefly bacilli. The other was an ordinary post-traumatic case, and showed the same appearances microscopically, with the ordinary methods of staining.

Numerous difficulties stand in the way of our full acceptance of Deutschmann's theory, or the migratory theory, as it is called. Two were obvious from its introduction. First, Why do the germs not produce meningitis? and secondly, Why is panophthalmitis not followed by sympathetic ophthalmia? To the first objection Deutschmann answered that the cocci lost in virulence as they ascended the nerve from the wounded eye against the lymph stream, and that when they reached the chiasma they were at once swept away by the lymph stream down the other nerve, to produce sympathetic ophthalmia in the other eye. This, however, is a very weak explanation, for even granting that it was so in some cases, one would expect the infection to be so severe in others that at least a slight meningitis would be produced, though this is never so. As regards panophthalmitis, which

exceedingly seldom causes sympathetic ophthalmia, Deutschmann supposed that the intensity of the inflammatory process destroyed the micro-organisms; and Gifford suggested that the channels of communication with the other eye were blocked by inflammatory products.

But cocci have been found in eyes suffering from panophthalmitis by Schirmer⁹ three weeks, by Schmidt-Rimpler¹⁰ four weeks, and by Axenfeld¹¹ five weeks after the onset of the inflammation, and have even been detected in the optic nerve. It has lately been shown by Andogsky,¹² of St. Petersburg, that if a culture of cocci is injected into the vitreous a mass of leucocytes "march out" from the nerve and protect the disc, but if the cocci are injected into the posterior part of the vitreous they succeed in entering the nerve. This tells against the migratory theory, for it is not often in wounds of the vitreous that we find sympathetic ophthalmia, but after wounds of the ciliary region. Further difficulties in the way of the migratory theory have risen as the subject has been further studied. Among these is the occurrence of sympathetic ophthalmia in spite of optic neurectomy, as in cases recorded by Rohmer¹³ (three weeks after), Troussseau¹⁴ (thirteen weeks after), and Schmidt-Rimpler¹⁴ (eighteen months after). To explain these Deutschmann made a series of experiments on rabbits, removing pieces of the optic nerve, and after varying times killing the rabbits, and injecting the sheath of the optic nerve from the meninges with Indian ink. He came to the conclusion that a fibrous connection was formed between the nerve ends, and that through this the injected fluid, and so presumably cocci, could pass; but his experiments were most carefully repeated by Velhagen¹⁵, and later by Bach⁶, who entirely failed to establish his results, and indeed showed plainly that, after a period of from two to five weeks, the injected fluid could not pass the cicatrised end of the nerve.

Another difficulty is the occasional occurrence of sympathetic ophthalmia in the absence of a wound of the other eye. Many cases have been reported, but most are probably cases of the same affection occurring in both eyes, one being later than the other, and both due to a common cause. But some undoubted cases of sympathetic ophthalmia have occurred after non-perforating injuries. Bronner,¹⁶ of Bradford, has recorded a case where sympathetic ophthalmia appeared in the right eye eighteen days after a kick on the left, which had caused no external wound, and persisted two years in spite of enucleation. Donaldson,¹⁷ of Londonderry, has recorded another case, where a blow on the eye caused a subconjunctival rupture of the sclerotic without external wound. The eye was enucleated in 20 days, but 27 days after the operation sympathetic ophthalmia appeared in the other. The enucleated eye showed a plastic cyclitis, which is the constant appearance of eyes which have caused sympathetic ophthalmia. Nieden¹⁸ has recorded a case of sarcoma of the choroid, causing iridocyclitis in the same eye, and followed by sympathetic ophthalmia in the other. The eye was enucleated, and submitted to Deutschmann for examination. He reported that both in the neighbourhood of the tumour and in the optic nerve were numerous short thick bacilli.

Deutschmann explains these cases of sympathetic ophthalmia without external wound of the eye by supposing that the micro-organisms are carried by the blood stream, and settle in the eye already weakened by tumour or traumatism. Panas¹⁹ has brought forward a great array of evidence to prove the frequent occurrence of endo-infection of the eye, as in the metastatic panophthalmitis sometimes seen in puerperal fever, ocular tuberculosis, gonorrhoeal iritis, etc., and no doubt bacterial infection may occur in an eye without external wound. He explains sympathetic ophthalmia, not very satisfactorily, as due to "the general dyscrasia of the patient."

In a case of this kind (that is, in a case of sympathetic ophthalmia) is the endo-infection of the sympathetically affected eye due to toxins or to attenuated germs carried in the blood stream? It is impossible to say. The presence of germs is not constant, but it is an undeniable fact that micro-organisms readily become localised in a traumatised eye..... When one eye has been irritated by a traumatism the other eye becomes the seat of various phenomena of vaso-dilatation. If the patient is already in a condition of toxæmia he will develop sympathetic ophthalmia; if he is not toxæmic all will be limited to some simple functional disorder.

To this de Wecker very pertinently objected that sym-

pathetic ophthalmia is most frequently found in vigorous patients. An explanation adopted by Schmidt-Rimpler,¹⁰ Baquis,⁷ Galezowski,²⁰ and others is that sympathetic ophthalmia is due to bacteria or their products carried by the blood vessels or lymph channels, acting on the second eye already subjected to irritation of the ciliary nerves. But a sympathetic irritation is not always seen before sympathetic ophthalmia sets in. Further, if it is present it disappears at once if the exciting eye is enucleated, whereas sympathetic ophthalmia may appear even a month later. Priestley Smith²¹ does not consider this latter objection to be very strong, remarking that in syphilis there is always a considerable interval between the primary infection and its remote manifestations.

According to the nerve theory sympathetic ophthalmia depends on irritation of the ciliary nerves in one eye causing a reflex or sympathetic irritation in the other. These nerves are divided into the long and short sets, both derived from the sensory portion of the fifth nerve, the former set direct from its nasal branch, and the latter through the lenticular or ophthalmic ganglion, which receives also twigs from the sympathetic.

There are several great difficulties in the way of this theory as well as those I have mentioned as explained by Deutschmann's theory. One often raised is that an iridocyclitis in one eye is a necessary condition for the occurrence of sympathetic ophthalmia, which is also an iridocyclitis, in the other; and in the great majority of cases, this iridocyclitis in the injured eye is of an infective origin. Mechanical irritation, apart from iridocyclitis, and unless it first causes the latter, will not produce sympathetic ophthalmia, so that we should have to assume that this nerve irritation, supposed to cause sympathetic ophthalmia, can itself only be caused by bacteria or their products, an assumption for which there is no analogy, as Schirmer²² points out. Nevertheless the adherents of the migratory theory admit the existence of a definite sympathetic irritation, causing physiological changes in the second eye, and as this sympathetic irritation is only seen in cases where sympathetic ophthalmia might be expected to follow, it must be assumed that it also is caused by bacterial irritation in the exciting eye, and by nothing else. And though it is true that it is generally clearly distinguished from sympathetic ophthalmia, and may last many months without being followed by sympathetic ophthalmia, still it is not always clearly distinguished, and, according to Fuchs,²³ often passes insensibly from simple irritation to actual inflammation.

Lately some positive evidence in favour of the ciliary nerve theory has been brought forward by Bach.⁵ He irritated one eye in a rabbit by irrigating with a solution of perchloride of mercury, by electric currents, by mechanical irritation, etc., and immediately removed the other eye under chloroform and examined it microscopically. On careful examination he found various changes—fibrinous exudations, and in some cases corpuscular elements of the blood, both in the anterior chamber and between the ciliary processes. He concluded:

That a prolonged irritation of one eye produces vascular dilatation and escape of the blood constituents, not only in this eye but in the other eye also, and that these changes are the first stage of an inflammatory process which, under long continuance of severe irritation, develops into sympathetic ophthalmia.

My own work has consisted of two proceedings—first, the pathological examination of eyes removed on account of injury; and, secondly, an experimental research to determine what changes, if any, follow prolonged irritation of one eye, both with and without traumatism.

I have examined eight injured eyes, four of which had already given rise to sympathetic ophthalmia in the other eye. In one case the injury to the exciting eye had been received two years before the onset of inflammation in the sympathising eye, and in the others less. Of the other four eyes, which had not caused sympathetic ophthalmia, three had received perforating injuries and one non-perforating—a blow on the eye causing hyphema and glaucoma—the eye which was useless being removed to relieve pain. In another of the four the eye had been injured eleven years before, and had lately become painful. In the third of this set (described and figured in the *Ophthalmic Review* for April, 1897) there were optic neuritis and commencing atrophy of the globe, and

the fourth had been wounded by a piece of lead from a rifle bullet, iritis developing after seven days. In none of these four did sympathetic ophthalmia arise after enucleation. All these cases were carefully examined for micro-organisms and all gave a negative result. In the cases where sympathetic ophthalmia had arisen in the other eye evidences of iridocyclitis were found, as was to be expected. These observations, therefore, disagree with those of Deutschmann, and support those of Randolph, Baquis, and Bach quoted above.

In the experimental research my aim was to discover whether prolonged irritation, external or internal, of a rabbit's eye would produce any visible organic changes in the ciliary region, either in the irritated eye or in the other. In the first experiments the eye was cocaineised, and a wound was made in the ciliary region without antiseptic precautions, and the cornea irritated with powdered jequirity seeds for varying periods, up to six months. The jequirity causes purulent conjunctivitis, and abundant cocci were present in the pus, but though in some cases the ciliary region was again wounded while the eye was bathed in pus, no iritis was caused. In one case there was an extensive ulcer of the cornea, which ultimately healed without perforating. The non-irritated eyes were examined at intervals with the ophthalmoscope, but no departure from the normal appearance of the disc was seen in any case.

Internal irritation was then tried, small shot being inserted into the eye, again without antiseptic precautions. In fact, soiled instruments were purposely used, to reproduce as far as possible the kind of wound which is most often found to cause sympathetic ophthalmia. When the incision was made in the sclerotic a few millimetres above the corneal margin, a little vitreous escaped on the shot being put in, but the wound healed rapidly, and in a few days no sign of it, or of external irritation, could be seen. When the wound was made in the ciliary region more external irritation was seen. In one case the wound opened after several days, the shot was expelled, and all signs of irritation disappeared. In another the shot was similarly expelled, but after some weeks, and irritation is still present at the end of five months, as shown by injection in the neighbourhood of the wound, but no signs of iritis are to be seen, either in the traumatised eye or in the other. This rabbit I have kept alive to observe how long the irritation lasts, and whether it will yet lead to any changes in the iris or ciliary body.

After periods varying from three to four months, the rabbits were killed, the eyes and parts of the optic nerves removed and hardened in formol, frozen, cut in two, further hardened in spirit, mounted in celloidin, and sectioned. The sections have been examined for micro-organisms and for deposits of fibrin or leucocytes in the ciliary region.

No trace of bacteria could be found in any case, and in only one were there signs of iritis, in the case where jequirity had caused the corneal ulcer. Here there was some exudation in the anterior chamber, probably akin to that seen in hypopyon ulcer, but there was no change in the ciliary body, which appeared quite normal.

We must conclude, therefore, that Bach's assumption is not warranted by the facts, and that, though a temporary irritation of one eye may cause a temporary exudation of fibrin and leucocytes both in the irritated eye and its fellow, it is not the case that long-continued irritation will cause abundant exudation and consequent organic changes in the second eye.

Further, it is plainly a matter of some difficulty to produce an irido-cyclitis in the eye of a rabbit, and in any further experiments I should be inclined to select some animal in which the ciliary muscle is better developed, and presumably more active, than it is in the rabbit.

SUMMARY.

Practically the whole weight of the experimental evidence, and the greater portion of the clinical evidence, which has accumulated since the publication of the migratory theory, either fails to support that theory or is actually opposed to it. Even Deutschmann, the propounder of the theory, has in his later papers come very near contradicting his earlier observations; and, if his theory is to stand, it can only be on the assumption that the migratory germs are of such a nature that they can neither be inoculated, cultivated, nor stained

by any known method—an assumption for which there is no ground whatever.

It seems to me that we are obliged to fall back on some form of the old theory, that sympathetic ophthalmia is due, partially if not entirely, to irritation of the ciliary nerves. No doubt there are difficulties in the way, which experimenters have so far failed to remove, but in seeking for an explanation of these failures several possibilities are open to us.

The common occurrence of sympathetic irritation in one eye after injury to the other is generally acknowledged, and it may be that the mechanism by which this irritation is conveyed differs in man from that in dogs, rabbits, and guinea-pigs, and this seems not so unlikely when we remember the anatomical arrangement of the whole ocular apparatus in man, designed to secure binocular vision. Or it may be that the explanation lies in the anatomical and physiological relationships of the nervous mechanism conveying the irritation, of which we as yet know nothing, and I believe that it is by the study of these relationships that our knowledge of the cause of sympathetic ophthalmia is most likely to be extended.

I have to acknowledge my great indebtedness to Dr. Lorrain Smith, not only for his permission to work in the Pathological Laboratory of Queen's College, but also for valuable aid and advice in performing the experiments, and in the course of the research.

REFERENCES.

- ¹ Von Graefe's Archiv, xxx, pt. 3, xxxi, pt. 2. Ueber die Ophthalmia Migratoria (L. Voss, Hamburg), 1889. Beiträge f. Augenhell., March, 1893.
² Archiv. of Ophthal., vol. xv, p. 281. ³ Ibid., vol. xvii, p. 187. ⁴ Quoted in Oph. Rev., 1891, p. 301. ⁵ Von Graefe's Archiv, xlii, pt. 1, 1896. ⁶ Archiv f. Augenhell., xxii, pt. 1. ⁷ Etude sur l'Ophthalmie Sympathique. Rep. of Internat. Med. Congr. at Rome. ⁸ Archiv f. Ophthal., xl, pt. 4. ⁹ Ibid., xxxviii, pt. 1. ¹⁰ Ibid., xxxviii, part 4. ¹¹ Ibid., xl, p. 46. ¹² Quoted by Berry in Edin. Med. Journ., March, 1897, p. 334. ¹³ Ann. d'Ocul., cxii, 14. Quoted by Velhagen, vide infra. ¹⁵ Archiv. of Ophthal., xxv, pt. 3. ¹⁶ Trans. Oph. Soc., 1894, p. 215. ¹⁷ Oph. Rev., 1897, p. 35. ¹⁸ Trans. Internat. Oph. Congr., 1894, p. 205. ¹⁹ La Semaine Médicale, 1897, p. 218. ²⁰ Ibid., p. 220. ²¹ Oph. Rev., 1893, p. 96. ²² Von Graefe's Archiv, xxxviii, pt. 4, 1892. ²³ Textbook of Ophthalmology, London, 1893, p. 293.

OPHTHALMITIS OCCURRING LONG AFTER ENUCLEATION OF THE FELLOW EYE FOR INJURY.*

By GEORGE FERDINANDS, M.D.,
Aberdeen.

ANY morbid condition occurring in an eye after its fellow has been removed for injury is of great interest to the ophthalmic surgeon. The two following cases which have come under my notice are worthy of record, especially as they tend to throw some doubt on the infallibility of preventive enucleation.

CASE I.—J. S., when 5 years old, injured her right eye while attempting to cut a garter with a knife. She was taken to the Aberdeen Royal Infirmary, where, according to the statement of the patient's mother, an operation was performed on the injured eye—probably an iridectomy—eight or nine days after admission. Three weeks later the patient was discharged. Not long after—the exact period cannot be ascertained—the parents noticed that the uninjured eye was “weak,” and the child could not bear light. Evidently the symptoms of sympathetic mischief had set in. The patient was taken back to the infirmary, where the exciting eye was removed, not, however, without some delay occasioned by the reluctance of the parents to consent to the removal of the injured organ. The sympathising eye then rapidly recovered, but, to use the patient's own words, “the eye was never very strong, especially in the winter.” On December 15th, 1891, that is, fourteen years after the accident, the patient consulted me for the first time. She wished to begin wearing an artificial eye but the socket would not admit one. On examination the socket was found occupied by a mass of hypertrophied conjunctiva, and a narrow band of cicatricial tissue limited the movement of the upper lid. The swollen conjunctiva was removed, the cicatricial band divided, and the patient was ultimately able to wear a comfortably-fitting glass eye, which she only used occasionally. On March, 1895, that is, three years and some months after the patient commenced the use of an artificial eye, she was led into my consulting room, and the following was her story:—“For months past her health had not been good, and her eye grew ‘weaker and weaker,’ until the sight got so bad that she could not see to move about.” Thereupon she thought it time to seek advice.

On examination the eye was found sensitive to pressure, the conjunctiva and sclerotic were injected, the cornea studded with keratitis punctata and the iris inflamed. V = fingers at 2 ft. barely. The socket showed no sign of inflammation neither was it sensitive to pressure. The patient was ordered atropine and put under mercury and iodide of

potassium. Under this treatment the eye made rapid progress, the vision being on June 21st $\frac{2}{5}$ and J. 1. Unfortunately this improvement was not permanent, and the eye has had several relapses, each of which only yielded temporarily to mercurial inunction. At present the vitreous shows signs of degeneration, and there are large floating opacities. V = $\frac{1}{8}$. The eye has to be kept continually under atropine, as its cessation causes a return of the active symptoms.

CASE II.—J. W., farm bailiff. In 1875 injured his left eye; in his own words, “the eye was reduced to a pulp.” It was removed four days after the injury. In 1896 he found that the artificial eye he was wearing irritated the socket, and as its use gave him “a pain over the brow” he discontinued wearing it. About this time he began to notice that he could not see to read so clearly as before, and this disability gradually increased until the eye showed “signs of inflammation.” The patient first consulted me on June 26th, 1897. Examination of the eye revealed well-marked irido-cyclitis and a subcorneal deposit. The eye was very sensitive to touch, and the patient complained of severe “neuralgic” pain in the brow and temple. V = fingers at 3 ft. The eye improved under mercury and atropine; the subcorneal deposit cleared up so that in July vision improved to $\frac{2}{8}$. In August, however, the inflammatory process returned in all its activity, in spite of treatment. In February, 1898, the cornea began to ulcerate. Just now the eye is quieter, and the corneal ulceration shows signs of healing. V = fingers at 4 ft. No view of the fundus could be obtained.

REMARKS.

The question of the difficulty of deciding whether such cases as narrated above are cases of true sympathetic ophthalmitis was fully discussed at the Ophthalmological Congress at Heidelberg. In the cases now recorded, the long interval between the removal of the injured eye and the onset of the inflammation is, I believe, without precedent. Of the thirty cases examined into by the Committee of the Ophthalmological Society, the interval between enucleation and the first appearance of sympathetic mischief was under three weeks in twenty-five cases, more than three and up to eight weeks in five cases. Since then Leopold Weiss¹ has recorded a case, which he regarded as one of sympathetic ophthalmitis, occurring nearly two and a-half years after enucleation for injury. So far as my own observation of the two cases goes, the symptoms and the progress are identical with those of sympathetic ophthalmitis. In Case I there is one point worthy of notice—that the inflammation appeared to follow the irritation caused by an artificial eye. The wearing of an artificial eye over a shrunken stump has been known to excite sympathetic mischief,² but in Case II the eye had evidently been wholly removed.

Hitherto in our treatment of injuries of the eye we have confined our attention almost exclusively to the injured organ itself. The recently-published experiments of Bach³—who found that irritation of an eye causes almost simultaneous sympathetic irritation of its fellow—while throwing fresh light on the probable cause of sympathetic inflammation, also indicates the direction in which additional preventive measures might be taken. The visual organs are so closely connected that clinically at least they should be regarded as one organ. In severe injuries of an eye, therefore, it may be well to extend our palliative treatment to its fellow.

REFERENCES.

- ¹ Archives of Ophth., xxii, 1. ² Lawson, Ophth. Hosp. Rep., 1867-69, vi. ³ Von Graefe's Arch., xlii, 1.

A CASE OF ACQUIRED NYSTAGMUS.

By ARCHIBALD S. PERCIVAL, M.B., B.C. CANTAB.,
Ophthalmic Surgeon to the Children's Hospital, Newcastle-upon-Tyne.

THE following case affords confirmatory evidence in support of the view that acquired nystagmus is essentially due to fatigue of the ocular muscles, and not to deficient illumination. The former view, as is well known, has been ably supported by Snell, Dransart, Nieden, and others.

J. A., a railway clerk, came to me complaining of dimness of sight. His visual acuity was normal when tested with test types under good illumination, and no abnormality of his fundus was detected on ophthalmoscopic examination. On inquiry as to the character of his work, he said that he was engaged all day in a well-lighted room, adding up long columns of figures in a large folio account book; he then declared that the figures that he looked at seemed to dance up and down.

On putting him into the position that he ordinarily assumed when at work, with his eyes running up the edges of the page of a large book, a vertical nystagmus at once became manifest. It is noteworthy that this nystagmus failed to appear on simply rotating his eyes upwards. He did not complain spontaneously of hemeralopia, but on inquiry he said he thought he saw less well than other people when the illumination was defective. I took him therefore into the dark room, and comparing his visual acuity with mine (which is presumably normal), I found that with

* Exhibited at a meeting of the Aberdeen Medico-Chirurgical Society