final drop only occupied forty-eight hours. On the other hand it was, in one case, prolonged to six days.

One significant fact may be added. The symptoms which have been suggested as essentially toxic were in some degree a constant feature, and their incidence and subsequent intensity bore an evident relation to the fall in blood pressure.

REFERENCE.

1 BRITISH MEDICAL JOURNAL, November 4th, 1899, p. 1282.

THE TREATMENT OF DIPHTHERIA.*

By A. JEFFERIS TURNER, M.D.LOND.,
Honorary Physician to the Hospital for Sick Children, Brisbane.

In the year 1891 a new era in the treatment of diphtheria was introduced by a remarkable series of investigations published by Professor Behring, now of Marburg University. Although, as Behring remarks, all the essential points in the antitoxin treatment of diphtheria had been by that time fully worked out in the laboratory, the subject did not at first attract very much attention. It was not until 1894, when the acquisition of serum from large animals, particularly horses, permitted the extension of the methods discovered in the pathological laboratory to the treatment of human beings that general interest was aroused. The treatment rapidly spread, and by the commencement of 1895 was on its trial throughout the whole civilised world. In the five years that have since elapsed the diminution in the hospital mortality of diphtheria has been nothing short of marvellous to all but the few who had made themselves acquainted with the thorough and unimpeachable experimental basis on which the new treatment had been established. The records of hospital experience have in this instance been in complete accord with the conclusions of the laboratory. In private practice also, in Australia at least, antitoxin has been very widely used with very gratifying results in the reduction of the mortality from this disease. Nevertheless, its use is not yet universal, and it is every now and then our sad experience to see children admitted into hospital dying from diphtheria, to whom an injection of serum has never been administered, although they have been under medical treatment. In England this appears to be the case to a larger extent. One reads with astonishment¹ that the reduction of the diphtheria mortality in London is almost entirely confined to cases treated in hospital, and that it appears very doubtful whether antitoxin is much used in general practice. Though the contrary statement is true in these colonies, I believe that the remedy, even by practitioners of repute, is not always administered in sufficient doses. In mild and early cases a small dose of 600 to 1,000 units is sufficient, but in advanced and desperate conditions many practitioners are satisfied with injecting 1,000 or 1,500 units, when 4,000, 6,000, or even 8,000 or more units are really needed. The treatment of diphtheria appears to be, therefore, still a proper subject for discussion, and I shall endeavour in the first place to lay before you as briefly as possible the facts on which my opinions are based.

The records of the Brisbane Children's Hospital have been very carefully kept for the past ten years, and the majority of the diphtheria cases during that time have come under my personal observation. I am, therefore, in a position to guarantee the similarity and continuity of the statistical material.

During the preantitoxin period of five and a half years the diagnosis was based solely on clinical observation. Doubtless a few cases not really diphtherial were included, and these would make the mortality appear lower than it really was. I do not think they made much difference in the totals.

The antitoxin period is divided into five sections, of which the results of the two first have been already published.² In them we have included no cases but such as were verified by bacteriological examination. The new treatment being on its trial, such a course seemed advisable. During the next two sections we have included only cases which were verified bacteriologically, or which on clinical grounds could be recognised as diphtheria beyond any reasonable doubt. In the last period we have had to rely on clinical evidence alone,³ but

this has been interpreted with great strictness, much greater than in the preantitoxin period. In the absence of bacteriological examination the diagnosis of early and mild cases must often be doubtful, but in the severer cases, including the laryngeal cases, I am convinced that no appreciable fallacy has been introduced. All cases of possibly catarrhal laryngitis have been excluded.

The first division of the antitoxin period may be regarded as to some extent transitional. We used a serum of unknown but certainly very low antitoxin strength, of which 23 c.cm. was usually injected. In the second division the same serum was mainly used in doses up to 100 c.cm., and much better results were obtained. Since then we have used a serum of known antitoxin strength, and severe cases have usually received 4,000 units. Latterly, we have increased this dose in some cases to 6,000, 8,000, or even 12,000 units, and we believe that the lowered mortality of the last division is partly attributable to this increase.

The following table gives the total mortality for both preantitoxin and antitoxin periods:

Cases of Diphtheria treated in the Children's Hospital, Brisbane.

Preantitoxin Period.	Admissions.	Deaths.	Mortality.
July, 1889, to June, 1891 (2 years)	73	34	Per Cent. 46.6
July, 1891, to June, 1893 (2 years)	114	48	42.1
July, 1893, to January, 1895 (18 months)	116	46	39.7
Total	303	128	42.2

Antitoxin Period.	Admissions.	Deaths.	Mortality.
Small doses : January, 1895, to November, 1895 (11 months)...	40	10	Per Cent. 25.0
Large doses : November, 1895, to September, 1896 (10 months)	43	5	11.6
September, 1896, to December, 1897 (15 months)	83	8	9.8
January to December, 1898 (12 months)	45	7	15.6
January to August, 1899 (8 months)	106	10	9.4
Total	317	40	12.6

In presenting these statistics, I must not omit to acknowledge my indebtedness to the hospital residents, L. N. Ashworth, M.B., B.S.Melb., T. E. Green, M.B., B.S.Melb., Hugh Bell, F.R.C.S.Eng., and P. Dixon, M.B. Ch.M.Sydney, to whose assiduous care the good results are greatly attributable, and but for whose careful records I should be unable to write the present paper.

Although the whole number of cases is small as compared with other statistics that have been published, they have the advantage of having been treated in the same institution and under the same personal observation. Their lesson appears clear enough, but it is necessary, in order to exclude any possible doubt, to consider objections that have been raised by various critics to other similar statistics.

It has been objected that the introduction of a new treatment is accompanied by greater attention to the cases, and more care on the part of the residents and nursing staff. As far as our Hospital is concerned, I can testify that the greatest possible pains were taken over these cases during the preantitoxin period. It will be shown presently that nearly half of the cases were so severe as to necessitate intubation or tracheotomy, and, allowing for this, the results obtained compared favourably with other institutions at the same date.

Again it has been urged that the introduction of bacteriological methods of diagnosis has led to the inclusion of mild cases which would not formerly have been recognised as diphtheria, and so caused an apparent reduction of mortality. This certainly does not apply to our present figures. Cases not recognisable clinically as diphtheria, but in which diphtheria bacilli are present, have occurred with us very rarely: we have only record of three such cases (which came

* Read in the Section of Medicine of the Intercolonial Medical Congress, held in Brisbane.

under observation in 1897), and these cases have been omitted from our statistics.

Lastly, it is asserted that the type of the disease has changed, and that as a whole it is of a milder character than formerly. For this assertion there is no foundation, in fact it is a pure assumption, which can be satisfactorily met. Some observers have attempted to meet it by classifying their cases as mild, severe, very severe, and so on. This is hardly satisfactory, as the classification depends too much on the individual opinion of the observer. There is, I think, a better way. All are agreed that diphtheria affecting the larynx is a serious disease. By including all cases with a distinct laryngeal stridor* it is possible to draw a sharp objective line including a large proportion of the cases, which may be known as laryngeal cases. These cases are always serious, and among them the mortality used to be very high. It has now been reduced more than two-thirds, as is shown by the following table:

Laryngeal Cases.

Preantitoxin Period.	Cases.	Deaths.	Mortality.
July, 1891, to January, 1895*	147	87	Per Cent. 59.2
Antitoxin Period.			
Small doses : January, 1895, to November, 1895	22	9	40.9
Large doses : November, 1895, to September, 1896	30	4	13.3
September, 1896, to December, 1897	37	7	18.9
January to December, 1898	20	6	30.0
January to August, 1899	68	7	10.3
Total	177	33	18.6

* Record incomplete before July, 1891.

In the preantitoxin period very few cases recovered without operative interference, only about 1 case out of 12. Now, although we never fail to intubate a laryngeal case as soon as it appears necessary, nearly half the cases recover without interference. No doubt with earlier treatment the proportion would be larger, for many of the hospital cases require operative interference immediately or very soon after admission.

Laryngeal Cases.

Preantitoxin Period.	Cases.	Recovered without Operation.	Proportion.
July, 1891, to January, 1895	147	12	Per Cent. 8.4
Antitoxin Period.			
Small doses : January, 1895, to November, 1895	22	3	13.6
Large doses : November, 1895, to September, 1896	30	12	40.0
September, 1896, to December, 1897	37	12	32.2
January to December, 1898	20	8	40.0
January to August, 1899	68	33	48.5
Total	177	68	38.4

Operative interference is still necessary in more than one half of the hospital cases. In private practice it should be much less frequently necessary. But even in hospital cases requiring operation the mortality has been reduced from nearly two-thirds to about one-fourth of the cases operated on.

I regard the total results obtained in these laryngeal cases as of the greatest value, inasmuch as they deal with a sharply-defined group of cases of a very grave form of diphtheria, and are not liable to erroneous diagnosis in any appreciable proportion so long as cases of possible catarrhal laryngitis are excluded, as we have been careful to do. They are furthermore fairly comparable with the hospital returns of laryngeal cases in other parts of the world, which the total admissions are not, for these are affected by the proportion of mild tonsillar

cases admitted. The total results may be conveniently stated as follows:—Of 100 cases of diphtheria with affection of the larynx treated in the hospital in the preantitoxin period, 8 recovered without operative interference, 33 recovered after either intubation or tracheotomy, and 59 died. Of 100 similar cases treated with antitoxin in sufficiently large doses, as many as 40 now recover without operative interference, another 45 after operation, and only 15 die.

Operation Cases (Intubation or Tracheotomy or Both.)

Preantitoxin Period.	Operation Cases.	Deaths.	Mortality.
July, 1889, to June, 1891 (2 years)	42	28	Per Cent. 66.7
July, 1891, to June, 1893 (2 years)	60	41	68.3
July, 1893, to January, 1895 (18 months)	64	40	62.5
Total	166	109	65.7
Antitoxin Period.			
Small doses : January, 1895, to November, 1895 (11 months) ...	19	9	47.4
Large doses : November, 1895, to September, 1896 (10 months)	18	4	22.2
September, 1896, to December, 1897 (15 months)	25	7	28.0
January to December, 1898 (12 months)	12	4	33.3
January to August, 1899 (8 months)	35	7	20.0
Total	109	31	28.4

The next table shows the difference in the mortality of all the diphtheria cases admitted under the old and new methods of treatment at each year of life.

Mortality of all Cases of Diphtheria at each Year of Life before and after the Introduction of Antitoxin.

Age Last Birthday.	Preantitoxin Period.			Antitoxin Period.		
	Cases.	Deaths.	Mortality.	Cases.	Deaths.	Mortality.
Under 1 year...	11	9	Per Cent. 81.8	7	2	28.6
1 year	44	32	72.7	35	10	28.6
2 years	56	37	66.2	48	10	20.8
3 "	44	49	111.4	44	6	13.6
4 "	53	14	26.4	47	3	6.0
5 "	38	9	23.7	54	4	7.4
6 "	23	10	43.5	42	4	9.5
7 "	16	5	31.3	19	1	5.3
8 "	6	1	16.7	8	—	0.0
9 "	7	—	0.0	6	—	0.0
10 "	—	—	0.0	5	—	0.0
11 "	1	1	100.0	2	—	0.0
12 "	3	—	0.0	—	—	0.0
13 "	1	1	100.0	—	—	0.0
Total	303	128	42.2	317	40	12.6

The decrease in mortality is striking at all ages, and is particularly well marked during the first two years of life, which had under old methods of treatment such a fearful death-rate.

To these statistics I might add others equally convincing from Australia, from America, from France, from Germany, from England. If hospital records can prove anything, the controversy regarding the efficacy of antitoxin in the treatment of diphtheria may be regarded as closed. But it yet remains to consider in what way the remedy should be used to obtain the best possible results in reducing the mortality of the disease. It will clear the way to the understanding of this problem if we consider it very briefly from the standpoint of recent pathology.

Diphtheria is known to be caused by a bacillus which secretes a powerful toxin. The local irritation produces a fibrinous exudation which forms the diphtheritic membrane, and the diphtheria bacilli are confined to this membrane and the fluid discharged from it. The inflammatory processes have a tendency to spread locally, especially into the air passages. In this way the toxic organism may reach the lungs, but it does not invade the other organs of

the body, which contain no more than a few stray individuals, of no practical importance, who may happen to be washed there by the blood current. Apart from the mechanical obstruction of the breathing all the danger of the disease results from the absorption of diphtheria toxin from the local focus of disease. The antitoxin contained in the serum derived from artificially-immunised horses has no direct effect on the diphtheria bacilli, but neutralises the toxin produced by them. This it does by some form of chemical action, neutralising the toxin in a test tube in strictly equivalent proportions, the two behaving in this respect exactly like an acid and an alkali. As a matter of experience it is found that when their toxin is thus neutralised the diphtheria bacilli in the local lesions become harmless; they are dealt with by the tissues as ordinary saprophytic bacteria, and the case so treated makes a speedy recovery to the normal condition. When mixed in a test tube the proportion of antitoxin required to neutralise many times the lethal dose of toxin is very small. When the two are injected into an animal simultaneously or with only a short interval, the proportion is somewhat larger, but still remains small; for antitoxin and toxin behave to each other in the blood stream as they do in the test tube. But if a multiple of the lethal dose of toxin be injected twenty-four hours or more before the antitoxin, the proportion of the latter required to save life becomes enormously increased, and a thousandfold the dose sufficient in the former instance may be now inadequate. The explanation of this is that the toxin is no longer wholly contained in the blood stream. It has entered into some kind of combination with those tissue cells which it particularly affects. In the case of tetanus, for example, a minute dose of antitoxin may be sufficient if injected simultaneously, while the largest possible dose may be insufficient when the toxin has attached itself to the substance of the cerebro-spinal motor cells. That it does so attach itself is shown by the fact that if the toxin is mixed with an emulsion of cerebro-spinal tissue in sufficient proportion before injection, it becomes harmless. In tetanus, the combination once formed is unfortunately very stable, but in the case of the diphtheria toxin it appears that by maintaining a very high degree of antitoxin strength in the blood, the toxin may be neutralised even after it has fixed itself in the tissues. But to do this, very large doses are necessary, and the longer the interval allowed before their use, the larger must they be.

Applying these considerations to practice, we should, in the first place, inject our remedy as early as possible, as soon as we have made our diagnosis, and frequently in doubtful cases before a positive diagnosis is possible. In very early cases an ordinary dose of 600 or 1,000 units is quite enough. The course of the disease is rapidly arrested and convalescence is speedy and complete. But it is quite different with cases of long duration, presenting extensive membrane, extending, it may be, into the nose or larynx, or in cases showing symptoms of toxæmia. In these, the patient, in spite of appearances in some cases, is in a condition of imminent peril. The toxin is already fixed in his tissues, and death from implication of the central nervous system, heart, or kidneys will very likely occur. We may now be too late to avert the fatal result, but fortunately we may often do so by using very large and, if necessary, repeated doses. In a really bad case of this description I should administer from 6,000 units at once, and perhaps inject an additional 2,000 or 3,000 units or more within the next twenty-four or forty-eight hours. No time must be lost in producing a high degree of antitoxin strength in the patient's blood. According to my experience, about twenty-four or sometimes forty-eight hours elapse before improvement begins; it is then often rapid. The antitoxin appears to require this time to produce its effect, and in cases which die within twenty-four or forty-eight hours of administration the largest doses sometimes do not appear to have any effect. Notwithstanding this, the good effects of large doses in advanced cases have been well illustrated in our experience. During the first eleven months of antitoxin treatment our dosage was too small, and the mortality only showed a reduction of 17 per cent.; but, by using much larger doses, we immediately obtained a further reduction of 13 per cent., and this has been well maintained since. The following two cases will serve to illustrate recovery from grave toxæmia:

M. D., aged 4 years, admitted on the fifth day of illness. Membranous patches on tonsils and pharynx. Breathing with stridor and recession, colour dusky, 5,000 units injected. Intubation soon after admission. That evening, though breathing was easy, the child had a bad colour, lips were pale and dusky, pulse rapid and weak, and she was very restless. These symptoms pointed to grave toxæmia, and 3,000 additional units were injected. Prognosis unfavourable.

Second day, no improvement. Tube coughed up once, and replaced. Râles and rhonchi in chest.

Third day, condition still very grave. Urine with one-sixth albumen; 4,000 units injected. Tube was coughed up, and she had several attacks of dyspnoea, which were not relieved by reintubation. After a severe attack with cyanosis, Dr. Dixon performed tracheotomy, and extracted a large cast of membrane from the trachea. After operation pulse was extremely weak.

Fourth day, slight improvement, gradually leading to convalescence. The total amount of antitoxin administered was 12,000 units.

E. C., aged 6 years, admitted on the eighth day of illness. Membranous patches on tonsils and pharynx. Breathing with stridor and recession; 6,000 units injected.

Second day, breathing worse. Intubated; during operation coughed up a large piece of membrane. Another 3,000 units injected. Pulse poor. Urine scanty: 2 ounces since admission, one-third albumen. Ordered pilocarpin 1.20 gr. and hot-air bath, which caused free perspiration.

Third day, urine 3 ounces, about one-quarter albumen. Hot-air bath repeated. Another 3,000 units injected. Pulse still poor.

Fourth day, urine 3 ounces, one-sixth albumen. Hot-air bath repeated. Seems brighter. Intubation tube removed.

Fifth day, better. Perspiring freely. Urine 5 ounces, trace albumen. From this point steady improvement to convalescence. In all 12,000 units were administered.

There is much to be learnt from an analysis of the fatal cases. In 1897, 1898, and 1899 we have had 25 deaths; of these, 16 occurred within forty-eight hours, and 8 within twenty-four hours of admission. They may be classified according to the main cause of death as follows:

Primary toxæmia	6 cases
Bronchial membrane (in two cases combined with pneumonia)	4 "
Pneumonia	5 "
Secondary toxæmia (with scanty urine or suppression)	9 "
Accidental asphyxia (from coughing out tracheotomy tube during night)	1 "

Total ... 25 cases

The first two headings caused 10 deaths, all of which occurred within forty-eight hours of admission. One, an infant of 18 months, was admitted on the third day of illness. At this age the disease runs an exceptionally rapid course. Two were admitted on the fourth, the remainder on the fifth to seventh day of illness. In all the cases under this heading the disease was apparently too far advanced on admission to be combated. They died as most cases used to die before the antitoxin treatment was introduced.

Pneumonia was responsible for 9 deaths, of which 7 occurred within forty-eight hours. Since the introduction of the new treatment this cause of death has come into much greater prominence; for while the antitoxin will neutralise the diphtheria poison, it has no effect on this complication, which is due to a secondary infection. In two cases examined by me this was found to be due to the pneumococcus, but other observers have found the streptococcus pyogenes to be frequently the excitant cause. Two of these cases, both children aged 2 years, were admitted according to the parents' statements, on the second day of illness. If the statements were correct, they show that even early treatment of the diphtheria by antitoxin will not always prevent a fatal issue from pneumonia. Clinically these cases may be recognised by their extremely rapid shallow breathing, with high temperature and rapid failing pulse. Physical signs are sometimes to be elicited. Several cases were verified *post mortem*. I have seen cases complicated with well-marked pneumonia recover, but it has more often proved fatal.

Under secondary toxæmia I have included a very important class of cases which I have observed before the introduction of serum treatment, but which has assumed a higher relative importance lately. They are nearly always cases of long duration on admission. The earliest was admitted on the fourth day of illness, but this is exceptional, one on the fifth, the remainder had been ill a week or more. They have had extensive areas of membrane in nearly every instance. Some have at first appeared to be doing well, but after several days' treatment, the urine is observed to be scanty and loaded with albumen; sometimes there is complete anuria. At the same time the child develops a painfully-marked restlessness, is totally sleepless, but retains consciousness to the end, while the pulse fades away and often becomes imperceptible at the wrist for many hours before death. These are often classified

as cases of heart failure, but they are not properly regarded as cases of simple heart failure.

That the condition of the kidneys is an important factor in the result I have no doubt; but they are not cases of uræmia in the usual acceptance of this term. Convulsions I have never observed, and coma is absent, being replaced by an exactly opposite condition. I have never seen a case in which these symptoms had developed recover under any form of treatment; but of late I have endeavoured to anticipate their occurrence in cases with scanty albuminous urine by the injection of pilocarpin (gr. $\frac{1}{16}$) and hot-air baths, as in the case of E. C. given above. I believe nasal cases are particularly liable to develop these symptoms, and such cases ought to receive a large initial dose of antitoxin, although their condition at the time may not appear serious.

If we classify our cases since the commencement of the antitoxin period according to the day of the illness on which antitoxin was first administered, we obtain the following:

	Cases.	Deaths.	Mortality.
Treatment commenced on 1st day ...	7	—	
" " 2nd " ...	39	2	
" " 3rd " ...	50	6	
" " 4th " ...	51	4	
Total ...	147	12	8.2 per cent.
" " 5th " or later ...	131	22	16.8 "
Cases of uncertain duration...	33	6	18.4 "

The two cases fatal from pneumonia, although antitoxin was administered on the second day of illness, have already been mentioned. Of the six deaths among those admitted on the third day, one occurred during the first period of antitoxin treatment when the doses administered were too small. Of the remaining five, four died from pneumonia; the last case was an infant of 18 months, admitted moribund from toxæmia on the third day of illness, and has been already alluded to. It is the only case we have seen since 1895 in which death occurred from uncomplicated diphtheria treated with antitoxin during the first three days of the disease.

These results emphasise the importance of early treatment. Every medical practitioner ought to have a supply of diphtheria serum and a suitable syringe at hand. In the large towns there is no difficulty in obtaining a supply, and it is easy to keep a few bottles of the liquid serum and to replace them as needed. The liquid preparation is the easiest for use, and may be kept for six months or more without losing its efficacy. Practitioners in the bush townships should keep a supply of the dried serum, which can probably be relied on to retain its strength unimpaired for years. It is absurd that a practitioner several days' journey from the metropolis, meeting with a case of diphtheria, should have to telegraph for antitoxin. This is too often not of any use to him by the time he gets it.

All other forms of treatment are of very minor importance. Routine applications to the throat have been altogether abandoned in the Children's Hospital. We only use them when the membrane is sloughy and offensive. It has been argued on theoretical grounds that as antitoxin only neutralises the poison, something should be done to kill the bacilli. But I have found by bacteriological examination that in the majority of cases the bacilli disappear within a few weeks without any local treatment; and in those cases in which they persisted longer I have been unable to hasten their disappearance by such treatment. Local applications are, in nine cases out of ten, unnecessary; and unnecessary treatment is bad treatment. When, however, there is much nasal discharge, gentle syringing with salt and water (a teaspoonful to a pint) is advisable. Internal medication has also been abandoned as a routine measure. When the pulse is showing signs of failure, we have given digitalis, strychnine, and alcohol in moderate doses, and these may be of some service. When the urine is very scanty and albuminous, I have lately administered pilocarpin hypodermically (the dose should not exceed gr. $\frac{1}{16}$), followed by a hot-air bath, administered by covering the child with a cradle and blankets, under which

the moist hot air of a spirit lamp flame is conducted through a bent metal tube. In cases dying from bronchial membrane or pneumonia, inhalation of oxygen gas is of value in relieving distress and promoting euthanasia. I have not seen it save life.

To sum up the whole matter, the treatment of diphtheria consists in the administration of antitoxin early and in sufficient dose. A trustworthy preparation of known antitoxin strength should be employed. There need be no fear of administering an overdose; the only objection to giving an unnecessarily large dose is its cost. To give too little for this reason is a very poor economy. The saving in the number of doctor's visits, not to mention funeral expenses, much more than covers the cost of the serum. In the case of poor patients I have been accustomed to explain the danger of the disease, and have never found them object to the expense of the remedy. Those who cannot afford this are suitable cases for hospital treatment. I would certainly decline to take any responsibility, or to continue to attend a case, if the treatment were objected to, and I hope other practitioners would do likewise. I would appeal particularly to those engaged in club practice, who are more bound to their patients, to explain to them the need for immediate injection of serum or immediate removal to hospital. Too many patients of this kind have been brought to us too late.

We may say that if patients and medical men did their duty, the former in calling assistance early, the latter in applying the remedy early and in sufficient dose, the death-rate from diphtheria might be reduced to a small fraction of what it is at present, although it has been materially lowered in this colony of late years. Indeed, with the necessary qualifications, we may say no child ought to die of diphtheria. There are no doubt some cases complicated by pneumonia at a very early stage, some others at a distance from medical aid, some few in which the diagnosis in the early stages is extremely obscure. Cases sometimes occur of laryngeal diphtheria without any visible membrane, and these are sometimes very hard to distinguish from catarrhal laryngitis. In cases of doubt a full dose of antitoxin should be given. Sometimes an attack of diphtheria becomes engrafted on a previous catarrh or tonsillitis, and is then very likely to be overlooked in the early stages. But these are all rare exceptions, for the most part death from this disease is avoidable and ought to be prevented.

NOTES AND REFERENCES.

¹ Goodall, BRITISH MEDICAL JOURNAL, 1899, vol. i, p. 107. ² *Intercolonial Medical Journal*, October, 1896. The figures given of the preantitoxin period contain a few unimportant errors which are here corrected. ³ Owing to pressure of other work, I ceased making these examinations, but I hope they will be soon resumed by the hospital residents, for which purpose a laboratory is being fitted up. ⁴ Cases showing merely hoarseness, croupy cough, or even visible membrane in the epiglottis are excluded. It is always possible to distinguish laryngeal stridor from pharyngeal, stertorous breathing.

MEMORANDA:

MEDICAL, SURGICAL, OBSTETRICAL, THERAPEUTICAL, PATHOLOGICAL, Etc.

A CASE OF MALIGNANT PUSTULE: NECROPSY, WITH BACTERIOLOGICAL INVESTIGATIONS.

On June 4th I was called to see a girl, aged 17, said to be suffering from œdema of the glottis. I found her moribund, and presenting symptoms which suggested the possibility of anthrax—namely, a pustule at the angle of the mouth, swelling and redness of the cheek, general cyanosis and insensibility which was of very rapid onset. The pustule was atypical.

The history was as follows: Eight days previously the patient spent the day in the country at a village C.; three days later she developed a pustule, rapidly became worse, and died within five days of the appearance of the local lesion. She did not dress meat, handle hides or wool, and had nothing to do with animals.

A *post-mortem* examination was made thirteen hours after death, when the pustule appeared more typical; it was black in the centre, about the size of a threepenny piece, but had