showing any evidence of malignant disease that, although in an occasional case malignant disease is reported as having been found at the operation, it is so rarely that I have come to look upon these cases as so certainly non-malignant as to warrant an optimistic pre-operative prognosis.

On the other hand, an *x*-ray experience of the malignant stomach is that it is far more frequently seen in males; that on going into the history carefully, the very large majority state that the symptoms have been present for a matter of a few months only; and that never in their lives have they had any "indigestion" before this attack. There is nothing in the history to suggest in any way a pre-existing ulcer. I say that this is the common history, and that only a comparatively few relate a long-continued history of intermittent attacks of digestive disturbance.

There are also other facts bearing on this question. Duodenal ulcers are extremely common; they are continually irritated; they are rarely malignant. Cancer of the oesophagus is so common that I have seen literally hundreds of cases in the *x*-ray department. The symptoms are sudden and the history almost invariably that of a few months at the most—nothing to suggest previous ulcer.

These facts have produced a very distinct impression and raised very great doubts in one's mind that gastric ulcer is a common antecedent to malignant disease of the stomach. I suggest that these *x*-ray facts will require a lot of explanation before such a cause for cancer of the stomach can be accepted in the vast majority of cases.

SUMMARY.

1. Hour-glass contraction from simple ulcer is a comparatively common condition—frequent in females, rare in males.
2. Its symptomatic history is a long one.
3. It is almost entirely an *x*-ray diagnosis.
4. It rarely ends in malignant disease—especially in females.
5. An hour-glass condition is found in a few cases, mostly in males, and with a short history, and which is caused by malignant disease. Its *x*-ray appearances generally differ considerably from those seen in the non-malignant type, and are suggestive.
6. The experiences of a large *x*-ray department are strongly opposed to the theory that it is usual for cancer of the stomach to be preceded by a simple ulcer.

MESSAGE OF THE HEART AND RESUSCITATION.

BY

J. A. GUNN, M.D., D.Sc.,

PROFESSOR OF PHARMACOLOGY IN THE UNIVERSITY OF OXFORD.

In a paper published in this JOURNAL (November 6th, 1920) Mr. W. E. Fisher describes cases of "Resuscitation in Death under Anaesthesia," and in regard to cardiac massage he says that "it will be of great help if an understanding of the processes concerned is quickened by a clear pronouncement by physiologists, to whom we must look for guidance and more definite directions." I feel sure that he is not aware of a paper published in 1915¹ by P. A. Martin and myself, in which we attempted to give clearer directions for performing cardiac massage than had hitherto been done, and anticipated some of the observations that he has made on the human heart. Since this paper has escaped the attention of others who have subsequently written upon the subject of cardiac massage—as is almost inevitable with the multiplicity of medical journals—I think one may be excused for recapitulating, in a journal more accessible to clinicians, some of the results there recorded, the more so as I may take this opportunity of adding some other observations on the subject of resuscitation based upon unpublished experiments.

I may first state part of the conclusions arrived at by Martin and myself.

"In performing massage of the arrested heart with a view to reviving the spontaneous contractions, the following points are of importance:

"Compression should be gradual and relaxation abrupt. The rate of compression should be at most less than half the normal

heart rate. Massage should be regularly interrupted at short intervals for a few seconds to allow the spontaneous beats to develop. When the heart is arrested, massage, properly performed, acts both as an artificial circulation and as a mechanical stimulus to the heart muscle.

"Massage can be performed in the following ways, arranged in order of efficacy: (1) By direct compression of the heart through an opening in the thoracic wall; (2) by compression of the heart above the diaphragm through an opening in the abdominal wall; (3) by simple compression of the abdomen; (4) by simple compression of the thorax.

"Direct thoracic massage combined with intrapericardial injection of adrenine can revive the heart in practically any case of chloroform poisoning in the rabbit, even though the treatment be not begun until fifteen minutes or more after the heart has completely ceased to beat.

"In about 70 per cent. of cases recovery can be obtained if treatment is commenced ten minutes after arrest of the heart by the following procedure: artificial respiration through a tube in the trachea; intrapericardial injection of adrenine; massage through an opening in the abdomen. If, when the heart has been started, the blood pressure continues low and the heart beats remain (chiefly as a consequence of this) feeble, speedy and permanent improvement of the heart and blood pressure can be obtained by intravenous injection of either adrenine or pituitrin.

"Treatment on these lines is suggested for cardiac arrest in the human subject, occurring either in chloroform poisoning or in other conditions in which general recovery might be anticipated to result from resuscitation of the heart."

Before the publication of our paper the expression "massage of the heart" had been used as if the manner of performing it were indifferent, but we found experimentally that attention to points of detail may make all the difference between success and failure in difficult cases of resuscitation of the arrested heart. The grounds for certain of the conclusions given above may therefore be of value.

One question of the first importance is this, that "rhythmic compression of the heart effects in the first place an artificial circulation; it produces movement of the blood in the vessels to an extent which depends upon the way in which it is done." In animal experiments such as we performed an indication of the mechanical efficiency of massage in effecting a circulation of the blood could be obtained (the blood pressure being at zero) by the amplitude of the wave of blood propelled into the aorta by each compression of the heart. A record of this wave was obtained by a mercury manometer connected with a carotid artery. But a more direct proof of the fact that massage of the heart acts, to begin with, as a substitute for the spontaneously beating heart in causing a circulation of the blood was obtained in the following way: A solution of a blue-green dye "was injected into the cavity of the right ventricle of an arrested heart. Cardiac massage was then performed, and after a few compressions the blue-green colour appeared first in the lungs and then in the carotid artery. Massage had therefore propelled the blood from the right ventricle through the lungs and left side of the heart into the systemic circulation."

Fisher describes this action of massage as "an influence upon the circulation that I have not seen previously observed." It is all the more interesting that he has described independent observations on the same phenomenon but judged by different effects—namely, flushing of the skin and rise of intraocular tension occurring as the result of massage prior to spontaneous beating of the heart.

This action of massage has an importance not merely academic. I have started the heart beating an hour after it has been arrested by chloroform. I have perfused the exsected heart of a stillborn child four hours after the child was born, and started the heart beating again. The difficulty is not so much in starting the heart, though this may of course be difficult, but in starting the heart sufficiently soon that the other tissues have not meantime been rendered incapable of recovery by stoppage of the circulation. So far as is known, it is the cells of the central nervous system that are most susceptible to stoppage of the circulation, so that, from the practical point of view, the circulation must be started before the cells of the central nervous system are irretrievably damaged. The current opinion is that after complete stoppage of the circulation for about fifteen minutes the cortical cells cannot be revived. While my own experiments have also pointed to this conclusion, I am not convinced that it is true; and, for reasons that I need not enter into here, I should not be surprised if some day it were shown that even the cortical cells can survive arrest of the circulation for a much longer period. What is

known is that they are difficult to revive. It is not proved that they are dead.

Many other tissues can withstand absence of circulation for a long period. Thus I have revived the movements of the isolated guinea-pig's uterus after seven days' keeping in cold storage, and the human Fallopian tube after thirty hours;² the rabbit's intestine after five days;³ the human vermiform appendix after thirty-six hours;⁴ and the heart after twenty-four hours⁵ or longer. At higher temperatures they do not survive so long,⁶ but even at temperatures between ordinary room temperature and body temperature they survive for hours. In tissues such as these there need be no anxiety in regard to the possibility of restoring their functioning capacity after prolonged arrest of the circulation. The liver requires further investigation.

In the meantime the question of resuscitation may be taken to depend upon how long the cortical cells can survive arrest of the circulation. And it is in this connexion that there arises the importance of the capacity of cardiac massage to act as an artificial circulation. For the difference between complete and incomplete stoppage of the circulation is enormous when gauged by the effects on the cells of the central nervous system. It has been found, for example, that when an attempt has been made to occlude the cerebral circulation by clamping the vessels to the brain, much more prolonged survival of these cells may occur if a mere trickle of blood through a small vessel be intentionally or accidentally left. Other phenomena point to the same conclusion. For example, in hibernating animals both circulation and respiration are reduced to a very low level for months, and yet the animals revive rapidly and completely when the circulation and temperature, etc., are restored. Now the experiments by Martin and myself are confirmed by Fisher's observations, and it can be taken as proved that massage of an arrested heart, if properly performed, can act as a fairly effective mechanical means of conducting an actual circulation of the blood. This being so, it is more than probable that, for the purpose of deciding whether the cells of the central nervous system are capable of complete restoration or not, the time can be calculated not as from the start of spontaneous heart beats but as from the time of beginning massage. This is a point of fundamental importance, and puts a different and more hopeful complexion upon the whole question of resuscitation.

Method and Rate of Compressing the Heart.

In the paper already quoted we said further:

"Judged merely from the point of view of its acting as an artificial circulation, massage would be most efficacious when carried out uninterruptedly, because the maximum amount of circulation of blood would thereby be accomplished, and therefore the speediest elimination of chloroform. However, we have repeatedly found that this is not the case, the reason being that this procedure fails to take due cognizance of the second function of massage—that, namely, of acting as a mechanical stimulus to the heart itself. The latter is the real object aimed at in performing massage; and, though a mechanical circulation may be a necessary preliminary to obtaining conditions under which the heart can resume its action, it is clear that, even from the point of view of circulating the blood, cardiac massage is an inadequate substitute for the beating heart. It is essential that this mechanical stimulation should be applied in such a way as most effectively to encourage the spontaneous contractions of the heart. This is not attained by continuous massage. It has been found, for example, that massage which has started feeble spontaneous beats, if continued without intermission, may cause the beats to disappear again. No doubt it is possible to overdo mechanical stimulation, and continuous massage does not give the heart a sufficient opportunity of developing spontaneous beating. We have found, as a result of a large number of experiments, that when the feeblest beats have begun, even beats which fail to move the manometer, it is better to cease massage for some seconds so as to allow them to develop, and to apply short periods of massage intermittently, when it is usually found that perhaps by the third or fourth series the heart suddenly regains a normal rhythm and the blood pressure rises rapidly."¹

The most effective rate of compression is much slower than the normal heart rate, for two reasons: One is that a slow rate is necessary to allow complete filling of the ventricles, and the other that it is a subnormal rate of beat that one is attempting to elicit—namely, the rate at which the arrested heart will begin beating again. The advantage of intermittent massage had not previously been suggested. Possibly if attention were paid to this,

attempts to revive the human heart would be more successful.

Action of Adrenalin.

There is no known drug comparable with adrenalin in efficacy to start an arrested heart or to antagonize the action of chloroform. I have shown that adrenalin can start beats in a perfused mammalian heart—for example, when kept for twenty-four hours—when mere perfusion has failed. When the isolated mammalian heart is perfused with concentrations of chloroform or chloral which almost or even completely arrest the heart, the addition of adrenalin to the chloroform or chloral solution is able to restore the heart beats to their normal vigour.⁵ This is an extraordinary effect. It is not an antagonism of the type, for example, of atropine *versus* pilocarpine. For chloroform and chloral depress the heart muscle, and in the experiments referred to adrenalin is able to antagonize a concentration of chloroform or chloral which practically or completely stops the heart while the latter solution is still passing through the heart, and its action therefore getting deeper. I do not think that this power of adrenalin—which is equivalent to stimulation of the sympathetic nerve of the heart—to antagonize a muscular paralysis has been sufficiently realized.

When the circulation is arrested in the intact animal, the difficulty, of course, is to get the adrenalin at the heart, even intravenous injection being practically useless when the heart has stopped. One method is to inject adrenalin into the pericardial sac before beginning massage, but this might be difficult to one who has had no experience of giving such an injection, and I have myself no experience of performing it in man.

Seeing, however, that massage causes an artificial circulation, adrenalin will reach the heart in time from a vein if massage be continued. This will occur sooner the nearer to the heart the vein is into which the adrenalin is injected. If it were injected into an external jugular vein it would very probably reach the arrested heart after a few effective compressions by massage.

One point in regard to adrenalin which may cause confusion is that a combination of adrenalin and chloroform is, as has been pointed out by Levy, more likely to produce fibrillation of the heart than chloroform alone. This, however, refers only to cardiac arrest from chloroform occurring in the initial stages of anaesthesia. This danger does not occur, in my opinion, in cardiac arrest occurring at other times.

There is another point in regard to the action of adrenalin about which there seems to be a good deal of misconception, and that is as to the permanence of its effect. It is true that in the normal animal with a good blood pressure the rise of blood pressure produced by adrenalin is very transient, lasting only a few minutes. But this is not true in certain abnormal conditions of the circulation—for example, when the animal is being resuscitated after arrest of the heart. Here the blood pressure may remain low after spontaneous beats of the heart have begun. This low blood pressure is due partly to temporary paralysis of the vasomotor centre and partly to feebleness of the heart's contractions. These two effects interact on one another. This vicious circle is broken by adrenalin. The temporary rise of blood pressure provides a better blood supply to both heart and vasomotor centre, and this may be sufficient to enable both to function properly, and in these circumstances the rise of blood pressure produced by adrenalin may be permanent. I believe it may be so in other similar conditions of the circulation.

Figs. 1 and 2 are reproduced (by permission) from the paper referred to. They show the sequence of events in a difficult case of restoration of the heart after it had been arrested by chloroform, and massage begun fifteen minutes later. The effects of intermittent massage are shown and the permanent effects of adrenalin injection.

Artificial Respiration in Man.

I have taught my students for years that the time will come when every hospital will be provided with an appliance for mechanically performing artificial respiration in man. So far this prophecy has remained unfulfilled. The usual method employed in the laboratory for this purpose in the case of laboratory animals is a pump or bellows which forces air into the trachea, and so distends the

lungs. It is a method theoretically imperfect, which in practice works well and gives no trouble. I have seen it stated in physiological textbooks that this method of performing artificial respiration is inadmissible in man, but I have not seen any convincing evidence given for this statement. In the case of adults with very rigid chest walls there would be theoretical objections to it, but, with the patient in the horizontal position, very probably an excursion of the lung sufficient to give the minimum necessary aeration of blood could be obtained by that expansion of the lung allowed by little else than movement of the diaphragm. If necessary, oxygen could be used instead of

tube. It was this method that I used in resuscitation experiments on rabbits.

If the heart has stopped, for example, in the course of chloroform anaesthesia, to get the heart started again is only the beginning of resuscitation. The next thing is to get the natural respiration evoked. Now when the circulation has been stopped for several minutes, the respiration does not commence immediately upon the heart beginning to beat again. It takes time for the respiratory centre to recover. In animal experiments I have seen thirty minutes elapse between the starting of the spontaneous heart beats by massage and the beginning of

FIG. 1.

FIG. 1—Rabbit. Heart arrested by chloroform. Showing complete resuscitation, manometer record of blood pressure, and pulse wave in carotid artery. The periods of massage are marked. By the end of Fig. (b) feeble spontaneous beats had commenced, and by the end of (c) the heart was beating regularly. Blood pressure remained low. Intravenous injection of adrenine produced an abrupt rise of pressure (d). The pressor effect was maintained (e, taken twenty minutes later).

air. But, in any case, in children the method of positive artificial respiration ought to prove just as serviceable as in animals like the dog.

Having in view the possible adoption of this method of performing artificial respiration in man in cases of resuscitation, etc., I have modified the usual technique in order to dispense with tracheotomy and the time involved in performing it. Instead of tying a short cannula into the trachea through a tracheotomy opening, a long tube is passed through the mouth and larynx down to just above the bifurcation of the bronchi. This tube should be of about two-thirds of the internal diameter of the trachea. When intermittent positive artificial respiration is used—for example, by a Brodie's pump—each blast distends the lung and the air escapes along the outside of the tracheal

D

natural respiration. More usually it is between five and ten minutes. If artificial respiration is not carried on during this interval, the heart will of course fail again from asphyxia. It is therefore imperative that artificial respiration should be kept up after the heart has begun beating again and until normal respiration begins.

There is no question that the problem of resuscitation is enormously facilitated by the employment of mechanical artificial respiration. With an apparatus such as Brodie's respiration pump with its modern improvements, whereby the amount of air pumped and the rate of pumping can be altered at will, and the air warmed or charged with a volatile anaesthetic if desired, one has no anxiety with regard to the respiration. Artificial respiration can be intermitted and restarted in a moment. If it be found

possible to use this method in man—and I shall be convinced of the impossibility only when I see it fail—there would be a great increase in the percentage of recoveries after attempted resuscitation in man.

Other Factors requiring Attention in Resuscitation.

In experiments which I performed in the latter part of 1915 I found that there were certain other points which have an important influence on the speed and completeness of resuscitation, and which I may take this opportunity of pointing out.

In the first place, when natural respiration begins (after arrest of the heart, massage, etc.) the respirations to begin with are slow. When the tube through which artificial respiration has been performed is allowed to remain in the trachea they tend to continue slow. I found on looking over the data of experiments that when this tube was removed from the trachea there was invariably a sudden and pronounced increase in the rate of respiration. This, no doubt, was partly due to increase in the available lumen of the trachea consequent upon the removal of the tube. But it seemed greater than could be accounted for by that alone, so that it is possible that the presence of the tube, from reflex irritation of the air passage, tends to slow the natural respiration. Whatever the theoretical explanation the practical conclusion is clear: As soon as natural respiration has become established the tracheal tube should be removed.

In the second place, there is one danger point in the course of resuscitation which I believe is of great importance and which has not to my knowledge been noted before. After considerable experience in resuscitating animals killed by chloroform, I found that I could anticipate with some confidence that I would be able to resuscitate an animal, by the methods described, to the stage that the heart beats and natural respiration were resumed. But here an unexpected difficulty arose, because I found repeatedly, but not invariably, that after the first gasp or two of natural respiration the heart, hitherto beating well, suddenly ceased beating and could not be restored even by massage. I had this experience with three rabbits in succession in one day. This was distinctly disappointing. In casting about for an explanation the only one that suggested itself—and it is not entirely satisfactory—is this: After stoppage of the circulation for several minutes the central nervous system is paralysed. The centres recover when the circulation is resumed and the blood aerated. When the respiratory centre recovers, probably other medullary centres recover somewhere about the same time. It is conceivable that when the respiratory centre commences to function the vagus centre commences at the same time or only just later, and that after its prolonged abeyance of function there is a sudden and excessive discharge of vagal impulse which stops the heart. Acting on this hypothesis, whether it be correct or not, in subsequent experiments I gave an intravenous injection of atropine to prevent vagal inhibition, and after that I did not experience this type of cardiac failure. The result may, as so often happens, have been due to an increase in skill in performing the experiments of which I was not cognisant. But in any case, seeing that atropine can do no harm, I think it would be advisable provisionally to adopt this safeguard in resuscitation in man.

Lastly, in all resuscitation experiments great care must be taken to maintain the body temperature. After stoppage of the circulation and temporary paralysis of the whole central nervous system, the animal does not easily recover its power of heat regulation.

SUMMARY.

To sum up, I would venture to lay down the following directions for resuscitation in cardiac arrest in man or animals, whether from chloroform anaesthesia or from any other condition in which cardiac arrest is "accidental."

If the heart has stopped, artificial respiration should be resorted to immediately by the usual methods. If, at the end of three or four minutes, no pulse can be felt, and especially if no heart beats can be heard on auscultation, adrenalin should be injected into an external jugular vein, by an assistant if available; if no assistant be available, I believe the gain from adrenalin more than compensates for the loss of time, provided the materials are at hand, as they ought to be. In the meantime, or subsequently,

as the case may be, the abdomen should be opened high up, one hand inserted in the opening, passed up over the left lobe of the liver, and the heart felt for above the diaphragm. The heart should then be massaged intermittently. I endorse all that Mr. Fisher says in regard to not relying upon artificial respiration alone and in regard to boldness of procedure if the heart has stopped. When it has stopped there can be no further risk. As soon as spontaneous beats of the heart, however feeble, are felt massage should be interrupted for several beats to allow the contractions to establish themselves; otherwise, continuous massage may abolish them again. The heart should then be massaged again for a short time, when the beats will probably improve. When the heart has begun beating it is safer to use too little than too much massage. The hand should not be removed from the abdomen till the heart is beating well. When the heart continues to beat regularly and strongly an intravenous injection of atropine should be given. Artificial respiration must be kept up continuously, also after the heart begins beating. After five minutes it may be interrupted to see if natural respiration will begin, but not for long. No harm is done by keeping up artificial respiration, even if it inhibits the natural respirations, whereas, on the other hand, it is not safe to interrupt artificial respiration very long, if natural respiration does not commence.

If it be found that positive artificial respiration can be used in man by, for example, a Brodie's pump of larger size than that used in the physiological laboratory, the problem of resuscitation will be greatly facilitated.

Natural respiration may be expected to begin in five to ten minutes after the heart begins to beat again, but may be delayed for thirty minutes.

During the manipulations, and afterwards if these procedures be successful, care must be taken to maintain the body temperature.

All attempts, successful or not, at resuscitation after cardiac arrest in man should in the present state of knowledge be recorded. When it is possible—and it cannot well be possible unless assistance is available—accurate notes should be kept not merely of the sequence of events, but of the actual times of occurrence.

REFERENCES.

- ¹ Gunn and Martin, *Journ. of Pharmacol. and Exper. Therapeutics*, 1915, vii, p. 31. ² Gunn, *Proc. Roy. Soc.*, 1914, lxxxvii, p. 551. ³ Gunn and Underhill, *Quart. Journ. of Exper. Physiol.*, 1911, viii, p. 275. ⁴ Gunn and Whitelocke, *Brit. Journ. of Surg.*, 1914, ii, p. 92. ⁵ Gunn, *Quart. Journ. of Physiol.*, 1913, vii, p. 75.

THE PRACTICAL GAINS OF CLINICAL CARDIOLOGY SINCE 1900.*

BY

W. GORDON, M.A., M.D., F.R.C.P.,

PHYSICIAN TO THE ROYAL DEVON AND EXETER HOSPITAL; PHYSICIAN TO THE WEST OF ENGLAND EYE INFIRMARY.

DURING the last twenty years a great deal of interesting and valuable work has been done in medical cardiology. What I propose is briefly to summarize the practical gains which we, as clinicians, have derived from it. In so limiting my field, I am forced to exclude from it much that is interesting, as belonging to physiology rather than to medicine. But I hope that what my paper thus loses in attractiveness it may atone for by usefulness.

CARDIOLOGY IN 1900.

I need hardly remind an audience well read in the literature of twenty years ago that all the factors involved in cardiac failure were then clearly recognized; that no one called in question so obvious a truism as that the cardiac muscle was of predominant importance; or doubted the danger of such mechanical impediments as a leaking aortic orifice or a contracting mitral. No thoughtful person then referred to "back pressure" as a cause of heart failure, but only as a consequence and complication of it. Gaskell's brilliant work had shown that the heart muscle was more independent of nervous control than had previously been imagined, and Kent and His had demonstrated the path along which the normal stimuli of contraction travelled. The effects of vascular disease, in and

* Read at a meeting of the Devon and Exeter Medico-Chirurgical Society, to open a discussion, December 16th, 1920.