dietetic treatment, though of course there are exceptions. It is interesting to note that the intestinal symptoms seem to be much more beneficially influenced than the mouth symptoms.

As regards the mode of action of the intensive bicarbonate treatment, I have very few remarks to offer. At one time I believed that large doses of bicarbonate by decreasing the acidity of the intestinal contents might check the growth of monilia fungi, considered by some authors to be the true cause of the malady, and by Low and myself to be merely the cause of certain symptoms, as, for instance, the frothiness of the stools. If recent hypotheses concerning the etiology of the malady be proved correct, then a different explanation will have to be found. At any rate, there appears to be little doubt that in sprue a certain degree of acidosis is often present, and that an intensive alkaline treatment is indicated.

Conclusion.

The administration in sprue of massive doses of sodium bicarbonate by the mouth and intravenously, in conjunction with the usual dietetic measures, gives satisfactory results in a large number of cases, especially as regards the intestinal symptoms of the malady.

ANGINA PECTORIS WITH AURICULAR FIBRILLATION.

BY

HAMILTON M. SANDISON, M.B., CH.B. EDIN.

With a Note by FREDERICK W. PRICE, M.D., F.R.S. EDIN.

THAT auricular fibrillation may coexist with angina pectoris is demonstrated in the case of M., a woman aged 72. The value of digitalis when its administration is controlled, not by its empirical dose but by its therapeutic effect, was evident in the treatment of the case.

M. has suffered more or less all her life from cardiac affection, but, notwithstanding the advice of her physician, has for the past ten years led a strenuous life. On several occasions in recent years, she has been compelled by a feeling of general and cardiac weakness to rest for two or three weeks in bed, and to take 7 or 10 minims of tincture of digitalis thrice daily. She regained strength during these rests in bed, but the action of the heart remained rapid and irregular.

In July, 1920, she had a more serious breakdown, associated with breathlessness on exertion, palpitation, frontal and occipital headaches, and several paroxysmal attacks of pain of varying severity. This pain, undoubtedly that of true angina, was situated in the region of the heart; it was sudden in onset, and associated with a feeling of tightness in the front chest, mental perturbation, intense anxiety, and a sense of numbness in the left arm.

The pulse was completely irregular, with a ventricular rate of 124, many of the ventricular contractions not being registered at the wrist. The systolic blood pressure was undoubtedly raised, most of the beats registering 210 mm.; the wall of the brachial artery was thickened. The left border of the heart was one fingerbreadth outside the nipple line, the right border two fingerbreadths to the right of the sternum, and the note in the neighbourhood of the junction of the manubrium sterni and right intercostal cartilages and spaces was impaired. There was an apical systolic murmur, propagated to the axilla, and the second sound in the aortic area was rather accentuated. There was an absence of albumin in the urine.

Dr. Frederick W. Price, who was called to see the case, confirmed the diagnosis of auricular fibrillation and true angina pectoris. It is accordingly of special interest to find that in the chapter on auricular fibrillation in his valuable work *Diseases of the Heart*, he states: "It would appear, however, that patients suffering from auricular fibrillation are exempt from definite attacks of angina pectoris." Dealing with the etiology of auricular fibrillation, in the chapter referred to, two main groups of cases are described: "(1) Those with a history of rheumatism, and (2) patients suffering from myocardial degeneration. In the case of the former the condition occurs more commonly in early adult life, and the patients frequently have valvular disease—more often mitral than aortic disease, and more especially mitral stenosis. In the second group of cases the condition is more common after middle age." M.'s case probably belongs to the first group,

since there is a definite rheumatic history, and in view of the long duration of the illness and the presence of mitral regurgitation.

The etiology of the angina pectoris is, perhaps, not so certain; but the supernormal blood pressure, the evidence of thickening of the wall of the brachial artery, and the accentuation of the aortic second sound point to arterial degenerative changes, while the impairment of percussion note in the aortic area suggests the possibility of fusiform dilatation of the ascending aorta.

M. was confined to bed for four weeks, and then very gradually allowed to resume a more normal life. Only a very small quantity of fluid was allowed with each meal, and the total intake of fluid did not exceed 1½ pints a day. Tincture of digitalis was administered during the first week in 15-minim doses thrice daily. At the end of the week, as there was no appreciable improvement, the dose was increased to 18 and ultimately 20 minims. During the second week there was obvious improvement; the ventricular rate gradually falling to 84. As there was now some complaint of general depression, headache, and giddiness, the digitalis was stopped for forty-eight hours, and then recommenced in 15 minim doses. It was found possible to continue with this dose, and the ventricular rate ultimately diminished to 74, and the pulse became much less irregular. The systolic blood pressure came down to 180 mm., a drop of some 30 mm. Coincident with the remarkable slowing and steadying of the heart's action there was a most gratifying improvement in M.'s general condition.

Note by Dr. Price.—I desire to express my thanks to Dr. Hamilton Sandison for allowing me the opportunity of reading his notes of this case, which he kindly called me to see in consultation. The case was undoubtedly one of auricular fibrillation and angina pectoris. The first statement quoted by Dr. Sandison from my book was written towards the end of 1917, and represented my experience up to that time. Since then, however, I had observed two cases of auricular fibrillation (proved by electro-cardiographic examination) who also suffered from typical attacks of angina pectoris, before seeing the case of M., which made the third.

ANEURYSM OF EXTERNAL CAROTID ARTERY TREATED BY LIGATURE OF COMMON CAROTID ARTERY AND INTERNAL JUGULAR VEIN.

BY

W. A. PEVERLEY, AND J. K. HAWORTH,
M.B., B.S. DURH., MALVERN, NATAL. M.S., M.D. DURH.,

THE following case we think may be of interest and importance.

A Zulu woman, aged about 30 years, came with a tumour of the neck which she said came on after having had a tooth out. There were no signs of injury to her jaw, but she had a swelling the size of a Tangerine orange just below the angle of the lower jaw on the left side. The swelling was expansile in every direction and pulsated visibly. There was a loud systolic bruit over it. The impulse and the bruit disappeared on compressing the artery against Chassaignac's tubercle, and the swelling could be emptied on gentle pressure. It filled up from below on releasing the pressure. The pulsation in the left superficial temporal artery was delayed. The case seemed definitely to be one of aneurysm of the external carotid artery, the only other possibility being a pulsating tumour of the carotid body, and the subsequent history disposed of that possibility.

Following the principle of simultaneous ligature of artery and vein, one of us (Peverley) tied the common carotid artery and internal jugular vein at the seat of election. The after-course was uneventful; the pulsation at once disappeared, and now, six weeks after operation, the tumour is only the size of a small nut, and is hard and fibrous. She had no signs of any cerebral disturbance.

The artery seemed healthy, and she had no syphilitic history, though she had had two stillborn children. A Wassermann test was not done.

It does not appear to be the usual practice to tie the vein as well, though Ceci (Paris Surgical Congress, 1904) proposed the theory and Boari (*Policlinico*, 1905) has found it most successful. It is interesting to note in this regard that Lynn and Shipley (Ochsner, *General Surgery*, 1917) carefully avoided tying the internal jugular vein, thinking, presumably, it added to the risk of the operation. The idea of ligaturing the vein as well as the artery in this case was merely an enlargement of Sir George Makins's principle of simultaneous ligature of vein and artery in