neighbourhood of the wound, gastric distension, spasm and sometimes vomiting due to angulation of the pylorus and duodenum by adhesions.

To avoid these complications, to get rid of a potential source of recurrent infection in the biliary channels, it appears advisable to remove a gall bladder which has been attacked by cholecystitis of whatever type. The contra-indications are not local but general. In a small proportion of cases it may be advisable, in view of the patient's age or feebleness, to limit the operative procedure to the somewhat less exacting cholecystostomy after removal of calculi if present. With this exception, cholecystectomy should be the rule in acute, subacute and chronic cases. It is necessary to make certain that after its removal no obstruction will remain in hepatic and common ducts. If this is demonstrated beyond doubt, there is no need, even in cases where there is jaundice, to provide external drainage of one or other of these ducts; the indication for external drainage is an associated pancreatitis, and if this is present it is necessary to drain the common duct externally by rubber tube.

Patients who have had the gall bladder removed do not suffer any demonstrable disadvantage from its absence; in the case of calculous cholecystitis they are relieved of the double risk of the re-formation of stones and the persistence

of bile duct infection; in the case of cholecystitis without calculi they are relieved of a diverticulum which would in all probability prove the source of a recrudescence of their intoxication.

An operation field which permits a proper examination of the bile ducts is adequate for cholecystectomy. The operation may present considerable difficulties when the gall bladder is shrunken and lies embedded in adhesions—it is in this condition that its removal is most necessary if recurrent attacks of infection are to be avoided. Under other circumstances its removal is simple. The raw surface left can often be obliterated by suture, or sometimes an omental graft is useful for the purpose. If this is done, and the stump of the cystic duct touched with the cautery, a clean operation field is left in which adhesions are unlikely to form. The operation is altogether a cleaner performance than cholecystostomy in which the gall bladder is opened *in situ*, and some infection of the wound is difficult to avoid. If it is true that the gall bladder is of no particular functional value in the human subject, that it is the nidus of persistent infections in the biliary channels, that its removal is not a more serious operative risk than cholecystostomy, the argument for cholecystectomy as the normal operation in cholecystitis is overwhelming.

ASTHMA AND ANAPHYLAXIS.*

BY
FRANK COKE, F.R.C.S.

THE whole subject of anaphylaxis is of very recent origin; it is also very complex. It will therefore be well to have some general description of this strange phenomenon before passing to the claim that asthma be considered as a symptom of anaphylaxis or protein poisoning, and not otherwise a separate entity.

Charles Richet,¹ in 1902, injected a dog with a dose of poison derived from the tentacles of the actiniae and with no harmful effect whatever. Twenty-two days later he reinjected the dog with exactly the same dose, expecting to reach larger doses later and a condition of prophylaxis to the poison. To his surprise, hardly had he finished the injection when the animal became extremely ill and died in thirty-five minutes. This being the exact opposite of prophylaxis, he called it anaphylaxis.

Numerous experimenters in many countries have since investigated this question; detailed accounts may be found in the books of Richet,¹ Besredka² and Danysz.³

Apart from Richet's experiments, an explanation is found for many curious facts noted previously by Hyman⁴ and others, even by Magendie in 1839. It is found that any form of protein, serum, white of egg, milk, or cereal, has the power of rendering an animal sensitive to another dose of the same protein, which, when given, will instantly produce the anaphylactic syndrome—even to the point of death—in seconds. The next peculiar point is the absolutely specific nature of anaphylaxis. An animal sensitized with horse serum will not react to an injection of beef or any other serum except that from a horse. But no matter what kind of protein is used for the sensitizing and exciting dose, the anaphylaxis resulting is always the same.

The symptoms of anaphylaxis vary in the different animals. The typical picture, applicable even to man, appears as follows:

Provided a sensitizing dose has been given, and a sufficient incubation period of about ten days has elapsed, at the second

or exciting dose the animal is in trouble almost before the injection is finished. It becomes restless, scratches its muzzle as if irritated; the respiration is quickened, then difficult; vomiting takes place, especially in dogs; there is diarrhoea, with the passage of blood; micturition occurs; tetanic spasms and somersaults give place to paralysis and collapse; cessation of breathing brings death. There is marked eosinophilia and other blood changes.

There is at present no complete explanation of this remarkable syndrome. A diagram will serve to visualize the whole subject and to give a rough working explanation of many of the points in connexion with anaphylaxis.

The diamonds, A, B, C, D, represent varying quantities of antigen, horse serum, which are injected into the guinea-pig, E to F, below, such antigen being marked by horizontal lines. The body—that is, the blood and cells—is incapable of making use of this raw protein, or of eliminating it as such in the urine, nevertheless it disappears in the blood. Therefore it must undergo some change of a digestive nature inside the blood or cells. (There is no room for exceptions to rules in a short article.) The lower thin line, at F, denotes the potentiality of the blood to produce this internal digestive mechanism and, may be, this is in normal working order to deal with infinitesimal quantities

of undigested (by the bowel) proteins that pass through into the blood and cells from time to time or arrive there through the lungs. Under the stimulus of the protein injected (A), this mechanism (vertical lines) becomes active as it has never been before, and the protein disappears, G to H, until a time comes at the end of some ten days, H, when this digestion is finished and abruptly ceases. An excess of this digestive substance, J, then passes freely into the blood, whilst the further formation of

it sinks to normal, a mere potentiality, as shown by the thin line, K.

There is certainly a moment in the diagram, about H, possibly, too, in the real anaphylaxis it is meant to portray, when there must be antigen and antibody, protein and digestive substance, present together enough to cause a small anaphylaxis. It is at this moment when the serum rashes appear. Is this possibly their explanation?

The blood then contains the antibody or the toxogenin of Richet, the sensibilisin of Besredka, the precipitin of Friedberger, or more simply (here) a digestive substance. The animal is now sensitive. A further injection, B, of antigen horse serum meeting the antibody, J, in the correct proportion, and meeting it suddenly, will immediately cause the anaphylactic syndrome to take place, the large black bar, L. It is now apparent why no anaphylaxis occurs until the incubation period is over. If more antigen is injected during the incubation period, all that will occur is that the time when sensitization

*The substance of a lecture delivered to the Tunbridge Wells Division of the British Medical Association, November 25th, 1920.

appears will be postponed until the increased quantity of horse serum has been digested. But once sensitive, the animal will remain so for years, even for the rest of its life.

Apart from the various theories of the physico-chemical or biological upset that takes place in anaphylaxis, it is certain that the central nervous system plays an important part, for under an anaesthetic anaphylaxis will not develop, the animal waking up with the crisis safely passed and with the gain of some measure of immunity or rather desensitization.

No matter what protein is used in the experiments the resulting shock and all its symptoms are always the same.

Passive anaphylaxis. If the blood of the guinea-pig which is sensitive to horse serum, about J, be injected into another guinea-pig that fresh guinea-pig at once becomes sensitive to horse serum, and a dose of horse serum injected, at once without any incubation period, will immediately cause anaphylaxis. Further than this, if the guinea-pig's blood sensitized to horse serum be withdrawn and mixed in a flask with horse serum, the anaphylactic poison, whatever it is, is formed there *in vitro*, and if now injected into a new guinea-pig the latter is seized with anaphylaxis at once.

Anti-anaphylaxis of Besredka.² A certain sized dose of antigen is needed, and it must be injected quickly to cause anaphylaxis. Anti-anaphylaxis is the using up of this excess of digestive substance or antibody J and M by small doses C C, when the animal will become desensitized, N, and large doses, D, can then be given with impunity.

It will thus be seen that the horse serum is used in three different ways: as a sensitizing dose A, as a lethal exciting dose B, or as a curative desensitizing dose C C. In the desensitization by antigens of human beings who are variously sensitive to these antigens, great care is necessary to make certain that the size of the dose is suitable.

Such is a brief description of anaphylaxis, and it must be clearly understood that the diagram in no way enters into the controversy of how anaphylaxis really takes place; the digestive theory may be wrong from top to bottom, but in the meantime it does give a working explanation of the experimentally observed facts.

The Clinical Side of Anaphylaxis.

True anaphylaxis in man after a second injection of protein is rare, but many cases are on record.

A practitioner in India¹ gave himself a dose of antiplague serum, and in a year's time gave himself another. He was instantly seized with fainting fits, coma and asphyxia, and died in a few hours. With the frequency with which serums are now given more of such accidents may be expected. They may be avoided by small preliminary injections, C C, the anti-anaphylaxis of Besredka, by giving the injection very slowly,³ a special syringe—the Woodyatt pump—having been devised for this purpose, and by noting any sensitiveness by the skin reactions, as detailed below. If actually developed, anaphylactic shock may be treated by anaesthetizing the patient, whilst adrenalin or epinephrin will probably keep minor manifestations in abeyance.

ASTHMA AND ANAPHYLAXIS.

Apart from these very rare cases, there are a large number of complaints which have greater or lesser claims to be considered as anaphylactic symptoms of protein poisoning, and no longer as separate diseases. Chief amongst these is asthma. The case for asthma will be given very fully, and at various points of the argument other complaints that seem to come under the same headings will be noted. Anaphylaxis results from sensitization to foreign proteins. Half the cases of asthma can be proved to be due to sensitization by foreign proteins. In Allbutt's *System of Medicine*, asthma and hay fever are considered together in the same article. Hay fever is due to sensitiveness to pollen, not as a mechanical irritant, but from its protein content. An attack of hay fever can be excited in a susceptible patient by an injection of the pollen protein hypodermically and in mid-winter.⁴

Now the sensitiveness to the pollen can be tested for by placing pollen in the conjunctival sac. Test sets for the timothy grass are supplied from St. Mary's Hospital laboratories, where much of the work that led up to the American investigation was performed. Later, this method was transferred to a scratch on the skin, when an urticarial wheal can be produced. Further, it had long been noticed that some cases of asthma were undoubtedly due to the presence of animals—for instance, cats and horses. The next step was to see if a skin reaction could be obtained by the proteins of horse dandruff or cat hair, and the

hundred and one other proteins with which we come in daily contact. To cut a long story short, this method has proved eminently satisfactory and feasible, and by its means the cause of 50 per cent. of asthmatics can be determined,⁵ Dr. Freeman being the first to lecture on this subject in England.

Technique of the Skin Reaction.

The inside of the arm and forearm are cleaned, and a scratch $\frac{1}{2}$ in. long made with a scalpel, not going deep enough to draw blood. On to this the protein and solvent are placed. The tests should be an inch and a half apart. On the two arms a number of tests can be made at a sitting, though sometimes it will take three or even four hours to complete a test. A positive reaction consists of an urticarial wheal not less than $\frac{1}{2}$ in. across. This appears in from ten minutes to half an hour. Sometimes they are as large as a halfpenny with irregular branches coming out of them.

These reactions vary enormously in different people, so much so that some experience is necessary to judge the degree of reaction present. The proteins I use fall into four main groups: Some 70 foods, 20 bacterial proteins, 10 kinds of animal hair, and 10 pollens; also various face powders, etc. A full testing may take four sessions of an hour each.

Very rarely the history is clear enough to enable one to mark the offending protein with the first test, but much time is saved in the long run by a complete examination.

For instance, one patient could only give me a very short time. He was very sensitive to hen feathers. He gave up his feather pillows and feather bed, and his two eiderdowns, which he always used for warding off a chill and the asthma at night! Much of his asthma went, but not all. Further testing showed him sensitive to rice and beans, the rice undoubtedly being responsible for his last attack. He is now clear of asthma.

Multiple reactions are very common, as the following case will show.

A child 5 years old, with a strong family history of asthma; eczema and urticaria always; asthma continuously day and night for one year. The skin reactions were banana, cabbage, eggs, and rhubarb + + + +; barley, beans, rye, and oats + + +; beetroot, carrot, chicken, lentil, orange, potato, wheat, and rabbit fur + +; articles to which he was not sensitive were apples, beef, Bird's custard powder, Colman's cornflour, cocoa, codfish, coffee, haddock, herring, milk, lettuce, pear, pork, rusks, rice, lamb, tapioca, veal, horse dandruff, and hen feathers. On a diet chosen from articles to which he did not respond this boy has had no asthma since the first day he came to see me. It surprised me that he should be sensitive to cabbage, but I find that is quite a common thing to be sensitive to. His mother was perfectly certain that fish upset him, I was equally certain that it did not; he now takes large quantities of it. This child had been seen by many eminent medical men in London and Edinburgh both for his asthma and his eczema, but without the skin tests it is difficult to see how a correct dietary could ever have been drawn up.

An explanation is now given to the extraordinary likes and dislikes of these asthmatics in their food, their personal surroundings, and their locality.

A girl, aged 9, had asthma continuously for six years except for a break of eleven months whilst at Exmouth. She came back to London and the asthma recommenced at once, so she went back to Exmouth. There she had asthma this second time, rather worse than she had ever had it anywhere before. This is quite a common kind of history. I was asked to test her in St. George's. She gave horse + + + + +; dog + +; cat + + +. Her "animal history" was that she had always had a cat at home but none at Exmouth the first time. The second time she visited Exmouth there was a cat and a dog there which she played with all the time. She had then been in hospital six weeks with no asthma. She was given the hospital cat to nurse from 5 to 6 p.m. At 1 a.m. she woke up with a typically bad attack of asthma. This explains why so few have asthma in hospital; they leave the article they are sensitive to at home.

One more case. I found a girl very sensitive to dog hair and to nothing else. She got rid of the dog and her asthma went. Some months afterwards she went to the sea and had bad asthma the first night. In the old days she would have stopped there and had asthma for a few days, and then gone home because the "sea air" did not suit her. Having great faith, however, she made inquiries, and found that the people who had left that morning had four dogs, and all four had slept on the rug which covered her bed.

The absolutely specific nature of anaphylaxis is well copied by these asthmatics. Now 50 per cent. of all asthmatics and some 90 per cent. of asthmatic children give reactions. One of the most interesting reactions is that given to human hair, several married patients who are sensitive to it being convinced that they are best when sleeping apart from their husband or wife, as the case may be. It is undoubtedly "foreign"—that is, different from

the blood serum. Hen feathers, chicken meat and eggs, even the white and the yolk, are all different as proteins, and I have had patients sent to me who are only sensitive to one of these.

The following case illustrating passive anaphylaxis, published by Ramirez,⁹ is of great interest:

A man was transfused with 600 c.cm. of blood from a donor, with no immediate discomfort. A fortnight later, on going out for a drive, he was seized with violent asthma, never having had such a thing previously. It was found that the donor he was transfused from was an admitted horse asthmatic, sensitive to 1 in 50,000 horse dandruff protein. His sensitive blood at J in the diagram had been transferred to this new man and had sensitized him—to wit, a skin reaction to 1 in 20,000 of the protein.

No less interesting is the fact that another man who received even more blood from this same donor did not develop asthma. What was there in his blood that had avoided sensitization? Or was he sensitized but did not show it by asthma? Their blood groupings are not stated, but presumably they were the same.

Certain individuals are sensitive to other things than proteins.

I examined a patient who gets asthma at once on taking half a grain of aspirin. Another was recently unconscious for ten hours in an asthmatic seizure after being given a kindly night-cap of 5 grains of aspirin. In other people aspirin, quinine, and iodoform or antipyrin^{11,12} give rise to a profuse urticaria. If the serum of these people be injected into animals, the animals become sensitized to the drug. They become extremely ill when these drugs are then administered to them, whereas the controls give no symptoms. Pagniez¹⁰ states that, serum from asthmatics having been injected into guinea-pigs, asthmatic attacks can be produced in them by injecting them with a solution of the Charcot-Leyden crystals, which the French consider to be made of tyrosin.

These asthmatics show many other idiosyncrasies to food and other proteins quite comparable to the other symptoms of anaphylaxis. It appears that, given the entry of a foreign protein, it is a matter of luck or predisposition what symptoms are called forth.

The connexion between asthma, hay fever, and paroxysmal sneezing is well known. Any of them may be due to pollens, animals' hair or the dust of tea or flour. They simulate the nasal irritation of anaphylaxis. Sometimes a simple hoarseness takes the place of both. Vomiting is common.

One child, who is only sensitive to rabbit fur, vomits, as the father says, at the sight of a bowl of rabbit broth. Another patient, who reacts to a solution of 1 in 10,000 of white of egg, is sick even with the glazing on the top of a bun or with a spoonful of soup cleared with white of egg, an inconvenience that overtakes him in a few seconds. Another patient, who has not taken honey for thirty years because it makes him sick instantly, in spite of this interval still gives a +++ reaction to honey. Those patients who are sensitive to animal hairs must avoid circuses and the Zoo. I had a patient who had to be actually carried out of the monkey house. Another who went to a performance of the "Garden of Allah" only saw the first act.

Eczemas and urticarias are very common with asthma, but even alone they give much the same proportion of skin reactions¹³ and are, in fact, merely symptoms from the same protein cause. The ragweed pollen may cause eczema without any asthma or hay fever.¹⁴ A patient who gives a reaction to one member of a group will very often be sensitive to other members of the same group—that is, the hairs, cereals or legumens—probably because they contain some of the same proteins as those in the article tested. Epilepsy is common with asthma and migraines, and is also in the same category.

While the Americans have devoted their energies to the skin tests, the French go to prove all these complaints as anaphylactic by various blood changes.

Widal, Abrami,¹⁵ and their collaborators have discovered a "crise hémoclasique," as they term it, which takes place in anaphylactic shock, in protein shock, and also in asthma, urticaria, migraine, and epilepsy, in paroxysmal haemoglobinuria, in malaria and other parasitic complaints when large quantities of protein are thrown suddenly into the blood stream, and also in those absorbing proteins from their own wounds; as a side issue on this, the rapid removal of tourniquets, massage, and squeezing of the limb with bandages is contraindicated even in civilian practice.

The "hémoclasique crise initiale" consists of a fall in blood pressure, a leucopenia, the leucocyte count falling

from 12,000 to 4,000 or even less, and many other less easily observed changes in the coagulability of the blood, its refractive index, the colour and substance of its clot.

The "marchand de mouton"¹⁶ had dealt in sheep for nearly forty years, during the last three of which he had become so sensitive to the smell of sheep that he had to give up his trade. In hospital he, of course, had no asthma. For the experiment he entered a pen of sheep. His leucocyte count and blood pressure were taken every few minutes. After two hours both fell, the leucocyte from 12,000 to 6,000. They then rapidly returned to the normal. During this time he felt no difference in himself. At the end of three and a half hours, when the crisis was completely over, he developed a very bad attack of asthma.

Another man had migraine, especially after chocolate,¹⁷ but only at times. He had chocolate each morning in large quantities, followed each time by a normal digestive leucocytosis, until there came a morning when the blood pressure and leucocyte count both dropped. Shortly after this gastric disturbances and a bad attack of migraine came on.

In all cases the blood crisis passes without any knowledge of inconvenience to the patient, but, having passed, the asthma or migraine at once develops. With certain cases of paroxysmal haemoglobinuria all that is necessary to initiate the paroxysm is to place the patient's hands in a bowl of very cold water. The blood crisis follows at once, and later the haemoglobin appears in the urine.¹⁸ But because these blood changes are present in this variety of conditions and after the injections of proteins into the blood, it does not, of course, necessarily mean that they must all be true anaphylaxis. Besredka² says that naturally intravenous injection of large quantities of foreign protein, or what not, will give rise to a shock, but he refuses to call these various intravenous protein shocks anaphylactic, unless the same shock is caused when the substance is injected intrathecally.

Cowie and Colhoun,¹⁶ in their article on the blood changes in connexion with the non-specific protein therapy by means of injections of antityphoid vaccine, note this same initial leucopenia, etc., of the "crise hémoclasique" but differentiate this protein shock from that of anaphylaxis by the absence of an eosinophilia. This increase of the coarsely granular cells staining with eosin is always present in anaphylaxis. Of other complaints in which it is noted in man, the asthmatic attack is the one of all others in which it is most extensively found. The sputum may be packed with them, in the blood they may be half the number of white cells present. Eosinophilia may, then, be another link to bind these anaphylactoid conditions together. Meschoowitz¹⁷ notes that it is present in the subjects of asthma and hay fever, and in their secretions, in the urticarial wheals and bullae of a great many skin diseases, in mucous colitis, in pellagra, and after an injection of tuberculin, in tetany, hemiparesis and epilepsy. It is also a point of great diagnostic value in intestinal parasites. Note the common occurrence of epilepsy, urticaria, even asthma, with worms—that is, an absorption of foreign protein?

Now what of the 50 per cent. of cases in which we do not get a reaction? One is surprised to find how little interest is taken in those cases by the Americans. The type of asthma, excluding the cardiac or renal air hunger, is exactly the same as those in which we get reactions. Probably the fault lies in our not having a sufficient list of proteins to try them with. There are many different proteins in a grain of wheat. I have six in my tests. Also think of the number of protein derivatives there are in the digestion of foods, the scores of microbes inhabiting the bowel, the hundreds of non-pathogenic organisms, many unnamed and unknown to us, but each with its specific protein to which we may be sensitive. There is also the question of a period of desensitization after attacks to be considered.

Chronic Anaphylaxis.

Danysz, the French bacteriologist, has recently published a book which is creating considerable excitement amongst the few who have read his work on *Maladies Chroniques*.

He attributes to sensitization to intestinal microbes or other proteins, not only the attacks and exacerbations of chronic complaints from frequent anaphylactic upheavals, but states that the actual presence of antibody or our "digestive substance" in excess is the cause of most chronic complaints. Fresh antigen may then precipitate big attacks of asthma, urticaria, and so forth, or may by smaller doses cause such complaints as psoriasis, eczema, neurasthenic and menstrual disorders or stomachic upsets.

It is necessary always to bear in mind the double poisoning of microbes and other substances, one in virtue of their toxins, the other in anaphylactic response to their protein content.

The following experimental work backs up Danysz's claim:

Rabbits are given doses of horse serum daily for two or three months with no ill effect. In this way a maximum quantity of the digestive substance is formed. Now if the animals be left for three months without any injections it is found that some will have paralysis, others skin troubles with alopecia, others arthritis, in fact at the end of a year they will nearly all be dead. They waste and in every way copy the chronic complaints that man is heir to.

How do Asthmatics become Sensitive?

In many there is a strong family history, not to the actual protein but to asthma and like diseases. A horse asthmatic father may have one child with hay fever, another with asthma and sensitive to potatoes and hen feathers. There is often a history of fever or inflammation of mucous membranes which has allowed the sensitizing protein to pass through.

Schloss's boy¹⁹ had albumen water for diarrhoea when only a few days old, when egg albumen can pass through the intestinal mucous membrane; this was his sensitizing dose. He next had his first spoonful of egg, the exciting dose, at fourteen months, and it nearly killed him.

The long-continued presence of a protein may make people sensitive to it. I have a patient who has been a baker for twenty years; he is now an asthmatic, very sensitive to wheat and oats.

We will now go over a case of asthma in the light of this new work.

More important than the age is the age of onset. The younger the patient the more simple the cause, the freer from bronchitis and bacterial infections or sensitizations. Ninety per cent. of children give reactions. But no matter what age the patient may be, the chance of finding the cause by the skin reactions increases in proportion to his youthfulness at his first attack. Patients 60 years of age will give reactions if they started their asthma in childhood; whereas in those commencing after 40 very few give reactions except clear cases of occupational sensitization, as the butcher and the baker. The chief points in the history are to know if the complaint is continuous or arises only at certain times of the year or in certain places. If the asthma is constant, have there ever been times when the patient has been free—on a sea voyage, for instance?

The family history merely denotes the predisposition and will not help us to the protein at fault. The pre-natal history will not help either, but is intensely interesting. Like everything else in asthma, it is completely paradoxical.

One patient's mother had seven children. Her husband was a horse asthmatic. With one child, a boy, she had eczema all the time she was pregnant; that boy is a horse and dog asthmatic, all the rest of the children being normal. Another lady has had asthma ever since she was married, very badly and always, excepting for nine months when she was pregnant. The child, a boy, is healthy. I have had a similar history with a migraine patient. Another lady had only had hay fever twice in her life, on both occasions during the whole summer that she was pregnant. One of her sons is a horse asthmatic, the other has had hay fever. Another patient, sensitive to milk, had to have chloroform and morphine every day and night during her two pregnancies. She did not nurse either child. One child was very ill for the first six weeks with gastric troubles after milk; the other, on milk also, died from asthma at 6 weeks of age. This was probably true transmission of sensitiveness to the offspring, as can be produced experimentally in animals. It soon passes off, as it did in the first child.

A history must be made of any distastes for special foods, of the animals present in the house, of microbic diseases, or of the effect, as known, of pollens. Next to the skin reactions, histories of treatment are interesting when seen in bulk. Every article that can be a cause of the asthma must be tested for. Too much reliance must not be placed on the history. One child was very ill in the country during the haymaking season, but better at the seaside. He was not sensitive to pollens, but to oats, etc. He had much oatmeal in the country, and none at the seaside. A complete physical examination must be made; one must not miss tuberculosis even if the reactions give the cause of the asthma.

There is often a history of colds that are said to cause the asthma. More often than not the sneezing and the catarrh is true asthma; certainly, in children, the asthma is often preceded by a year or more of colds and bronchitis. Exercise and dust will often bring on an attack, but they are not the cause. If one finds the cause, the mother will write to say: "It is so nice to see little Tommy running about with the other children and not getting asthma." If the cause can be discovered and it is one that can be removed, the patient will lose the asthma from that moment. Conversely, if the asthma does not go, the cause or all the causes have not been discovered.

There is not space to discuss, nor do we know, the actual mechanism whereby these various proteins act, but there would seem to be two types of patients—those whose mucous membranes respond as quickly as do skin reactions, a matter of minutes, and others whose attack comes some hours after meeting the protein to which they are sensitive, as in my cat case and in the "marchand de mouton." Lastly we must not forget the remarkable part played by the nose in asthma. Occasionally the nasal surgeon will cure patients for varying lengths of time by simply cauterizing the nasal mucous membrane.

Whether the smell of animals actually contains protein which enters with the breath, or whether some central nervous mechanism is set in motion directly through the olfactory nerve is still a matter for investigation. Very little work has been done in connexion with the sense of smell, but the mechanism is extraordinarily sensitive. Now, normal people in the presence of a horse must breathe in just as much of the smell or protein of the horse as these asthmatics. Is it simply that their blood is not sensitive to the horse protein? Or is the protein stopped at the mucous membrane of normal persons? Or does it get through, but is summarily dealt with in normal persons? Again, can this possibly have any bearing on the facies that stamps the ostler and the groom so unmistakably?

Starting with one drop of a 1 in 100,000 solution, I have produced a bad attack of asthma by reaching an overdose at 5 minims of a 1 in 10,000 solution. If the patient is sensitive to articles which cannot be avoided, that is, chiefly horse-hair or pollen, he can be desensitized to either by gradually increasing but very small doses, always controlled by skin reactions. Another treatment is that by the intravenous administration of peptone,²⁰ on the idea that all these proteins are broken up into peptones during digestion, and therefore the one peptone may, so to speak, catch them all at a lower level than the specific proteins. The third method is also a method of anti-anaphylaxis, but instead of using small doses of peptone, as antigen, mixtures of microbes from the ordinary intestinal flora are used hypodermically. This is the method of Danysz. He uses autogenous and heterogenous cultures of all the intestinal microbes that will grow on a sloped gelatine or peptone culture. They are used as antigens, not for any toxins they contain, but simply for their protein contents.

By these various methods most cases can be improved, some made gloriously well. The delicate child cooped up in a stuffy nursery each winter with the cat, the dog, or the feather bed that is causing all the trouble, becomes normal straight away. It is impossible to tell which case will give reactions, or which will improve under any particular method of treatment. The method of Danysz must be used with very careful dosage or big attacks of asthma may come on, the patient even being made worse for a time.

With regard to the treatment of disorders allied to asthma, still scantier claims must be made because of the small number of cases coming under any particular heading. The following case is of interest:

A gentleman, at one time a fine athlete, has suffered from neurasthenia and gastric disturbances since the war, feeling blown out after his meals, and in a condition that might well be called hypochondriacal. On testing he is found to be very sensitive to potatoes. Removing these from his diet has made him a different man.

The treatment of Danysz in some cases does give the extraordinary results claimed for it, especially in chronic eczema and other skin diseases. I am, however, convinced that we have yet much to learn in the dosage of these entero-antigens.

If we look back at our diagram and see that the size of the dose alone decides whether it shall be a sensitizing dose,

an exciting dose, or a curative dose, we may well be puzzled in finding the correct dose in a sensitive patient.

Summary.

1. By the skin reactions half the cases of asthma can be proved to be due to some foreign protein.
2. If these causes can then be removed the patients are relieved of their asthma at once.
3. Much can be done to relieve or even cure other cases which do not give positive skin reactions, by the usage of various proteins, as antigens, to desensitize them with.
4. Asthma, hay fever, urticaria, eczema, migraines, epilepsy, and paroxysmal haemoglobinuria, have many common traits and will probably all be proved eventually to be symptoms of anaphylaxis and not separate diseases at all. This list will most likely be considerably enlarged.

In conclusion I wish to thank those many gentlemen whose patients I have tested and treated, but especially my friend Dr. Latham, through whose interest my attention and work have been so fully turned to this matter.

This new advance in medicine may be only a matter of inches, but it does allow us to see round a fresh corner and to obtain a view, and a direct view, of great possibilities in connexion with these chronic complaints, the cause and cure of which have so long baffled explanation.

REFERENCES.

- ¹ Richet, *Anaphylaxis*, translated by Murray Bligh. ² Boeredka, translated by Gloyne. ³ Danysz, *Maladies Chroniques*, 1920. ⁴ Hayman, *De Sang et des alterations autotomiques*, 1889. ⁵ Longcope, *International Congress*, London, August 11th, 1913. ⁶ Lewis, *Journ. Amer. Med. Assoc.*, February 1st, 1919. ⁷ Freeman, *Lancet*, September 16th, 1911. ⁸ Goodale, *Boston Med. and Surg. Journ.*, 1916. ⁹ Talbot, *N. Y. State Journ. of Med.*, 1916; Chandler Walker, *Journ. Med. Research*, 1917, and some sixteen other papers; Wodehouse, *Journ. of Immunology*, 1917, etc.; Freeman, *Lancet*, 1920. ¹⁰ Pagniez, *Presse Médicale*, February 23rd, 1916. ¹¹ Vidal, *Presse Médicale*, February 4th, 1920. ¹² Cordier, *Journ. de Lyon*, 1920, p. 531. ¹³ Sutton, *Journ. Amer. Med. Assoc.*, vol. 75, p. 1434. ¹⁴ *Presse Médicale*, 1914, p. 525. ¹⁵ Cowie and Colhoun, *Arch. Intern. Med.*, January, 1919. ¹⁶ Moscheowitz, *New York Med. Journ.*, January 7th, 1911. ¹⁷ *Semaine Médicale*, 1913, p. 613. ¹⁸ Schloss, *Journ. Dis. Children*, 1912, 341, iii. ¹⁹ Auld, *BRITISH MEDICAL JOURNAL*, 1917, i, p. 580. ²⁰ Pagniez, *Presse Médicale*, November 19th, 1919. ²¹ Vidal, *Presse Médicale*, July 11th, 1914.

CHRONIC RETENTION OF URINE ARISING FROM PROSTATIC DISORDER:

ITS CAUSE, AND TREATMENT BY THE PRACTITIONER WITHOUT OPERATION.

BY

HERBERT T. HERRING, M.B., B.S.,

LONDON.

CHRONIC retention of urine is most commonly produced by some disorder of the prostate gland, and among these troubles enlargement is that most frequently met with. Retention therefrom was generally attributed to a pure mechanical cause, that is, the urethra was blocked or plugged by the intervention of a mass of abnormal gland tissue, which was thrust into the orifice during micturition, thus preventing the escape of urine. Many mechanical theories, based on this principle, have been advanced. At one time it was a large middle lobe which dropped into the orifice like a ball valve. At another it was side-flaps springing from the neck, which met together in the middle line like flood-gates. At a third it was the regular enlargement of the glands round the orifice which prolonged the urethra as a solid projection into the vesical cavity and prevented the escape of fluid, much as an excise-man's inkstand does when upset. And finally, the post-prostatic bar was accused of causing the mischief, for it was said that the bar came forward over the outlet during vesical contraction and effectually blocked it, while at the same time the pressure produced a deep pocket or pouch behind the prostate. None of these explanations appear to be entirely satisfactory. For their supposed action is not supported by the facts disclosed during the use of a catheter; neither do they account for many of the symptoms frequently met with; and finally, the removal of the irregularities—valves, flaps, nipples, bars, etc.—does not invariably restore the lost function. The whole prostatic tissue must be enucleated to ensure success. It would seem, therefore, that although mechanical obstruction may assist, it is not the principal cause of retention.

Before giving yet another explanation, on a different line, a brief sketch of the various pressures existing in the bladder and urethra is necessary to make the argument clear.

Urine within the bladder, when the organ is quiescent, is always subjected to pressure—namely, that due to the force exerted by muscle tone, and to that of the weight of the viscera above. For convenience these two forces are considered as one, and may be termed "tonal bladder pressure." It is found that this tonal pressure increases as the bladder fills—thus, for instance, when empty the pressure may register 15 mm. of mercury, but when holding 10 ounces of urine it may have risen to 30 mm.

The escape of urine is prevented by the resistance the prostatic urethra offers to dilatation. The channel is always tightly closed, and is only open when the pressure on the urine in the bladder exceeds its power of resistance. It therefore follows that as urine does not escape until the act of micturition—that is to say, not until a muscular contraction is evoked—this prostatic resistance must always be greater than the maximum tonal pressure, and if the latter is known by experiment, then the pressure of prostatic resistance must be in excess of that figure. The resistance throughout the length of the prostatic urethra is not equally distributed: it is least at the commencement and increases gradually towards the apex of the gland, where it is the greatest. The total resistance of the whole passage may be said very roughly to be between 40 and 50 mm. of Hg, and it is possible that its maximum is attained with an empty bladder, decreasing slightly as the organ is distended.

When the rise of the tonal pressure, due to the gradual filling of the bladder, approaches that exercised by the prostatic resistance, urine begins to make its way into the first part of the prostatic urethra, which then becomes an integral part of the bladder. As the invasion advances the reflex centre controlling the muscular power of the bladder is stimulated, and the desire to micturate is induced. Should the call be inconvenient, extra resistance is immediately imposed on the urethra by the contraction of the various bands of voluntary muscle distributed around the passage and also by the compressor urethrae muscle. The augmented opposition to the pressure in the bladder, and possibly the expulsion of the urine which has invaded the passage, checks the escape, and the slight wave of contraction which has swept over the bladder wall slowly dies away and with it the desire to micturate. If, on the other hand, the call is complied with, the muscular bladder wall is thrown into contraction, the prostatic resistance is overcome and the urine issues in a steady stream. During the act the voluntary muscles, if desired, may be called into play, so that more resistance is interposed; the stream then lessens, and finally stops, and urine is voluntarily left behind in the bladder. The muscular expulsive power of the bladder is found to behave similarly to that of the tonal pressure, for it is strongest when the bladder is full, and falls as it empties, but it differs in that it ceases directly evacuation is complete. The expulsive force, for instance, with 10 oz. of urine in the bladder may record 180 mm. of Hg pressure, but with 5 oz. it has fallen to 50 mm. or less, and with an empty organ there is none, only that of the tonal pressure of 15 mm. being registered. Further, under normal conditions the whole power the vesical muscle is capable of developing is not invariably called forth, the amount generated being tempered to the work that has to be accomplished, and normally there is always a good margin of reserve power.

The act of micturition is therefore effected, first, by the steady rise of the tonal pressure as the bladder fills, which gradually forces urine into a portion of the prostatic urethra, there stimulating the area governing the bladder muscles; secondly, by the contraction of this bladder muscle, which develops sufficient power to overcome the prostatic resistance.

Normally the two forces—bladder expulsive power and prostatic resistance—are so balanced that they allow of the complete evacuation of the bladder and nothing more. If, however, the balance is disturbed, as is frequently the case in disease, disorder of the act of micturition must follow. Thus, if the prostatic resistance is increased, the expulsive power remaining the same, then a point will be reached during evacuation and before the bladder is empty